



RIA. Revista de Investigaciones Agropecuarias

ISSN: 0325-8718

ISSN: 1669-2314

revista.ria@inta.gob.ar

Instituto Nacional de Tecnología Agropecuaria

Argentina

Laclau, Pablo; Murillo, Natalia; Bértoli, B.; Vignolio, Osvaldo
Tolerance of pedunculate oak (*Quercus robur*) saplings to herbicides
RIA. Revista de Investigaciones Agropecuarias, vol. 46, no. 3, 2020, September-, pp. 387-396
Instituto Nacional de Tecnología Agropecuaria
Argentina

Available in: <https://www.redalyc.org/articulo.oa?id=86465821013>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Tolerance of pedunculate oak (*Quercus robur*) saplings to herbicides

LACLAU, P.¹; MURILLO, N.²; BÉRTOLI, B.³; VIGNOLIO, O.⁴

ABSTRACT

Spraying herbicides to reduce weed competition is a usual practice in agricultural regions. However, tree tending under agroforestry or forest plantations is hampered by the extensive use of these pesticides. Here, we assessed the survival, symptoms expression and growth of pedunculate oak (*Quercus robur*) saplings after spraying different herbicides at recommended doses in two seasons: winter and spring. Three independent herbicide trials were carried out on oak sapling plots (E_WINTER 2013, E_SPRING 2012, E_SPRING 2013) in Ayacucho, province of Buenos Aires, Argentina. Plants were kept under uniform conditions in a nursery for about three months. We alternatively tested the herbicides (doses in sub-index, cc.ha⁻¹ or g.ha⁻¹) flumioxazin (FLUM₁₀₀), glyphosate (GLIF_{2000, 4000, 6000}), metsulfuron-methyl (METS₈), dicamba (DIC₁₂₀), propaquizafop (PROP_{500, 1000}), flurochloridone (FLUO_{1000, 1500, 4000}), atrazine (ATR_{2000, 4000}), acetochlor (ACET₂₀₀₀), imazethapyr (IMAZE_{500, 1000}) and imazapyr (IMAZA_{33, 100}). At the end of each trial all plants survived, regardless of treatment, although the symptoms varied depending on the trial and product. Glyphosate associated with chlorosis, necrosis and deformation, propaquizafop with necrosis, and dicamba, flurochloridone, imazethapyr and imazapyr with chlorosis. ATR₂₀₀₀ and FLUO₁₅₀₀ treatments advanced the re-sprout phases. In terms of growth, saplings sprayed with IMAZE₁₀₀₀, IMAZA₁₀₀ and GLIF₆₀₀₀ had lower total dry weight than the control (CT) ($p < 0.05$), in E_SPRING 2012. In E_WINTER 2013 differences of dry weight were not significant, but in E_SPRING 2013, FLUO₁₅₀₀ sprayed saplings had a lower dry weight of roots than CT ($p < 0.05$). Total plant height and stem diameter presented similar trends. According to the whole set of symptoms assessed, the application of FLUM₁₀₀, ACET₂₀₀₀, ATR₂₀₀₀ or METS₈ would be compatible with oak saplings production. The low effect of herbicides in wintertime might be related to different ways of avoidance to herbicide penetration: root depth, the absence of foliage or to the protective structure of buds (perulae); in spring, to root depth, leaf cuticle thickness and hairiness, and to the (high) capacity of the oaks to re-sprout. Although the results obtained are promising for oak plantation in sites exposed to common herbicides, further research is needed to assess long term responses, or to repeated applications over time.

Keywords: weed control, agroforestry, symptoms, growth, survival.

RESUMEN

El uso de herbicidas en áreas agrícolas es una práctica usual para reducir la competencia de malezas. Sin embargo, la plantación de árboles en sistemas agroforestales o en macizos es amenazada por la aplicación extensiva de estos agroquímicos. En este estudio se evaluó supervivencia, síntomas y crecimiento inicial de plantines de roble pedunculado (*Quercus robur*) tratados con distintos herbicidas bajo las dosis usualmente recomendadas y en dos épocas del año: invierno y primavera. Con este fin se llevaron a cabo tres ensayos

¹Instituto Nacional de Tecnología Agropecuaria (INTA), Estación Experimental Agropecuaria (EEA), Balcarce, Agencia de Extensión Rural (AER) Tandil. Rodríguez 370, (7000), Tandil, Argentina. Correo electrónico: laclau.pablo@inta.gob.ar

²Instituto Nacional de Tecnología Agropecuaria (INTA), Estación Experimental Agropecuaria (EEA) Balcarce, Agencia de Extensión Rural Nicanor Otamendi, Urquiza 2213, (7603) Comandante Nicanor Otamendi, Argentina. Correo electrónico: murillo.natalia@inta.gob.ar

³Vivero Bértoli. Av. Dindart 1650, (7150), Ayacucho, Argentina. Correo electrónico: tanoaya@gmail.com

⁴Universidad Nacional de Mar del Plata (UNMdP), Facultad de Ciencias Agrarias, Unidad Integrada Balcarce (UIB), Ruta 226, km 71.5, (7620) Balcarce, Argentina. Correo electrónico: vignolio.osvaldo@inta.gob.ar

independientes sobre plantines de roble (*E_WINTER* 2013, *E_SPRING* 2012, *E_SPRING* 2013) en la localidad de Ayacucho, provincia de Buenos Aires, Argentina. Las plantas se mantuvieron en el vivero durante aproximadamente tres meses bajo condiciones uniformes. Se probaron alternativamente los herbicidas (dosis en subíndices, en cc.ha^{-1} o g.ha^{-1}): flumioxazin (*FLUM*₁₀₀), glifosato, (*GLIF*_{2000, 4000, 6000}), metsulfuronmetil (*METS*₈), dicamba (*DIC*₁₂₀), propaquizafop (*PROP*_{500, 1000}), flurocloridona (*FLUO*_{1000, 1500, 4000}), atrazina (*ATR*_{2000, 4000}), acetoclor (*ACET*₂₀₀₀), imazetapir (*IMAZE*_{500, 1000}) e imazapir (*IMAZA*_{33, 100}). Al finalizar los ensayos todas las plantas sobrevivieron, independientemente de los tratamientos, aunque los síntomas variaron según los ensayos y productos aplicados. El glifosato se asoció con clorosis, necrosis y deformación de las hojas, propaquizafop con necrosis, y dicamba, flurocloridona, imazetapir e imazapir con clorosis. Los tratamientos *ATR*₂₀₀₀ y *FLUO*₁₅₀₀ adelantaron las fases de rebrote. En cuanto al crecimiento, los plantines tratados con *IMAZE*₁₀₀₀, *IMAZA*₁₀₀ y *GLIF*₆₀₀₀ alcanzaron un menor peso seco total que el control (CT) ($p < 0.05$) en el ensayo *E_SPRING* 2012. En el ensayo *E_WINTER* 2013 no se observaron diferencias significativas en el peso seco, pero en *E_SPRING* 2013, con *FLUO*₁₅₀₀ los plantines alcanzaron menor peso seco de raíces que el control ($p < 0.05$). Las alturas totales y los diámetros de tallo presentaron tendencias similares. Según los síntomas observados en el total de los ensayos, *FLUM*₁₀₀, *ACET*₂₀₀₀, *ATR*₂₀₀₀ y *METS*₈ serían compatibles con los plantines de roble. Los modos de evadir la penetración de los herbicidas en el período invernal pueden relacionarse con la profundidad de las raíces, con la ausencia de hojas y con la protección de las yemas por pérulas; y en la primavera con la profundidad de las raíces, con el espesor de la cutícula y con la pilosidad foliar, y con la alta capacidad de rebrote de los robles. Si bien los resultados alcanzados son promisorios para la implantación de robles en sitios expuestos a herbicidas de uso corriente, es necesario conducir otros estudios para evaluar la reacción de las plantas a largo plazo, o sujetas a aplicaciones repetidas en el tiempo.

Palabras clave: control de malezas, sistemas agroforestales, síntomas, crecimiento, supervivencia.

INTRODUCTION

Weed competition is a major issue for native regrowth or planted forests establishment in temperate regions (Vasic *et al.*, 2014, 2013; Willoughby *et al.*, 2006; Dixon *et al.*, 2005) and their control require combined strategies (Vasic *et al.*, 2014, 2013; Vencill *et al.*, 2013; Devine and Harrington, 2010; Tu *et al.*, 2001; Jiménez and Cabezuelo, 1995). In pedunculate oak (*Quercus robur*) saplings, competition results in lower sprouts growth, plant dry weight, stem diameter, nitrogen concentration, plant water potential and stem conductance (Venegas and Palazuelos, 2009; Lof, 2000). These effects are similar to those caused by low soil water potential, while weeds also reduce light, temperature and the availability of nitrogen in the soil (Venegas and Palazuelos, 2009; Lof, 2000). In trials with different European oak trees and other temperate hardwoods, Cogliastro *et al.* (1990) found, after three growing seasons, that the growth of the trees was higher in treatments with herbicide applications or mechanical control of weeds, which was associated to a larger availability of foliar nitrogen and phosphorus.

The use of pre or post-emergence herbicides is a usual and effective management technique to reduce competition from weeds, particularly in temperate cropping regions (Rosales Robles and Esqueda Esquivel, 2008), and when it is possible, it is combined with mechanical removal (Vasic *et al.*, 2013; Vencill *et al.*, 2012; Jiménez and Cabezuelo,

1995). However, direct spraying or the drift of herbicides can significantly affect the survival or growth of forest saplings (Matschke and Macháčkova, 2002; Jiménez and Cabezuelo, 1995) of both natural forests and plantations. There is little information on the effects of herbicides on forest saplings (Matschke and Macháčkova, 2002). In the Argentine pampas and other major agricultural regions of the world, the incorporation of trees under agroforestry systems or in plantations close to crops is hampered by the extensive application of herbicides.

Some post-emergence herbicides, such as cycloxydim, clopyralid and metazachlor have been tested on seedlings of *Q. robur* and *Prunus avium* with 6-8 expanded leaves, and on *Fraxinus excelsior* and *Acer pseudoplatanus* with 2-4 expanded leaves (Willoughby *et al.*, 2006). Survival was not affected in any case, although metazachlor reduced growth after repeated applications. In a test with a variety of coniferous and broadleaved species, clopyralid alone or combined with graminicides (cycloxydim, fluazifop-P-butyl, propaquizafop) did not affect the survival or growth of saplings (Dixon *et al.*, 2005). Another essay, on one to three years old pedunculate oak naturally regenerated plants monitored throughout three growing seasons, showed phytotoxicity from tribenuron-methyl, but not from nicosulfuron or imazamox (Vasic *et al.*, 2014). In controlled trials on two-year-old oak saplings (*Q. rotundifolia*), Jiménez and Cabezuelo (1995) found (i) high toxicity of

systemic herbicides like glyphosate, triclopyr or glufosinate-ammonium and residual herbicides as pendimethalin and diuron, causing mortality four months later, (ii) temporary or mild phytotoxicity of contact and residual herbicides as thiazopyr, or oxyfluorfen, and (iii) absence or very mild symptoms with simazine, flupoxan, and fluzifop-P-butyl. The application of simazine presented significant differences in height growth compared to the remaining treatments.

On the other hand, Matschke and Macháčková (2002) found, within one year after 2-4-D and glyphosate applications, a reduction of the concentration of indole acetic acid and cytokinins in oak (*Q. robur*) leaves, particularly with glyphosate. These authors highlighted that some morphological and physiological effects can be expressed in the next season, due to damages to dormant apical meristems. Also, in a trial with different sulfonylureas and asulam, damages of oak saplings occurred with metsulfuron-methyl and asulam (Lawrie and Clay, 1993). Iliev *et al.* (2013) searched the effects of the application of oxyfluorfen, fluzifop-P-butyl, quizalofop-P-ethyl and fenoxaprop-P-ethyl in field tests on two-year-old saplings of *Q. cerris* during 100 days of the growing season. While survival was not significantly affected, oxyfluorfen caused less growth in height, while the remaining herbicides (from the arilofenoxi propionic family) induced similar or greater plant height.

In this study, we explored the survival, symptoms expression, and growth responses of pedunculate oak saplings to the application of recommended doses of different herbicides under two seasons conditions, late winter and late spring. Both seasons are related to the opportunity of application of herbicides on winter and summer crops in temperate areas of the pampas of Argentina.

MATERIALS AND METHODS

Three independent trials of tolerance of oak saplings to herbicides were conducted in Ayacucho, province of Buenos Aires, Argentina. Different products and dosage (treatments) of herbicides were tested on a single application (table 1). One trial was initiated in August, late winter (E_WINTER 2013), and the remaining two in November, late spring (E_SPRING 2012, E_SPRING 2013). The trials lasted 98, 111, and 90 days, respectively. These seasons coincide with pre-sprouting (late winter) or unfurled leaves (spring) phenological phases of the oak saplings (Díaz-Maroto *et al.*, 1993) and match with the opportunity of herbicide application for winter and summer grain crops in the pampas. One-year-old saplings raised in the nursery from local seeds were used. Plants were chosen randomly from the nursery among the best qualified (unique and straight stems with no forks, upper buds healthy and flawless stems) and transplanted to individual plastic pots with 8 drainage holes around the bottom, 20 cm high and 12 cm diameter (volume: 2.26 liter). The substrate was ground dark soil from an A horizon of a local Typic Argiudoll. Each treatment plot was constituted by lots of 15 (E_SPRING 2012) or 30 saplings (E_WINTER 2013, E_SPRING 2013). The herbicides were sprayed using a Giber H® manual sprayer 1.5 kg.cm⁻² previous pressure, and 3 kg.cm⁻² maximum pressure. Each application was done in a single pass over the plants arranged in a row, simulating a sprinkler of a field spraying machine. The doses were prepared with neutral water, in the range recommended by the manufacturer on the basis of 100 l.ha⁻¹ of the application broth. A precision scale and a graduated cylinder were used for the measurement of the products. Control plants were treated with neutral water only. Both lots of treated and control plants were labeled and distributed randomly in

TRIAL	E_WINTER 2013	E_SPRING 2012	E_SPRING 2013
PERIOD AUG-DEC 2013	PRE-SPROUT	UNFURLED LEAVES	
	NOV 2012-MAR 2013	NOV 2013-FEB 2014	
COMMON NAME/ commercial (ACRONYM)	TREATMENTS (products x dose)		
FLUMIOXAZIN / Sumisoya (FLUM)	FLUM ₁₀₀		
GLYPHOSATE / Roundup (GLIF)	GLIF ₄₀₀₀	GLIF ₂₀₀₀ , GLIF ₆₀₀₀	GLIF ₄₀₀₀
METSULFURON_METHYL/ MetsulfuronAgros(METS)	METS ₈		
DICAMBA/ Hunter Agros (DIC)	DIC ₁₂₀		
PROPAQUIZAFOP/ Agil (PROP)		PROP ₅₀₀ , PROP ₁₀₀₀	
FLUROCHLORIDONE/ Rainbow (FLUO)		FLUO ₁₀₀₀ , FLUO ₄₀₀₀	FLUO ₁₅₀₀
ATRAZINE / Mazina 50(ATR)		ATR ₂₀₀₀ , ATR ₄₀₀₀	ATR ₂₀₀₀
ACETOCHLOR / Acetoclor Helm (ACET)		ACET ₂₀₀₀ , ACET ₄₀₀₀	ACET ₂₀₀₀
IMAZETHAPYR / Pivot (IMAZE)		IMAZE ₅₀₀ , IMAZE ₁₀₀₀	
IMAZAPYR / Clearsol(IMAZA)		IMAZA ₃₃ , IMAZA ₁₀₀	

Table 1. Herbicides, treatments (product x dose) and time period of the trials. Treatments are identified by an abbreviation of the common name of the herbicides with a sub-index representing the dose of active product (cc.ha⁻¹, except for imazapyr and metsulfuron-methyl, in g.ha⁻¹).

an open sector of the nursery on a polyethylene mesh to prevent soil rooting. During the trial period, the plants were kept under uniform conditions of luminosity, undergoing regular irrigation by aspersion. The presence of the following symptoms of damage on the leaves of the upper third was visually monitored every 12-15 days: chlorosis (isolated), spread chlorosis, necrosis, outbreak burn, and leaf deformation. At the end of each trial, the total plant height and the collar diameter of the stem were measured in three (E_SPRING 2012) or nine (E_WINTER 2013, E_SPRING 2013) randomly picked plants per treatment. Also, the dry weight of these saplings was determined by oven drying (60°C, seven days) the foliage, stem and roots. Besides, in the last two trials the average phenological phase, total plant height, and stem diameter were recorded over time. Qualitative data were analyzed by hierarchical clustering and principal components analysis (PCA) in bi-plot graphics. Quantitative data were analyzed with ANOVA models considering the treatments as a source of variation of a completely randomized design. Phenology variation was analyzed assuming a divided plot design considering the treatment as main plot factor and the observation date as the subplot (repeated measures over time). Assumptions of normality and homoscedasticity were probed through Shapiro-Wilks and Levene tests. All analyses were performed by using version 3.0.3 of R® (R CoreTeam, 2014), and *lsmeans* package (Lenth, 2014) for multiple comparison tests.

RESULTS

Sapling survival and symptoms

Sapling survival at the end of each trial was 100%, regardless of treatment. However, the treatments induced different symptoms, like chlorosis (partial or spread), necrosis (total, in the edges or in spots), leaf deformation and burning. Fig. 1 shows dendrograms of hierarchical clustering of the treatments according to the observed symptoms in the 2nd - 3rd and the 8th - 9th weeks after the herbicide application. In E_SPRING 2012 (top graphs), 11 days after the treatments, the control (CT) clearly differed from ACET₄₀₀₀ and from the group comprising the remaining treatments (cophenetic correlation 0.863). However, 58 days after herbicide applications, the control saplings (CT) presented similarity with the saplings treated with acetochlor (both doses), flurochloridone and atrazine (both only in low doses) (cophenetic correlation 0.782). In E_WINTER 2013, 64 days after treatments (at the stage of unfurled leaves) CT grouped with FLUM₁₀₀ and METS₈ and apart from GLIF₄₀₀₀ and DIC₁₂₀ (cophenetic corr. 0.872). In E_SPRING 2013, after 18 days of herbicide application, CT, ATR₂₀₀₀ and ACET₂₀₀₀ formed a common group distinctive of GLIF₄₀₀₀ and FLUO₁₅₀₀ (cophenetic corr. 0.734) and after 62 days, this last treatment joined the GLIF₄₀₀₀ group (cophenetic corr. 0.631).

On the other hand, PCA with bi-plot graphics (fig. 2) shows the correspondence between treatments and types of symptoms observed in leaves and new shoots. Results of E_SPRING 2012 showed no alignment between CT and any of the observed symptoms for both dates. On day

58, the treatments PROP₁₀₀₀, ACET₂₀₀₀, ACET₄₀₀₀ and ATR₂₀₀₀ reversed the initial symptoms while others, such as GLIF₂₀₀₀, IMAZA₃₃ or IMAZA₁₀₀ showed symptoms not expressed before. In E_WINTER 2013, at day 64, CT saplings and those treated with FLUM₁₀₀ and METS₈ did not present symptoms in leaves. In E_SPRING 2013, CT, ATR₂₀₀₀ and ACET₂₀₀₀ showed no symptoms at day 18, although at day 64 CT associated with necrosis and chlorosis.

Phenology

When analyzing the effects of treatments on the phenological phases of the saplings over the time elapsed since herbicide application, an interaction between treatment and date was found in experiment E_WINTER 2013 ($p < 0.0001$) (fig.3a). Only in the last observation (day 98), there were significant differences between treatments ($p < 0.05$); the phenological phase of saplings sprayed with GLIF₄₀₀₀ and FLUM₁₀₀ was more advanced than that of CT saplings, while it was more delayed for those treated with METS₈. That date was associated with the change of the phase of unfurled leaves of the second regrowth to the start of the third regrowth (phases 6 and 7 of the y-axis of Fig.3). On the other hand, there were neither significant differences nor interactions between treatment and date in experiment E_SPRING 2013 ($p < 0.06$) (fig.3 b).

Dry weight

Table 2 shows values of dry weight found in each trial, and differences among treatments ($p < 0.05$, Tukey's test). In E_SPRING 2012 there were treatment effects for the dry weight of leaves (ANOVA; $p = 1.81 \times 10^{-7}$), stem ($p = 0.000125$), roots ($p = 0.00044$) and for total dry weight ($p = 2.02 \times 10^{-5}$). IMAZE₁₀₀₀ and GL₆₀₀₀ treated saplings presented the lowest dry weight of leaves, which were not significantly different from those treated with IMAZA₁₀₀, ATR₂₀₀₀, IMAZA₃₃, FLUO₄₀₀₀ and IMAZE₅₀₀. As for the dry weight of stem GLIF₆₀₀₀ and IMAZE₁₀₀₀ treatments showed the lowest values, with significant differences only with CT, ACET₄₀₀₀ and ATR₄₀₀₀. The lowest values of dry weight of roots were found in GLIF₆₀₀₀ and IMAZA₁₀₀ treatments, with significant differences only with CT, ACET₄₀₀₀, PROP₁₀₀₀ and ATR₄₀₀₀. For the dry weight of the whole plant, GLIF₆₀₀₀ presented the lowest value, but not significantly different from IMAZA₁₀₀, FLUO₄₀₀₀, IMAZE₁₀₀₀ and IMAZE₅₀₀. In E_WINTER 2013, no treatment effects were found, and as for E_SPRING 2013 only FLUO₁₅₀₀ showed significant effects in the dry weight of roots, presenting a lower value than CT ($p < 0.05$).

Diameter and height

In E_SPRING 2012 there were treatment effects for stem diameter ($p = 2.38 \times 10^{-8}$) and total plant height ($p = 1.3 \times 10^{-8}$). The plants treated with GLIF₆₀₀₀ and IMAZE₁₀₀₀ showed the lowest diameters, significantly different from CT. IMAZE₁₀₀₀, GLIF₆₀₀₀ and IMAZA₁₀₀ treatments presented the lowest total




Figure 1. Cluster analysis (Manhattan distance) for testing: E_SPRING 2012 (above), E_WINTER 2013 (center) and E_SPRING 2013 (below), 11-18 days (left) and 58-64 days after treatment (right), with a three group leveling.

plant height, significantly different from CT. The combined variable diameter² x height also showed treatment effects ($p=1.97 \times 10^{-6}$), and CT differed from IMAZE₁₀₀₀ and GLIF₆₀₀₀ treated plants, showing the highest value. There were no treatment effects in stem diameter or total plant height in E_WINTER 2013. In E_SPRING 2013, effects in total plant height were not found, but the stem diameter and the diameter² x height of CT and ATR₂₀₀₀ treated plants were

significantly higher than those of FLUO₁₅₀₀, GLIF₄₀₀₀ and ACET₂₀₀₀ treated plants.

DISCUSSION

Survival

All of the plants survived at the end of the trials, regardless of the treatment. The survival capacity of the saplings




Figure 2. PCA with bi-plot graphics of E_SPRING 2012(above), E_WINTER 2013 (center) and E_SPRING 2013 (below), 11-18 days (left) and 58-64 days (right). Every inner radius represents the variation axis of the observed symptoms.

could be attributed to factors such as the spatial avoidance of underground biomass, with saplings developing fine roots away from the action zone of the pre-emergence herbicides (i.e. atrazine, acetochlor), or temporary avoidance, in case of winter applications. In winter, the absorption of contact herbicides would be restricted by the absence of leaves and the lignification of the stems. For example, glyphosate does not penetrate through the woody bark (Tu *et al.*, 2001). Also, aerial ingress through the buds would be limited by the protection of these meristems by scales (Feucht, 1988). On the other hand, tolerance and resistance to herbicides (*sensu* Vencil *et al.*, 2013) in actively growing saplings with expanded leaves would be given both by the extensive regrowth ability of white oaks (Qi *et al.*, 2006; Collet *et al.*, 1997) and by the leaf cuticle barrier, with thick adaxial palisade parenchyma (Qi *et al.*, 2006). While other barriers to herbicide penetration, such as the density and type of trichomes, are lower in *Q. robur* than those of other white oaks as *Q. petraea* or *Q. pubescens*. Doupuey and Badeu (1993) reported the presence of hair-

iness in leaf lamina, veins and petioles. Foliar trichomes of this species would be simple, serial and solitary in both sides (Penas *et al.*, 1994).

Symptoms and phenology

The analysis (cluster and PCA) of the observed symptoms in the upper leaves and sprouts showed variations between treatments (fig.1). Glyphosate treatments (different doses) associated rather with chlorosis, necrosis and deformation, propaquizafop with necrosis, and dicamba, flurochloridone, imazethapyr, and imazapyr treatments with chlorosis (imazapyr also caused outbreak burning) (fig.2). FLUM₁₀₀, ACET₂₀₀₀, ATR₂₀₀₀ and METS₈ treatments grouped together with CT, thus they seem to be more compatible for oak saplings survival and growth than the rest of the evaluated herbicides (fig. 2).

On the other hand, in E_WINTER2013 the phenological curves differed from the control only in the last observa-




Figure 3. Phenological phases of saplings by treatment overtime (mean $\pm$ SE) of E_WINTER 2013 (a) and E_SPRING 2013 (b). Phases (y-axis) are: (1) leaves unfurled (first outbreak of the year), (2) upper swollen bud, (3) elongated green bud, (4) beginning of sprouting (leaf primordia of the second regrowth), (5) first unfurled leaf, full sized (second regrowth), (6) final sprouting of the upper third (all leaves developed), (7) swollen bud (third regrowth), (8) elongated green bud (third regrowth), (9) leaf primordia of the third regrowth.

tion date (fig.3); at that time, GLIF₄₀₀₀ and FLUM₁₀₀ treated plants advanced the third regrowth flush when compared to CT ($p=0.06$), while plants under METS₈ treatment remained in the previous stage (phase 6, all leaves of the second regrowth developed) (fig. 3, left). In the E_SPRING 2013 trial, the CT curve separated from the all the rest at phase 7 (swollen bud of third regrowth), and differences clearly aroused from the third flush of growth (phase 8) (fig. 3, right), showing premature aging of leaves caused by herbicides spraying. Both trials results suggest a phenological response to some herbicides at late flushing stages due to previous harm to foliage.

Dry weights, diameters and heights

No significant differences between treatments were found in dry weight, stem diameter or total plant height in the winter trial (E_WINTER 2013) (table 2), suggesting that the above mentioned late effects in foliar damage and phenology would have not affected early plant growth under any treatment. Instead, some treatments in any of the springtime trials somehow affected the growth of saplings (table 2), showing total or partial depression of growth in weight or size of plants treated with different products and doses (GLIF₆₀₀₀, GLIF₄₀₀₀, IMAZA₁₀₀, IMAZE₁₀₀₀, FLUO₁₅₀₀, ACET₂₀₀₀).

The phytotoxicity of glyphosate we found for *Q. robur* has also been reported for plants of the genus *Quercus* and other hardwoods in previous studies (Matschke and Mach-

ačkova, 2002; Jiménez and Cabezuelo, 1995). On the contrary, reported damage caused to oaks by metsulfuron (Lawrie and Clay 1993) was not observed in our winter trial. The effect of the herbicides we tested, grouped by their mechanism of action (Papa, 2007; Tu *et al.*, 2001; CIAT, 1982), was not consistent with results previously published. For example, among the inhibitors of lipid synthesis (Group I), it was reported that quizalofop-P, fluazifop-P or cycloxydim did not damage and even promoted the growth of lateral roots (Lawrie and Clay, 1993; Willowghby *et al.*, 2006) while in the present study propaquizafop, of the same pole group, caused leaf damage. Among the destroyers of cell membranes (Group II), all reported herbicides (metsulfuron-methyl, asulam, oxidazon, oxyfluorfen) expressed different levels of phytotoxicity (Jiménez and Cabezuelo, 1995; Lawrie and Clay, 1993) although here, flumioxazin and metsulfuron-methyl appeared to be compatible with the oaks.

CONCLUSIONS

Oak seedlings behaved tolerant to various agrochemicals during early growth stages, underpinning the possibility of herbicide management in alley crops or in traditional crops adjacent to tree plantations. The adverse effects observed here, after direct spraying of the saplings, would be smaller under field conditions following preventive techniques such as directed crop spraying, use of protective screens or anti-drift sprinklers (Leiva, 2015). On the other hand, most of

TREAT	DRY WEIGHT (g.pl-1)									HEIGHT (cm)	DIAMETER (mm)	D ² x HEIGHT (cm ³)			
	LEAVES		STEMS		ROOTS		TOTAL DW		R/S						
E_SPRING 2012															
CT	2.36	ab	5.67	ab	12.69	a	20.72	a	1.58	38.1	ab	10.6	a	42.8	abc
ACET ₂₀₀₀	2.84	a	5.30	abc	10.04	abc	18.18	ab	1.23	36.1	abc	9.3	abc	31.2	abcd
ACET ₄₀₀₀	2.62	a	7.09	a	10.95	ab	20.67	a	1.13	39.7	a	10.9	ab	47.2	a
PROP ₁₀₀₀	2.43	ab	4.43	abc	11.08	ab	17.94	ab	1.62	35.7	abcd	9.1	abc	29.6	abcd
PROP ₅₀₀	3.12	a	4.89	abc	10.05	abc	18.05	ab	1.25	33.9	abcd	8.6	abcd	25.1	abcd
ATR ₂₀₀₀	2.08	abc	5.18	abc	10.30	abc	17.56	abc	1.42	39.5	a	10.3	ab	41.9	abc
ATR ₄₀₀₀	2.58	a	6.14	ab	10.52	ab	19.24	a	1.21	37.6	abc	9.9	a	36.9	abc
IMAZA ₁₀₀	1.06	bc	3.42	bc	4.45	c	8.93	cd	1.00	29.1	cde	8.7	abcd	22.0	bcd
IMAZA ₃₃	1.96	abc	4.56	abc	10.19	abc	16.71	abc	1.56	32.5	abcde	9.7	ab	30.6	abcd
FLUO ₁₀₀₀	2.32	ab	4.77	abc	10.07	abc	17.16	abc	1.42	31.5	abcde	8.9	abc	25.0	abcd
FLUO ₄₀₀₀	1.87	abc	4.09	bc	9.82	abc	15.77	abcd	1.65	31.0	abcde	8.9	abc	24.6	bcd
GLIF ₂₀₀₀	3.21	a	4.92	abc	9.35	abc	17.49	abc	1.15	31.8	abcde	7.7	abcd	18.9	cd
GLIF ₆₀₀₀	0.88	c	2.42	c	4.36	c	7.66	d	1.32	25.5	e	6.7	d	11.4	d
IMAZE ₁₀₀₀	0.75	c	2.77	c	5.98	bc	9.49	bcd	1.70	27.3	de	7.4	cd	14.9	d
IMAZ _{E50} 0	2.18	abc	3.91	bc	9.72	abc	15.81	abcd	1.60	30.5	bcde	8.8	abcd	23.6	bcd
E_WINTER 2013															
CT	2.54	a	4.63	a	5.46	a	12.63	a	0.76	44.4	a	5.8	a	14.99	a
DIC ₁₂₀	2.44	a	4.93	a	5.19	a	12.55	a	0.70	45.5	a	5.9	a	15.89	a
GLIF ₄₀₀₀	2.53	a	4.96	a	6.27	a	13.76	a	0.84	45.7	a	5.9	a	15.69	a
METS ₈	1.84	a	4.34	a	5.29	a	11.47	a	0.86	41.8	a	6.2	a	16.21	a
FLUM ₁₀₀	2.42	a	4.96	a	5.07	a	12.45	a	0.69	46.3	a	6.0	a	16.40	a
E_SPRING 2013															
CT	2.83	a	6.62	a	8.92	a	18.37	a	0.94	46.9	a	6.7	a	22.7	ab
ACET ₂₀₀₀	2.70	a	6.14	a	7.87	ab	16.71	a	0.89	47	a	6.2	b	19.29	bcd
ATR ₂₀₀₀	3.16	a	6.70	a	8.73	ab	18.59	a	0.89	48.9	a	6.9	a	24.71	a
FLUO ₁₅₀₀	1.92	a	5.08	a	5.49	b	12.49	a	0.78	45.3	a	6.2	b	18.9	cd
GLIF ₄₀₀₀	2.21	a	6.26	a	6.38	ab	14.84	a	0.75	46.1	a	6.2	b	18.9	cd

Table 2. Mean dry weight (g.pl⁻¹) of leaves, stems, roots and total, root / shoot ratio, height (cm), stem diameter (mm), and diameter² x height (cm³) by treatment of oak saplings at the end of each test. Same letters show no significant differences (Tukey test p<0.05) in each variable (columns).

the responses were not distinctively clear, and even with the experimental control and randomization imposed, other (genetic and environmental) factors (i.e. leaf insect attack) could influence results.

From all treatments, FLUM₁₀₀, ACET₂₀₀₀, ATR₂₀₀₀ and METS₈ behaved suitable to be applied to nearby crops, while IMAZA₁₀₀, IMAZE₁₀₀₀, GLIF₄₀₀₀ and GLIF₆₀₀₀ would not be advisable due to harm risks to the oak seedlings.

The observed differences on herbicides effect on growth between winter and springtime applications also highlight the importance of the opportunity of spraying herbicides. Winter applications would not significantly affect the growth of the oak saplings. The development of roots in-depth,

growing away from the action radio of preemergence herbicides (a few centimeters under the ground, i.e. atrazine, acetochlor), the absence of leaves during the period of activity of contact or systemic herbicides (i.e. metsulfuron-methyl, imazethapyr, dicamba), or bud protection by lignified scales, might be some of the mechanisms that prevent phytotoxicity. During that season, flumioxazin and metsulfuron-methyl herbicides did not cause symptoms or loss of growth. Neither dicamba nor glyphosate affected plant growth, although in the last this case the saplings showed symptoms of phytotoxicity. On the contrary, post-emergence herbicides imazethapyr, glyphosate, propaquizafop, imazapyr and flurochloridone sprayed in spring time caused negative effects on the dry weight, plant height or stem diameter of

the oak saplings. All these herbicides caused visible leaf symptoms and some anticipated regrowth flushes. In the same sense, atrazine (pre-emergent) would have not affected the saplings, which under the usual nursery practice develop roots of at least 10 cm in depth, much deeper than the activity zone of the herbicide, of about 5 cm (Devine and Harrington, 2010; Aldhous, 1975).

Although the results achieved are highly promising for oak plantation in environments exposed to some commonly used herbicides, further studies are needed to assess the reaction of treated plants in the long term and (or) after repeated applications.

ACKNOWLEDGEMENTS

This study was financed by the Unidad para el Cambio Rural, Proyecto de Investigación Aplicada (PIA) 10002 and by the Instituto Nacional de Tecnología Agropecuaria (INTA), Project PNFOR 1104075. The authors wish to thank Fernanda Buckley, Diego Domínguez Daguer and Luciano Lorea for the care and monitoring of trials. Also we acknowledge the observations and suggestions of the two anonymous reviewers, which deeply contributed to improving the quality of the article.

REFERENCES

- ALDHOUS, J.R. 1975. Nursery Practice. Forestry Commission Bulletin N° 43. 184 p.
- CEDERGREEN, N. 2008. Is the growth stimulation by low doses of glyphosate sustained over time? *Environmental Pollution* 156, 1099-1104.
- CIAT. 1982. Los herbicidas: Modo de actuar y síntomas de toxicidad. Centro Internacional de Agricultura Tropical, Serie 04SW-01-04, segunda edición, 40 p.
- COGLIASTRO, A.; GAGNON, D.; CODERRE, D.; BHÉREUR, P. 1990. Response of seven hardwood tree species to herbicide, rototilling, and legume cover at two southern Quebec plantation sites. *Canadian Journal of Forest Research*, 20(8), 1172-1182.
- COLLET, C.; COLIN, F.; BERNIER, F. 1997. Height growth, shoot elongation and branch development of young *Quercus petraea* grown under different levels of resource availability. *Ann. Sci. For.*, 54:65-81.
- DEVINE, W.D.; HARRINGTON, C.A. 2010. Planting native oak in the Pacific Northwest. Gen. Tech. Rep. PNW-GTR-804. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 25 p.
- DÍAZ-MAROTO, I.J.; PRUNELL TUDURI, A.; GONZÁLEZ HERNÁNDEZ, M.P.; SILVA-PANDO, F.J. 1993. Estudio fenológico de especies arbustivas y arbóreas de los robledales atlánticos. *Actas Congreso Forestal Español, Lourizán. Ponencias y comunicaciones*. T.1: 239-244.
- DIXON, F.L.; CLAY, D.V.; WILLOUGHBY, I. 2005. The tolerance of young trees to applications of clopyralid alone and in mixture with foliar-acting herbicides. *Forestry* Vol. 78, N.º 2.12 p. DOI:10.1093/forestry/cpi031
- DUPOUEY, J.L.; BADEAU, V. 1993. Morphological variability of oaks (*Quercus robur* L., *Quercus petraea* (Matt) Liebl, *Quercus pubescens* Willd) in northeastern France: preliminary results. *Annales des sciences forestières*, 1993, 50 (Suppl 1): 35-40.
- FEUCHT, J.R. 1988. Herbicide injuries to trees. Symptoms and solutions. *Journal of Arboriculture* 14(9): 215-219.
- ILIEV, N.; DANCHEVA, D.; ILIEV I. 2013. Effect of selective herbicides in the production of Turkey Oak saplings (*Quercus cerris* L.). *Muzeul Olteniei Craiova. Oltenia. Studii si comunicari. Stiintele Naturii*. Tom. 29 (2):97-100.
- JIMÉNEZ, M.J.; CABEZUELO, P. 1995. Evaluación de la fitotoxicidad de herbicidas sobre plantones de *Quercus rotundifolia* Lam. (encina). Congreso 1995 de la Sociedad Española de Malherbología: 225-228.
- LAWRIE, J.; CLAY, D.V. 1993. Tolerance of some Broadleaved and Coniferous Forestry Tree Species to Herbicides with Potential for Bracken Control. *Oxford Journals Science & Mathematics Forestry*, Volume 67, Issue 3: 237-244.
- LEIVA, P.D. 2015. Recaudos por deriva en pulverizaciones agrícolas. Siete aspectos importantes para su manejo. INTA, Estación Experimental Agropecuaria Pergamino. Pergamino (BA), 5 p. (Available at: http://inta.gov.ar/sites/default/files/script-tmp-inta_pergamino_recaudos_por_deriva_en_pulverizaciones.pdf verified: May 5th 2019).
- LEIVA, P.D. 2010. Consideraciones generales sobre calidad de agua para pulverización agrícola. Grupo Protección Vegetal - INTA, Estación Experimental Agropecuaria Pergamino. Pergamino (BA). 6 p. (Available at: http://www.inta.gov.ar/pergamino/info/documentos/2010/Calidad_Aguas_para_pulverizacion.pdf verified: January 24th 2016).
- LENTH, R.V. 2014. Lsmeans: Least-Squares Means. R package version 2.00-1. (Available at: <http://CRAN.R-project.org/package=lsmeans> verified: May 10th 2019).
- LÖF, M. 2000. Establishment and growth in saplings of *Fagus sylvatica* and *Quercus robur*: influence of interference from herbaceous vegetation. *Canadian Journal of Forest Research*, 30(6): 855-864.
- MATSCHKE, J.; MACHÁČKOVÁ, I. 2002. Changes in the Content of Indole-3-Acetic Acid and Cytokinins in Spruce, Fir and Oak Trees after Herbicide Treatment. *Biologia Plantarum*, Volume 45, Issue 3. 375-382 pp.
- PAPA, J.C. 2007. El modo de acción de los herbicidas *Actas Jornadas de Actualización: Elementos fundamentales para el buen uso de fitoterápicos. Dosis, Modo de Acción y Prevención de Deriva*. Sociedad Rural de Tucumán 17-19.
- PENAS, A.; LLAMAS, F.; PÉREZ MORALES, C.; ACEDO, C. 1994. Aportaciones al conocimiento del género *Quercus* en la Cordillera Cantábrica. I. Tricomas foliares de las especies caducifolias. *Lagascalia* 17 (2): 311-324.
- QI, Y.; FAVORITE, J.; CHIN, K.L.; XIAO, Y. 2006. Physiological, anatomical, and ecological characteristics of southern live oak. In: CONNOR, K.F. (ed.). *Proceedings of the 13th biennial southern silvicultural research conference*. Gen. Tech. Rep. SRS-92. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station 640. 448-453 pp.
- R CORE TEAM 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. (Available at: <http://www.R-project.org/> verified: May 10th 2019).
- ROSALES ROBLES, E.; ESQUEDA ESQUIVEL, V. 2006. Clasificación y uso de los herbicidas por su modo de acción. Tamaulipas, Mexico. 16 p. (Available at: https://www.researchgate.net/publication/266229319_CLASIFICACION_Y_USO_DE_LOS_HERBICIDAS_POR_SU_MODO_DE_ACCION verified: May 5th, 2019).
- TU, M.; HURD, C.; RANDALL, M. 2001. Weed Control Methods Handbook: Tools and Techniques for Use in Natural Areas. The Nature Conservancy. (Available at: <http://tncweeds.ucdavis.edu>, verified: April 2001).
- VASIC, V.; KONSTANTINOVIC, B.; ORLOVIC, S. 2012. Weeds in Forestry and Possibilities of their Control, Weed Control, Dr Andrew PRICE (Ed.), InTech. DOI: 10.5772/34792

VASIC, V.; KONSTANTINOVIC, B.; ORLOVIC, S. 2014. Application of post-emergence herbicides in the regeneration of pedunculate oak (*Quercus robur* L.) forests. *Forestry* Vol. 87, *Forestry* 2014; 0, 1–9. DOI:10.1093/forestry/cpt060

VELINI, E.D.; TRINIDADE, M.L.B.; BARBERIS, L.R.M.; DUKE, S.O. 2010. Growth Regulation and Other Secondary Effects of Herbicides. *Weed Science* 58 (3): 351-354. DOI: <http://dx.doi.org/10.1614/WS-D-09-00028.1>

VENCILL, W.K.; NICHOLS, R.L.; WEBSTER, T.M.; SOTERES, J.K.; MALLORY-SMITH, C.; BURGOS, N.R.; JOHNSON, W.G.; MCCLELLAND M. R. 2012. Herbicide Resistance: Toward an Un-

derstanding of Resistance Development and the Impact of Herbicide-Resistant Crops. *Weed Science* 2012, Special Issue: 2-30.

VENEGAS, R.; PALAZUELOS, R. 1999. Control químico de malezas arbustivas y arbóreas en plantaciones de *Pinus radiata* D. Don. Bushy and arboreal weed control with herbicides in *Pinus radiata* D. Don plantations. *Bosque*, 20(1): 79-88.

WILLOUGHBY, I.; JINKS, R.L.; STOKES, V. 2006. The tolerance of newly emerged broadleaved tree saplings to the herbicides clopyralid, cycloxydim and metazachlor. *Oxford Journals Science & Mathematics, Forestry* Volume 79, Issue 5: 599-608.