



Bragantia

ISSN: 0006-8705

ISSN: 1678-4499

Instituto Agronômico de Campinas

Brunetto, Gustavo; Benedet, Lucas; Ambrosini, Vítor Gabriel; Comin, Jucinei Jose;
Melo, George Wellington Bastos de; Santos, Matheus Ademir dos; Lourenzi, Cledimar
Rogério; Loss, Arcangelo; Belli, Paulo; Schmitt, Djalma Eugênio; Couto, Rafael
Copper and zinc fractions in the profile of an Inceptisol cultivated with apple in southern Brazil
Bragantia, vol. 77, no. 2, 2018, pp. 1-15
Instituto Agronômico de Campinas

DOI: 10.1590/1678-4499.2017033

Available in: <http://www.redalyc.org/articulo.oa?id=90859245013>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org



Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Copper and zinc fractions in the profile of an Inceptisol cultivated with apple in southern Brazil

Gustavo Brunetto^{1*}, Lucas Benedet², Vítor Gabriel Ambrosini², Jucinei Jose Comin², George Wellington Bastos de Melo³, Matheus Ademir dos Santos², Cledimar Rogério Lourenzi², Arcangelo Loss², Paulo Belli Filho², Djalma Eugênio Schmitt², Rafael Couto¹

1.Universidade Federal de Santa Maria - Departamento de Solos - Santa Maria (RS), Brazil.

2.Universidade Federal de Santa Catarina - Centro de Ciências Rurais - Florianópolis (SC), Brazil.

3.Empresa Brasileira de Pesquisa Agropecuária - Uva e Vinho - Bento Gonçalves (RS), Brazil.

ABSTRACT: The present study aimed to evaluate the accumulation and distribution of Cu and Zn fractions in the profile of an Inceptisol by application of fungicides with a previous history of apple cultivation in southern Brazil. Stratified soil samples (0.0 – 0.025 m, 0.025 – 0.050 m, 0.05 – 0.10 m, 0.10 – 0.15 m, 0.15 – 0.20 m and 0.20 – 0.40 m depth) were collected in two apple orchards (8 years old and 18 years old), and in a native pasture area adjacent to the orchards. The soil was subjected to analysis of physical and chemical properties and to chemical fractionation of Cu and Zn. The fractions of soluble, exchangeable, organic, residual and total Cu and Zn accumulated

in the topsoil (0.0 – 0.025 m) of the orchards, being higher in the orchard cultivated for 18 years. Most Cu, especially in the topsoil of the 18 years old orchard, accumulated in the less stable organic fraction, which can potentiate toxicity to plants and transfer Cu e Zn to ground water. In deeper layers, Cu was also distributed in the residual and mineral fractions, which are more stable. Most Zn in the soil of the apple orchards was distributed in the mineral and residual fractions, which have low mobility and low potential toxicity to plants.

Key words: *Malus domestica*, trace elements, toxicity, environmental contamination.

*Corresponding author: brunetto.gustavo@gmail.com

Received: Jan. 31, 2017 – Accepted: Sept. 18, 2017



INTRODUCTION

In the state of Santa Catarina, in southern Brazil, an area of approximately 18.000 ha is cultivated with apple crops (*Malus domestica*). It is the largest apple-producing area in Brazil, and Santa Catarina is the second largest producer, after Rio Grande do Sul state. Current apple production amounts to 531.000 t, equivalent to 44% of national production (Epagri 2013; Acate 2014). Nevertheless, apple trees are grown in regions with frequent rainfall, which stimulates the incidence of foliar fungal diseases and may reduce fruit yield and harm fruit quality. Therefore, foliar fungicides, e.g., the copper-based Bordeaux mixture ($\text{Ca(OH)}_2 + \text{CuSO}_4$) and zinc-based mixtures, such as those with the active ingredient Mancozeb ($\text{C}_4\text{H}_6\text{MnN}_2\text{S}_4 \times \text{Zn}$), are applied throughout the year. Furthermore, fungicides containing Cu and Zn are applied after leaf senescence and during plant dormancy to promote dormancy breaking and branches disinfection.

The accumulation of Cu and Zn fractions, especially in the upper soil layers, can be stimulated by several factors: unidirectional fungicide applications; Zn and Cu being washed off the leaf surface by rainfall; the deposition of senescent leaves and pruned branches; the cycling of Cu and Zn by species of ground cover plants that co-inhabit the orchards; a low transfer of Zn and Cu to apples; and an absence of soil disturbance between orchard rows (Fernández-Calviño et al. 2010; Mackie et al. 2012; Ruyters et al. 2013).

In soil, Cu and Zn are retained by physical and chemical bonds, by mineral and organic components, and their availability depends on: the mineral contents, oxides and hydroxides of Fe, Al and Mn; carbonates (Bradl 2004); soil pH values (Toselli et al. 2009); cation exchange capacity (CEC); content; and composition of organic matter (Fernández-Calviño et al. 2010). However, when Cu and Zn contents increase in the soil, as a result, for example, of frequent fungicide applications, there can be accumulation and changes in the distribution of their fractions in such soil components. Adsorption of Cu and Zn in the soil occurs primarily in high affinity sites, usually forming inner sphere complexes. The remainder is adsorbed at sites with low binding energy and, consequently, it increases the availability and mobility of these elements in the soil profile (Brunetto et al. 2014). Thus, the increase of Cu and Zn in the soil over the years may increase the amount of soluble and exchangeable fractions of these elements, increasing toxicity to plants (Michaud et al. 2007; Nagajyoti et al. 2010; Yadav 2010; Fernández-Calviño et al.

2012) and water contamination through their transfer in the soil profile (Mileusnić et al. 2014; Marchand et al. 2014).

The fractions of Cu and Zn can be estimated by chemical fractionation, which removes Cu and Zn from the most labile to the most stable fractions (Tessier et al. 1979). The total amount of these elements in soils or sediments can be separated into bioavailable fractions (water-soluble and exchangeable), potentially bioavailable fractions (bound to clay minerals, oxides, carbonates and organic matter) and residual fractions (bound to the structure of minerals) (Casali et al. 2008; Frankowski et al. 2010; Brunetto et al. 2013; Duplay et al. 2014). However, in the Americas, particularly in subtropical and temperate regions, few studies exist on the accumulation and distribution of Cu and Zn fractions in the soil profile in apple orchards with a history of continuous fungicides application, whose compositions include Cu and Zn. The present study aimed to evaluate the accumulation and distribution of Cu and Zn fractions in the profile of an Inceptisol on two plantations previously cultivated with apple trees in southern Brazil.

MATERIALS AND METHODS

Study location

This study was conducted using soil collected from two commercial orchards, one with eight years of cultivation (O8) and another with 18 years of cultivation (O18), and a soil from native pasture (NP), located nearby the orchards, in the county of Urubici, in the Planalto Serrano region, state of Santa Catarina, southern Brazil (Long 49°35'30"W, Lat 28°00'05"S). At the three sites, the soil was classified as an Inceptisol, a Cambissolo Húmico in Brazilian classification (Embrapa 2013). The local relief was classified as slightly undulating, with a 4% slope. The climate is humid mesothermal with mild summers, Cfb type according to Köppen classification, with an average annual rainfall of 1360 – 1600 mm, average temperature of 15.4 °C, and between 20 and 29 frosts can occur throughout the year (Pandolfo et al. 2002).

Orchard O8 was implanted in 2003 with a planting density of 4,444 plants·ha⁻¹ (4.5 m between rows and 0.5 m between plants), and orchard O18 in 1993, with a planting density of 1,482 plants·ha⁻¹ (4.5 m between rows and 1.5 m between plants). The orchards had two



commercial varieties, with Gala and Fuji accounting for 70% and 30% of the plants, respectively, with trees grown under a central-leader planting system. The plants of orchard O8 were grafted onto a Marubakaido rootstock, with a 20-cm M9 filter, while in orchard O18 the plants were grafted onto a Marubakaido rootstock without filter.

The implantation of the orchards was preceded by an application of lime to raise the soil pH in water to 6.0. Lime was applied on the surface and incorporated into the soil with a plow and then with a harrow to a depth of approximately 0.3 m. The plantation received 1000 kg·ha⁻¹ of formula fertilizer (5-20-10; N-P₂O₅-K₂O), totaling 50, 200 and 100 kg·ha⁻¹ of N, P₂O₅ and K₂O, respectively. The fertilizer was applied to the soil surface and also incorporated to approximately 0.3 m. Maintenance fertilizer was added annually: approximately 50, 30 and 150 kg·ha⁻¹ of N, P₂O₅ and K₂O in the formulation 11-7-35. The fertilizer was applied on the soil surface without incorporation, in a row coincident with the canopy of the plants. Weed control in these rows was performed annually with the use of an herbicide (with glyphosate as the active ingredient). Between rows, weeds consisted primarily of *Trifolium repens*, *Trifolium pratenses*, *Paspalum notatum* and *Chaptalia nutans*. They were mowed throughout the apple tree cycle at approximately 0.10 m height from the soil surface. The native pasture area, which was used as reference, was located adjacent to the orchards, and natural vegetation was composed mainly of cover crops.

For fungal disease control during the production period of the orchard (October to April), was applied a spray volume of approximately 1000 L·ha⁻¹ of fungicides N-(4,6-dimethylpyrimidine-2-yl) aniline (PYRIMETHANIL) (45 mL 100 L); N-(trichloromethylthio) cyclohex-4-ene-1,2-dicarboximide (CAPTAN) (22 g 100 L); Sulfur (sulfur) (100 g 100 L); and especially Mancozeb (160 g 100 L). The fungicides were applied every seven days, always with rotation between the products. However, if it rained for more than nine consecutive hours, a new application was performed. During the winter period, the plant apple trees received 1000 L·ha⁻¹ spray volume of a copper-based fungicide (Copper II + Oxychloride – Cu oxychloride), one application at the beginning of the production cycle, during the budding of the leaves, and other at the end of the cycle, after leaf senescence, with a total of around 22 fungicide applications per year.

Collection and preparation of soil samples

In October 2011, eight years after the implantation of orchard O8 and 18 years after the implantation of orchard O18, six trenches measuring 0.5 m depth and 0.5 × 0.3 m side were opened in random spots close to the planting rows in each orchard. Six trenches were opened at random points on the NP. Stratified samples were collected in each of the trenches in the following layers: 0.0 – 0.025 m, 0.025 – 0.05 m, 0.05 – 0.10 m, 0.10 – 0.15 m, 0.15 – 0.20 m and 0.20 – 0.40 m. Soil samples were air-dried, ground in a porcelain mortar, passed through a 2-mm mesh sieve, and reserved for analysis.

Physical and chemical characterization of soil

The soil samples were quantified for the levels of clay, silt and sand by the pipette method (Embrapa 1997). Total organic carbon was determined by the Walkley-Black method (Embrapa 1997), and the levels of soil organic matter were measured by multiplying the total organic carbon content by 1,724 (Tedesco et al. 1995). Analysis of soil pH was performed using a soil:water suspension of 1:1 v/v ratio (Tedesco et al. 1995). The levels of available phosphorus (P) and exchangeable potassium (K) were determined in a solution extracted by Mehlich⁻¹ solution (HCl 0.05 mol·L⁻¹ + H₂SO₄ 0.0125 mol·L⁻¹); P was determined by colorimetry (Murphy and Riley 1962) in a spectrophotometer; and K in a flame photometer. The contents of exchangeable Ca²⁺, Mg²⁺ and Al³⁺ were determined in a solution extracted by KCl (1 mol·L⁻¹), whereas Ca and Mg were determined by atomic absorption spectrometry (AAS), and Al by titration with NaOH (Tedesco et al. 1995). Effective cation exchange capacity (CEC_{effective}) was calculated by summing the exchangeable cations of Al³⁺ + Ca²⁺ + Mg²⁺ + K⁺. Cation exchange capacity at pH 7.0 (CEC_{pH 7.0}) was calculated by the sum of exchangeable bases (Ca²⁺ + Mg²⁺ + K⁺) plus H + Al_{SMP} estimated from the equation proposed by Kaminski et al. (2002). Table 1 shows the determined physicochemical parameters.

Chemical fractionation of copper and zinc in soils

The chemical fractionation of Cu and Zn was carried out by adapting the methodology proposed by Tessier

Table 1. Physical and chemical attributes of an Inceptisol under native pasture (NP) and in orchard O8 (8 years old) and orchard O18 (18 years old), both cultivated with apple trees.

Area	Layer	Clay	Silt	Sand	OC	Al	Ca	Mg	P	K	pH	CEC _{pH 7.0}	CTC _{effective}
	(m)		(g·kg ⁻¹)			(cmol _c ·kg ⁻¹)			(mg·kg ⁻¹)			(cmol _c ·kg ⁻¹)	
NP	0 – 0.025	91	545	364	105.5	0.9	4.6	3.6	71	585	5.00	31.5	10.2
	0.025 – 0.05	106	560	334	73.1	3.2	2.4	2.4	3.9	266	3.96	32.9	7.2
	0.05 – 0.10	113	540	347	72.8	5.6	1.4	1.3	2.9	138	3.91	32.1	6.0
	0.10 – 0.15	111	537	352	69.8	7.2	0.8	1.0	1.9	112	3.80	38.4	5.6
	0.15 – 0.20	139	515	345	45.2	7.4	0.7	0.7	1.6	59	3.65	40.6	6.1
	0.20 – 0.40	151	483	366	41.5	8.2	0.5	0.6	0.6	49	3.74	33.6	5.7
O8	0 – 0.025	240	195	565	50.0	0.2	6.5	8.0	35	144	5.19	15.1	10.2
	0.025 – 0.05	257	182	561	47.1	0.1	6.8	3.1	19.7	81	6.11	12.8	10.3
	0.05 – 0.10	259	186	555	37.4	0.1	7.2	3.2	21.7	55	6.22	13.4	10.6
	0.10 – 0.15	272	171	557	38.1	0.1	6.0	4.5	8.6	38	6.41	13.5	10.7
	0.15 – 0.20	272	168	560	32.5	0.1	6.7	4.6	4.8	38	6.09	14.5	11.5
	0.20 – 0.40	213	192	595	13.2	0.2	5.4	3.7	1.5	24	5.29	14.4	9.4
O18	0 – 0.025	209	211	580	63.5	0.2	6.8	6.5	95.6	271	5.18	21.7	14.2
	0.025 – 0.05	216	199	585	55.4	0.1	7.7	2.4	95.3	115	4.93	18.7	10.5
	0.05 – 0.10	209	210	582	41.1	0.1	13.7	3.2	83.1	110	5.07	17.5	10.9
	0.10 – 0.15	212	201	587	27.1	0.2	8.2	3.1	83.3	65	5.37	16.7	11.7
	0.15 – 0.20	213	208	580	27.3	0.1	9.0	3.0	79.1	64	5.51	17.0	12.3
	0.20 – 0.40	241	185	574	29.6	0.2	6.7	3.4	51.8	40	5.47	14.3	10.4

et al. (1979). Soil samples (1 g, in triplicate) were transferred to 50 mL polypropylene tubes to perform the following sequential extractions: the soluble fraction, extracted with deionized water (8 mL) (Cu_{sol} and Zn_{sol}); the exchangeable fraction, extracted with 1.0 mol.L⁻¹ MgCl₂ at pH 7.0 (8 mL) (Cu_{E} and Zn_{E}); the mineral-bound fraction, extracted with 0.04 mol.L⁻¹ hydroxylamine hydrochloride (NH₂OH.HCl) in 25% acetic acid (v/v) at pH 2.0 (20 mL) (Cu_{Min} and Zn_{Min}); and the organic-bound fraction extracted with 0.02 mol.L⁻¹ HNO₃ (3 mL) + 30% H₂O₂ (8 mL) + NH₄OAc in 20% HNO₃ (v/v) (5 mL) (Cu_{OM} and Zn_{OM}). The supernatant was separated by centrifugation at 3200 xg for 30 min, and an aliquot was set aside to determine the Cu and Zn levels. Between each extraction, the samples were agitated with deionized water and centrifuged, and the supernatant was discarded. At the end of the sequential extraction, the remaining residues were dried at 65 °C for 24 h, and then, 0.50 g of the dry residue was subjected to digestion with concentrated HF + HClO₄ to determine the residual Cu and Zn fractions (Cu_{R} and Zn_{R}). To determine the pseudototal levels, 30% H₂O₂ + concentrated HF + concentrated HClO₄ solution was added to 0.5 g samples of dry soil (not subjected to sequential extraction of Cu and Zn) to digest the samples (Cu_{T} and Zn_{T}). In conducting the

study, all glassware was soaked in 10% HNO₃ for at least 24 h and then washed in deionized water three successive times before use. The Cu_{Sum} and Zn_{Sum} represent the sum of fractions Cu_{sol} , Cu_{E} , Cu_{Min} , Cu_{OM} and Cu_{R} and Zn_{sol} , Zn_{E} , Zn_{Min} , Zn_{OM} and Zn_{R} , respectively.

Statistical analysis

The results of the Cu and Zn fractions were submitted to analysis of variance (ANOVA), and when the effects were significant between the layers of the same area, the means were compared by Tukey's test ($\alpha = 5\%$). The differences between areas were evaluated using orthogonal contrasts: native pasture versus orchards and orchard O8 versus orchard O18 within each sampled soil layer. An analysis of variance was performed, and the differences in the proposed contrasts were tested at 5%, 1% and 0.1% probability of error for the F-test.

The fractions of Cu_{sol} , Cu_{E} , Cu_{Min} , Cu_{OM} , Cu_{R} and Cu_{Sum} and the fractions of Zn_{sol} , Zn_{E} , Zn_{Min} , Zn_{OM} , Zn_{R} and Zn_{Sum} were submitted to principal component analysis (PCA). For this purpose, the mean of six replicates was calculated for each layer and subjected to PCA. To minimize the discrepancy between the data, the values were normalized using Eq. 1:

→

(value – minimum value / maximum value – minimum value) \times 2 – 1 (1)

PCA is used to reduce data size and thus facilitate analysis by means of a correlation circle graph (Herlihy and McCarthy 2006).

RESULTS

Fractions of Cu in soil

In the native pasture (NP) soil, the highest levels of Cu_{Sol} , Cu_{R} , Cu_{Sum} and Cu_{T} were observed in the layer of 0.0 – 0.025 m and decreased at deeper levels (Table 2). The levels of Cu_{E} in the NP soil did not differ between layers. The highest levels of Cu_{Min} in the NP soil were found in the layers at 0.0 – 0.025 m

and 0.025 – 0.050 m, whereas for the Cu_{OM} fraction, the higher content was observed in the layers at 0.0 – 0.025 m to 0.10 – 0.15 m. In the soil of O8 and O18 orchards, the highest levels of all Cu fractions were observed in the layer at 0.0 – 0.025 m and decreased in deeper layers.

The Cu_{R} fraction had higher levels compared with the Cu_{Sol} , Cu_{E} , Cu_{Min} , and Cu_{OM} fractions in all soil layers of NP (Table 2). The levels of Cu_{Sol} , Cu_{E} , Cu_{Min} and Cu_{OM} were similar to each other, except for the layer at 0.15 – 0.20 m, in which the Cu_{E} fraction had higher levels than the fractions of Cu_{Sol} , Cu_{Min} and Cu_{OM} (Table 2). In all soil layers evaluated for O8, the levels of Cu_{R} were also higher than those found in the Cu_{Sol} , Cu_{E} , Cu_{Min} and Cu_{OM} fractions, with the exception of the layer at 0.0 – 0.025 m, where there were no differences in the level of Cu_{OM} . For the orchard O18, there were also higher levels of Cu_{E} compared with the Cu_{Sol} , Cu_{E} , Cu_{Min} and

Table 2. Cu fractions of an Inceptisol under native pasture (NP) and in orchard O8 (8 years old) and orchard O18 (18 years old), both cultivated with apple trees.

Layers (m)	Cu fractions (mg·kg ⁻¹)							CV (%)
	Cu _{Sol}	Cu _E	Cu _{Min}	Cu _{OM}	Cu _{Res}	Cu _{Sum}	Cu _{Total}	
NP								
0 – 0.025	0.17 Ad ¹	0.65 Ad	0.56 Ad	0.56 Ad	7.14 Ac	9.07 Ab	12.70 Aa	17.29
0.025 – 0.05	0.11 Bd	0.61 Ad	0.54 Ad	0.50 ABd	5.24 Bc	7.00 Bb	10.59 Ba	16.57
0.05 – 0.10	0.11 Bd	0.70 Ad	0.39 Bd	0.42 ABCd	3.85 BCc	5.48 Cb	7.93 Ba	17.26
0.10 – 0.15	0.07 Cd	0.63 Ad	0.35 Bd	0.41 ACd	3.36 CDc	4.82 CDb	7.99 Ba	26.07
0.15 – 0.20	0.07 Ce	0.65 Ad	0.27 Ce	0.31 CDe	2.50 CDc	3.79 DEb	7.28 Ba	20.40
0.20 – 0.40	0.05 Cd	0.63 Acd	0.14 Dd	0.18 Dd	2.23 Dbc	3.23 Eb	6.99 Ba	28.78
CV (%)	12.42	3.99	8.89	16.67	15.18	10.99	11.28	
O8								
0 – 0.025	0.50 Ae	1.98 Ae	11.89 Ad	18.43 Ac	21.64 Ac	54.44 Ab	86.04 Aa	7.16
0.025 – 0.05	0.27 Bf	0.92 Bef	3.54 Bde	4.80 Bd	17.30 Bc	26.83 Bb	71.01 Ba	8.66
0.05 – 0.10	0.26 Bf	0.60 Cef	2.66 Cde	3.39 Cd	13.01 Cc	19.93 Cb	58.24 Ca	7.57
0.10 – 0.15	0.25 BCe	0.42 CDe	2.46 CDd	2.19 Dde	9.72 Dc	15.05 Db	51.56 Da	8.85
0.15 – 0.20	0.22 Cd	0.26 DEd	2.07 DEd	1.81 Dd	7.64 Dc	12.01 DEb	43.22 Ea	9.49
0.20 – 0.40	0.15 Dd	0.09 Ed	1.79 Ed	1.47 Dd	7.13 Dc	10.62 Eb	28.67 Fa	13.43
CV (%)	5.83	14.15	6.04	5.93	11.05	6.11	5.04	
O18								
0 – 0.025	0.70 Af	1.15 Af	19.13 Ae	50.13 Ac	35.39 Ad	106.50 Ab	120.16 Aa	6.34
0.025 – 0.05	0.39 Be	0.97 Be	12.55 Bd	23.92 Bc	24.70 Bc	62.52 Bb	78.02 Ba	8.58
0.05 – 0.10	0.28 Cf	0.93 Bf	11.82 BCe	15.28 Cd	18.63 Cc	46.93 Cb	68.71 Ca	5.94
0.10 – 0.15	0.24 Ce	0.70 Ce	10.99 Cd	13.99 Cd	18.96 Cc	44.87 Cb	55.04 Da	8.10
0.15 – 0.20	0.24 Ce	0.68 Ce	8.64 Dd	13.92 Cc	12.14 Dcd	35.61 Db	40.50 Ea	9.95
0.20 – 0.40	0.23 Cd	0.66 Cd	6.59 Ec	5.84 Dc	5.53 Ec	18.85 Eb	22.52 Fa	12.09
CV (%)	7.86	6.36	3.91	4.55	11.67	4.47	5.39	

¹Means followed by the same capital letter in a column and lower case letter in a row do not differ by Tukey's test ($p < 0.05$). CV = coefficient of variation.

Cu_{OM} fractions only in the layers at 0.05 – 0.10 m and 0.10 – 0.15 m. In the soil layer from 0.0 – 0.025 m in orchard O18, the level of Cu_{Org} was higher than the levels of Cu_R, Cu_{Min}, Cu_E and Cu_{Sol} (Table 2).

The means of the levels of all Cu fractions in the six evaluated layers (0.0 – 0.025 m, 0.025 – 0.05 m, 0.05 – 0.10 m, 0.10 – 0.15 m, 0.15 – 0.20 m and 0.20 – 0.40 m), except for the Cu_E fraction, were higher in the soil of orchards O8 and O18, compared with the levels of soil under natural pasture (Tables 2 and 3). This result was also observed when comparing the means of Cu levels in the six evaluated layers between the two orchards, where all fractions, except for Cu_E, were higher in the orchard O18 compared with O8 (Tables 2 and 3). Furthermore, the Cu_{Min}, Cu_{OM}, Cu_E and Cu_{Sum} fractions were higher in the orchard O18 compared with O8 in the six layers (Tables 2 and 3). The exceptions were the levels of fractions Cu_E and Cu_{Total} (0.025 – 0.05 m), Cu_{Sol} (0.05 – 0.10 m) and Cu_{Sol} and Cu_{Total} (0.10 – 0.15 m and 0.15 – 0.20 m), which showed no differences (Table 3).

The native pasture layer at 0.0 – 0.025 m had the highest percentage of Cu in the residual fraction (Cu_{Res}) (Figure 1), but in the soil of O8 and O18, the percentage of Cu_{Res} decreased compared with that of the native pasture soil. There was also an increase in the Cu_{OM} and Cu_{Min} fractions. In the soil of O18, the percentage of Cu_{OM} was higher than that of the Cu_E and Cu_{Res} fractions. In the layer at 0.025 – 0.050 m, the Cu_{Res} percentages decreased in the soil of O8 and O18, compared with that of the native pasture soil, because of an increase in the percentage of Cu_{OM}, especially in the soil of O18. In the layer at 0.05 – 0.10 m, the percentages of Cu_{Res} were similar between the soil under native pasture and that of O8. However, in the soil of O8, the percentages of Cu_{OM} and Cu_{Min} were higher compared to the soil under

native pasture. The highest percentages of Cu_{OM} and Cu_{Min} were observed in the soil of O18, compared with the soils of the native pasture and O8. In the layers at 0.10 – 0.15 m, 0.15 – 0.20 m and 0.20 – 0.40 m, the percentages of Cu_{OM} and Cu_{Min} increased in the soil of O8, and especially in the soil of O18. In the latter, because of the increase in the Cu_{OM} and Cu_E fractions, a lower proportion of Cu_{Res} was present compared with NP and O8. For the Cu_E fraction, the soil under native pasture had the highest percentages in all evaluated layers.

The principal component analysis (PCA) of the Cu fractions showed a relation between the main component 1 (factor 1) and the main component 2 (factor 2), which was split among the areas (NP, O8 and O18) (Figure 2a), accounting for 98.87% of the results. The areas of native pasture and O8 were separated from O18 by factor 1 of PCA, which helps to explain 95.4% of the results. The result was strongly influenced by the variables Cu_{Sol}, Cu_E, Cu_{Min}, Cu_{OM} and Cu_{Res}. Factor 2 of the PCA, which helps explain only 3.4% of the results, separated the areas of orchards O8 and O18 from the native pasture area.

Zn fractions in soil

The highest levels of Zn fractions (Zn_{Sol}, Zn_E, Zn_{Min}, Zn_{OM}, Zn_R and Zn_{Sum}) and Zn_{Total} were found in the layer at 0.0 – 0.025 m in all areas assessed, with a subsequent decrease in the levels because of increasing depth (Table 4). The exception to this pattern was found in the soil of O8 and O18, where the Zn_{OM} and Zn_{Total} fractions did not differ between the layers at 0.0 – 0.025 m and 0.025 – 0.050 m. Furthermore, in the soil of O18, the Zn_{Res} fraction showed equal values between those two layers and the layer at 0.0 – 0.15 m.

Table 3. Orthogonal contrasts between the Cu fractions of an Inceptisol under native pasture (NP) and in orchard O8 (8 years old) and orchard O18 (18 years old), both cultivated with apple trees.

Contrasts	Cu _{Sol}	Cu _E	Cu _{Min}	Cu _{OM}	Cu _R	Cu _{Sum}	Cu _{Total}
NP vs O8 and O18 ¹	***	Ns	***	***	***	***	***
O8 vs O18 ¹	***	Ns	***	***	***	***	*
Layer 0.0 – 0.025 O8 vs O18	***	***	***	***	***	***	***
Layer 0.025 – 0.05 O8 vs O18	***	Ns	***	***	***	***	ns
Layer 0.05 – 0.10 O8 vs O18	ns	***	***	***	***	***	***
Layer 0.10 – 0.15 O8 vs O18	ns	***	***	***	***	***	ns
Layer 0.15 – 0.20 O8 vs O18	ns	***	***	***	**	***	ns
Layer 0.20 – 0.40 O8 vs O18	***	***	***	***	*	***	***

¹Means of six evaluated layers. *Significant at 5%; ** significant at 1%; *** significant at 0.1%; ns = no significant.




Figure 1. Relative percentage of each Cu fraction obtained from a sequential extraction compared with the sum of the amount extracted from six layers of an Inceptisol under native pasture and in orchards cultivated with apple trees (O8 and O18).

In all layers and areas assessed, the highest Zn levels were found in Zn_{Total} , except for the Zn_{sum} fraction, which showed values equal to those of the Zn_{Total} fraction in the layer at 0.05 – 0.10 m for O8. For the Zn_{Res} fraction, there were higher levels compared with the Zn_{Sol} , Zn_E , Zn_{Min} and Zn_{OM} fractions in all layers of the soil under natural pasture, O8 and O18. The level of the Zn_{Min} fraction in the soil under native pasture, O8 and O18 in all layers was lower than that of Zn_{Res} , but higher than the levels of Zn_{Sol} , Zn_E and Zn_{OM} . In the soil of O18, in all layers, the Zn_E levels

were higher than the levels in the soil layers of the native pasture and O8.

Contrast orthogonal analysis showed that the mean Zn content of the six measured layers (0.0 – 0.025 m, 0.025 – 0.05 m, 0.05 – 0.10 m, 0.10 – 0.15 m, 0.15 – 0.20 m and 0.20 – 0.40 m) and of all fractions (Zn_E , Zn_{Min} , Zn_{OM} , Zn_R , Zn_{sum} and Zn_{Total}) was higher in the soil of O8 and O18, compared with the soil under native pasture (Tables 4 and 5). This result was also found by comparing the means of Cu levels measured in the six

→

Table 4. Zn fractions of an Inceptisoi under native pasture (NP) and in orchard O8 (8 years old) and orchard O18 (18 years old), both cultivated with apple trees.

Layer (m)	Zn fractions (mg·kg ⁻¹)							CV (%)
	Zn _{Sol}	Zn _E	Zn _{Min}	Zn _{OM}	Zn _R	Zn _{Sum}	Zn _{Total}	
NP								
0 – 0.025	0.18 Ae ¹	3.21 Ae	9.38 Ad	1.89 Ae	16.24 Ac	30.90 Ab	39.57 Aa	9.06
0.025 – 0.05	0.10 Be	1.70 Be	5.04 Bd	1.78 Be	12.02 Bc	20.62 Bb	24.38 Ba	10.38
0.05 – 0.10	0.08 BCe	1.51 Bcd	4.03 Cc	1.26 Bcd	9.35 BCb	16.22 Cb	19.59 Ca	20.38
0.10 – 0.15	0.07 BCe	1.42 Be	3.46 Dd	1.03 Be	8.29 Cc	14.26 CDb	17.97 CDa	13.21
0.15 – 0.20	0.06 CDe	0.73 Cde	3.01 DEd	0.43 Ce	7.08 Cc	11.39 DEb	14.53 Da	17.68
0.20 – 0.40	0.03 De	0.66 Cde	2.77 Ed	0.27 Ce	6.76 Cc	10.48 Eb	14.05 Da	18.51
CV (%)	19.30	9.24	4.83	9.96	11.95	7.24	9.24	
O8								
0 – 0.025	0.16 Ae	3.56 Ae	14.65 Ad	1.83 Ae	48.55 Ac	68.74 Ab	75.70 Aa	6.48
0.025 – 0.05	0.12 Be	0.57 Be	6.63 Bd	1.42 Be	33.40 Bc	42.14 Bb	76.25 Aa	4.93
0.05 – 0.10	0.11 BCc	0.42 Cc	5.29 Cc	1.40 Bc	23.51 Cb	30.74 Ca	34.16 Ba	18.84
0.10 – 0.15	0.11 BCDd	0.39 Cd	3.47 Dd	1.22 BCD	18.66 CDc	23.85 Db	30.37 Ba	13.19
0.15 – 0.20	0.08 CDd	0.41 Cd	1.83 Ed	1.07 Cd	15.32 Dc	18.71 Db	28.86 BCa	10.35
0.20 – 0.40	0.08 Dc	0.13 Dc	1.41 Ec	1.00 Cc	16.22 Db	18.82 Db	22.82 Ca	15.87
CV (%)	13.58	3.24	4.94	9.83	9.92	7.72	6.46	
O18								
0 – 0.025	0.22 Ad	12.03 Ad	41.75 Ac	4.56 Ad	49.69 Ac	108.25 Ab	155.00 Aa	12.67
0.025 – 0.05	0.16 Be	6.49 Be	29.74 Bd	4.36 Ae	45.88 Ac	86.63 Bb	100.86 Ba	7.44
0.05 – 0.10	0.15 BCe	4.09 Ce	28.52 BCD	3.26 Be	43.71 Ac	79.73 BCb	96.99 Ba	5.77
0.10 – 0.15	0.12 BCDe	2.79 De	26.94 CDd	2.72 Be	41.54 Ac	74.11 Cb	93.25 Ba	11.86
0.15 – 0.20	0.11 CDd	2.13 Ed	25.42 Dc	2.63 Bd	30.70 Bc	60.98 Db	89.28 Ba	9.99
0.20 – 0.40	0.09 De	1.73 Fe	14.56 Ed	1.88 Ce	24.21 Bc	42.48 Eb	59.59 Ca	14.25
CV (%)	15.32	3.25	3.15	9.68	10.04	5.39	8.43	

¹Means followed by the same capital letter in a column and lower case in a row do not differ by Tukey's test ($p < 0.05$). CV = coefficient of variation.

Table 5. Orthogonal contrasts between Zn fractions of an Inceptisoi under native pasture (NP) and in orchard O8 (8 years old) and orchard O18 (18 years old), both cultivated with apple trees.

Contrasts	Zn _{Sol}	Zn _E	Zn _{Min}	Zn _{OM}	Zn _{Res}	Zn _{Sum}	Zn _{Total}
NP vs O8 and O18 ¹	***	***	***	***	***	***	***
O8 vs O18 ¹	***	***	***	***	***	***	***
Layer 0.0 – 0.025 O8 vs O18	**	***	***	***	ns	***	***
Layer 0.025 – 0.05 O8 vs O18	*	***	***	***	***	***	***
Layer 0.05 – 0.10 O8 vs O18	*	***	***	***	***	***	***
Layer 0.10 – 0.15 O8 vs O18	ns	***	***	***	***	***	***
Layer 0.15 – 0.20 O8 vs O18	**	***	***	***	***	***	***
Layer 0.20 – 0.40 O8 vs O18	ns	***	***	***	**	***	***

¹Means of six evaluated layers. *Significant at 5%; ** significant at 1%; *** significant a 0.1%; ns = no significant.

layers between the two orchards, where all the fractions were higher in orchard O18 compared with O8. Except for the Zn_{Sol} fraction in the layers at 0.10 – 0.15 m

and 0.20 – 0.40 m, and Zn_R in the layer at 0.0 – 0.025 m, all other fractions in all layers were higher in the soil of O18, compared with that of O8 (Tables 4 and 5).

→

In the layers at 0.0 – 0.025 m, 0.025 – 0.050 m, 0.05 – 0.10 m, 0.10 – 0.15 m, 0.15 – 0.20 m and 0.20 – 0.40 m, the highest percentages of Zn in soil under the native pasture, O8 and O18 were found in the residual fraction (Zn_{Res}), being that the highest percentages were in the soil of O8 (Figure 2). However, in the soil of O18 was an increase of Zn content in the mineral fraction (Zn_{Min}) in all layers, compared with the soils of the native pasture and O8.

The PCA of Zn fractions showed the relation between the main component 1 (factor 1) and main component 2 (factor 2), which suggested separation of the areas assessed




Figure 2. The ratio between the principal component 1 (factor 1) and principal component 2 (factor 2), mean levels of (a) Cu and (b) Zn in the layer at 0.0 – 0.40 m. Cu_{sol} and Zn_{sol} – soluble fraction extracted with H_2O ; Cu_e and Zn_e – exchangeable fraction extracted with $MgCl_2$ 1.0 mol·L⁻¹; Cu_{min} and Zn_{min} – mineral fraction extracted with $NH_2OH\cdot HCl$ 0.04 mol L⁻¹ in 25% CH_3COOH ; Cu_{om} and Zn_{om} – organic fraction extracted with HNO_3 0.02 mol·L⁻¹ + 8 mL of 30% H_2O_2 ; Cu_{sum} and Zn_{sum} – sum of all fractions; and Cu_{res} and Zn_{res} – residual fraction extracted with $HNO_3 + HCl + H_2O_2$.

(Figure 2b), supporting the PCA of Cu fractions. Factors 1 and 2 accounted for 98.96% of the results. Factor 1, which explained 93.19% of the data, separated the areas of native pasture and O8 from that of O18. The Zn_{sol} , Zn_e , Zn_{min} , Zn_{om} and Zn_{res} fractions contributed greatly to this result, whereas factor 2 contributed only 5.7% to the result and separated O8 and O18 from the native pasture area.

DISCUSSION

The highest levels of Cu fractions (Cu_{sol} , Cu_e , Cu_{min} , Cu_{om} , Cu_R , Cu_{sum} and Cu_{total}) were observed in the surface layer (0.0 – 0.025 m), especially in the soils of orchards O8 and O18. They can be attributed to years of applications of foliar fungicides, such as Mancozeb, Pyrimethanil and $[CuSO_4 \cdot 5H_2O + Ca(OH)_2]$, with the aim of preventing fungal diseases and also to promote dormancy breaking of plants (Verdisson et al. 2001; Komárek et al. 2010; Duplay et al. 2014). Thus, Cu may have accumulated on the soil surface because the applications were directional to the trees and Cu was washed off the surface of leaves by rainfall, and also by deposition of senescent leaves and pruned branches containing Cu derived from these applications (Brunetto et al. 2013). Furthermore, the higher level of Cu on the soil surface was also favored by the absence of soil disturbance in planting rows and between rows of apple trees over the years, by the cycling of Cu through deposition and subsequent decomposition of senescent leaves and pruned branches of apple trees, or even by the decomposition of roots in the soil (Amery et al. 2007; Yuan et al. 2007; Wang and Mulligan 2013; Pérez-Esteban et al. 2013). Therefore, the organic matter content in the upper layers of the soil are maintained and some Cu remains on the soil surface because its electron configuration is $[Ar]3d^{10}4s^1$, thus making it highly reactive with the functional groups of organic matter that contain S and N, in addition to the carboxyl and phenolic functional groups (Croué et al. 2003). Furthermore, organic compounds provide Cu with high binding energy and low desorption (Lair et al. 2006; Casali et al. 2008; Brunetto et al. 2013).

In the case of the orchards, especially O18, the higher Cu content, represented by Cu_{total} , can be attributed to the 22 annual fungicide applications throughout the years. In addition to the factors mentioned above, the fresh pulp of apple fruit is between 2 and 4 mg·kg⁻¹ Cu (Stüpp et al. 2013). In this sense, an orchard with production of

60 t·ha⁻¹ has exported 0.24 kg·ha⁻¹ Cu, a fact that contributes to a higher accumulation of this trace element in the soil. Cu accumulation in the orchards was also favored because the relief at the site of the orchards is gently undulating, and cover crops are planted between the orchard rows, favoring the deposition of plant residues on the soil surface, which minimizes the transfer of Cu by the drained solution (Loures et al. 2006; Pinho et al. 2006).

The higher Cu content in the soil – mostly in the organic fraction (Cu_{OM}), compared with other fractions (Cu_{Res}, Cu_{Min}, Cu_E and Cu_{Sol}), especially in the surface layer (0.0–0.25 m), with increasing age of the orchards (eight years for O8 and 18 years for O18) and, therefore, a larger number of Cu-containing fungicide applications and higher nutrient-cycling time – can be attributed to high affinity and reactivity with the functional groups of the soil organic matter (Pérez-Novo et al. 2008). Therefore, because Cu is part of the organic fraction, the adoption of conservation management practices in orchards is crucial, for example, the deposition and maintenance of plant residues on the soil surface helps maintain the levels of soil organic matter and the absence of soil disturbance minimizes the mineralization of organic matter (Ferreira et al. 2013; López-Piñeiro et al. 2013). Nevertheless, the migration of the Cu to a depth of 0.40 m in the soil of orchards O8 and O18 was observed through the level of Cu_{Total} and Cu_{OM}, Cu_{Min} and Cu_{Res} fractions, (Table 3, Figure 1). This may have happened due to the saturation of functional groups of organic matter (Besnard et al. 2001; Croué et al. 2003), by the saturation of functional groups of Fe and Mn oxides (Bradl 2004), and the silanol and aluminol groups of phyllosilicates (Sposito 1989; McBride 1994). Furthermore, the decrease in pH values in the soil profile (Table 1) may change the sorption capacity of the clay minerals, especially iron oxides (Sposito 1989). At acidic pH values, sorption is lower and desorption is facilitated (Brunetto et al. 2013).

The highest levels of Zn_{Total}, in particular, as has occurred with Cu in the most superficial soil layer (0.0–0.025 m), mainly in the soil of orchards O8 and O18, are due to the unidirectional annual applications of fungicides containing Zn in their composition, such as Mancozeb (Komárek et al. 2010), Zn being washed off the surface of leaves by rainfall, and deposition of Zn-containing leaves or pruned branches (Fernández-Calviño et al. 2012; Brunetto et al. 2014; Duplay et al. 2014). Furthermore, the higher accumulation of Zn_{Total} in the surface layers of the soil of the orchards, particularly orchard O18, is associated with a larger number of Zn-containing foliar fungicide applications for the preventive control of fungal diseases over the years (Brunetto et al. 2013). As seen for Cu, accumulation of Zn forms on the surface is also associated with

an absence of soil disturbance in the rows and between the rows of apple trees, in addition to Zn cycling in the soil by the ground cover plants located in the rows and between rows of orchards (Couto et al. 2015; Hristovski et al. 2014).

In all soil layers of NP and orchards O8 and O18, higher levels of Zn were observed in the Zn_{Res} and Zn_{Min} fractions (Table 4 and Figure 3b), and these fractions are more stable and characterized by low availability and mobility in soil (Ramos 2006; Fernández-Calviño et al. 2012). Zn_{Total} and the Zn_E, Zn_{Min} and Zn_R fractions (Table 4 and Figure 3b) showed migration of Zn to 0.40 m depth. This may have occurred because of the migration of Zn by the flow of water in the soil because a fraction of Zn tends to remain in the soil solution in free form or as soluble ionic pairs (Citeau et al. 2003) and also by turbulent water flow in macropores and by soil cracks or even bioturbation (Giroto et al. 2010; Brunetto et al. 2013). In addition, Zn can bind to organic acids with low molecular weight, increasing its mobility in the soil profile (Wang and Mulligan 2013). This may occur because the formation of metal complexes with organic compounds with low molecular weight alters the balance between the metals in the labile form in the solid phase and in the soil solution, thus, dissolved organic compounds can also increase the release of Cu in the solid phase, and consequently increase its mobility in the soil (Giroto et al. 2010). In addition, Zn mobility may have been facilitated because of the decrease in water pH in the deeper layers of the soil, contributing to the formation of stable (innersphere) complexes between Zn and the functional groups of organic and mineral fractions of soil (Brunetto et al. 2013).

The increase in the Zn_E fraction in the soil profile, which is a bioavailable fraction (Tessier et al. 1979) especially in the orchard O18 (Table 4), increases the potential for contamination of subsoil water (Banas et al. 2010; Fernández-Calviño et al. 2012). However, the increase in Zn_E in the upper layers, where the roots of apple trees and ground cover plant species are located in the rows and inter-rows of the orchards, may potentiate toxicity to plants (Nagajyoti et al. 2010; Fernández-Calviño et al. 2012). One example is the data obtained by Zhang et al. (2013) and Hansi et al. (2014). The first authors, studying the length of barley (*Hordeum vulgare*) roots grown in soil in a greenhouse with the addition of 0, 12.5, 25, 50, 100, 200, 400 and 800 mg Cu·kg⁻¹, observed that root elongation drastically decreases with increasing Cu in solution.

Decreases of roots systems is not desirable because it reduces the nutrient cycling by ground cover plant species and the amount of plant residues deposited on the surface of soil, which are responsible for breaking the kinetic energy of raindrops, as




Figure 3. Relative percentage of each Zn fraction obtained from a sequential extraction compared with the sum of the amount extracted from six layers of an Inceptisol under native pasture and in orchards cultivated with apple trees (O8 and O18).

well as reducing transportation of the solution on the soil surface. Moreover, it helps maintain the organic matter content in the soil (Brunetto et al. 2013). The increased Zn concentration in the soil can also cause toxicity to microorganisms, adversely affecting, for example, the decomposition of the crop residue deposited on the ground and the mineralization of soil organic matter (Gillera et al. 1998). Zn_E also increases in the most superficial soil layer, which can enhance the transfer of Zn on the soil surface by the solution drained or adsorbed in the sediment (Fernández-Calviño et al. 2008; Fernández-Calviño et al. 2012), which, in turn, enhances contamination of surface water adjacent to the orchards.

CONCLUSION

The Cu and Zn fractions increased in the surface soil layer (0.0 – 0.025 m) of apple orchards compared to native pasture cultivation, being associated with the time of cultivation.

Most Cu was distributed in the less stable organic fraction in the upper layers of apple orchards, which may potentiate toxicity to plants and transfer Cu to subsurface waters. In deeper layers, however, Cu was also distributed in the residual and mineral fractions, which are more stable.

Most Zn in the soil of apple orchards was distributed in the residual and mineral fractions, indicating low mobility, as well as potential toxicity to plants.

ACKNOWLEDGEMENTS

We are thankful to the CNPq (National Council for Scientific and Technological Development) for its financial support and for the research productivity grant awarded to the first author.

ORCID IDs

G. Brunetto

 <https://orcid.org/0000-0002-3174-9992>

L. Benedit

 <https://orcid.org/0000-0002-0560-8790>

V. G. Ambrosini

 <https://orcid.org/0000-0001-9069-850X>

J. J. Comin

 <https://orcid.org/0000-0002-6176-6208>

G. W. B. de Melo

 <https://orcid.org/0000-0001-5825-3500>

M. A. dos Santos

 <https://orcid.org/0000-0003-2814-0788>

C. R. Lourenzi

 <https://orcid.org/0000-0002-0347-0003>

A. Loss

 <https://orcid.org/0000-0002-3005-6158>

P. Belli Filho

 <https://orcid.org/0000-0003-4338-2830>

D. E. Schmitt

 <https://orcid.org/0000-0001-9665-9704>

R. Couto

 <https://orcid.org/0000-0003-1071-4165>

REFERENCES

- Associação Catarinense de Empresas de Tecnologia (2014). Agronegócio e tecnologia. Anuário 2014. Estado de Santa Catarina. Available at: http://www.acate.com.br/sites/default/files/anuarioacate_0.pdf. Accessed on April 27, 2016.
- Amery, F., Degryse, F., Degeling, W., Smolders, E. and Merckx, R. (2007). The copper-mobilizing-potential of dissolved organic matter in soils varies 10-fold depending on soil incubation and extraction procedures. *Environmental Science & Technology*, 41, 2277-2281. <http://doi.org/10.1021/es062166r>.
- Banas, D., Marin, B., Skrabber, S., Chopin, E. I. B. and Zanella, A. (2010). Copper mobilization affected by weather conditions in a stormwater detention system receiving runoff waters from vineyard soils (Champagne, France). *Environmental Pollution*, 158, 476-482. <http://doi.org/10.1016/j.envpol.2009.08.034>.
- Besnard, E., Chenu, C. and Robert, M. (2001). Influence of organic amendments on copper distribution among particle-size and density fractions in Champagne vineyard soils. *Environmental Pollution*, 112, 329-337. [https://doi.org/10.1016/S0269-7491\(00\)00151-2](https://doi.org/10.1016/S0269-7491(00)00151-2).
- Bradl, H. B. (2004). Adsorption of heavy metal ions on soils and soils constituents. *Journal of Colloid and Interface Science*, 277, 1-18. <http://doi.org/10.1016/j.jcis.2004.04.005>.
- Brunetto, G., Miotto, A., Ceretta, C. A., Schmitt, D. E., Heinzen, J., Moraes, M. P., Canton, L., Tiecher, T., Comin, J. J. and Giroto E. (2014). Mobility of copper and zinc fractions in fungicide-amended vineyard sandy soils. *Archives of Agronomy and Soil Science*, 60, 609-624. <http://doi.org/10.1080/03650340.2013.826348>.
- Casali, C. A., Moterle, D. F., Rheinheimer, D. S., Brunetto, G., Corcini, A. L. M., Kaminski, J. and Melo, G. W. B. (2008). Formas e dessorção de cobre em solos cultivados com videira na Serra Gaúcha do Rio Grande do Sul. *Revista Brasileira de Ciência Solo*, 32, 1479-1487. <http://doi.org/10.1590/S0100-06832008000400012>.
- Citeau, L., Lamy, I., Van Oort, F. and Elsass, F. (2003). Colloidal facilitated transfer of metals in soils under different land use. *Colloids and Surfaces A*, 217, 11-19. [http://doi.org/10.1016/S0927-7757\(02\)00554-X](http://doi.org/10.1016/S0927-7757(02)00554-X).

- Couto, R. R., Benedet, L., Comin, J. J., Belli Filho, P., Martins, S. R., Gatiboni, L. C., Radetski, M., Valois, C. M., Ambrosini, V. G. and Brunetto, G. (2015). Accumulation of copper and zinc fractions in vineyard soil in the mid-western region of Santa Catarina, Brazil. *Environmental Earth Sciences*, 73, 6379-6386. <http://doi.org/10.1007/s12665-014-3861-x>.
- Croué, J. P., Benedetti, M. F., Violleau, D. and Leenheer, J. A. (2003). Characterization and copper binding of humic and nonhumic organic matter isolated from the south platte river: evidence for the presence of nitrogenous binding site. *Environmental Science & Technology*, 37, 328-336. <http://doi.org/10.1021/es020676p>.
- Duplay, J., Semhi, K., Errais, E., Imfeld, G., Babcsanyi, I. and Perrone, T. (2014). Copper, zinc, lead and cadmium bioavailability and retention in vineyard soils (Rouffach, France): the impact of cultural practices. *Geoderma*, 230-231, 318-328. <http://doi.org/10.1016/j.geoderma.2014.04.022>.
- Empresa Brasileira de Pesquisa Agropecuária (1997). Manual de métodos de análise de solo. Centro Nacional de Pesquisas de Solo, Rio de Janeiro: Embrapa-CNPq.
- Empresa Brasileira de Pesquisa Agropecuária (2013). Sistema Brasileiro de Classificação de Solos (Brazilian System for Soil Classification). Centro Nacional de pesquisas de solo, Rio de Janeiro: Embrapa-CNPq.
- Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina (2013). Síntese Anual da Agricultura de Santa Catarina. Instituto de Planejamento e Economia Agrícola de Santa Catarina. Available at: http://docweb.epagri.sc.gov.br/website_cepa/publicacoes/Sintese_2014.pdf. Accessed on May 5, 2016.
- Fernández-Calviño, D., Pateiro-Moure, M., López-Periago, E., Arias-Estévez, M. and Nóvoa-Muñoz, J. C. (2008). Copper distribution and acid-base mobilization in vineyard soils and sediments from Galicia (NW Spain). *European Journal of Soil Science*, 59, 315-326. <http://doi.org/10.1111/j.1365-2389.2007.01004.x>.
- Fernández-Calviño, D., Pateiro-Moure, M., Nóvoa-Muñoz, J. C., Garrido-Rodrigues, B. and Arias-Estévez, M. (2012). Zinc distribution and acid-base mobilization in vineyard soils and sediments. *Science of the Total Environment*, 414, 470-479. <http://doi.org/10.1016/j.scitotenv.2011.10.033>.
- Fernández-Calviño, D., Soler-Rovira, P., Polo, A., Arias-Estévez, M. and Plaza, C. (2010). Influence of humified organic matter on copper behavior in acid polluted soils. *Environmental Pollution*, 158, 3634-3641. <http://doi.org/10.1016/j.envpol.2010.08.005>.
- Ferreira, A. O., Amado, T. J. C., Nicoloso, R. S., Sá, J. C. M., Fiorin, J. E., Hansel, D. S. S. and Meneffee, D. (2013). Soil carbon stratification affected by long-term tillage and cropping systems in southern Brazil. *Soil and Tillage Research*, 133, 65-74. <http://doi.org/10.1016/j.still.2013.05.011>.
- Frankowski, M., Ziola-Frankowska, A., Kowalski, A. and Siepak, J. (2010). Fractionation of heavy metals in bottom sediments using Tessier procedure. *Environmental Earth Sciences*, 60, 1165-1178. <http://doi.org/10.1007/s12665-009-0258-3>.
- Gillera, K. E., Witter, E. and McGrath, S. P. (1998). Toxicity of heavy metals to microorganisms and microbial processes in agricultural soils: a review. *Soil Biology and Biochemistry*, 30, 1389-1314. [https://doi.org/10.1016/S0038-0717\(97\)00270-8](https://doi.org/10.1016/S0038-0717(97)00270-8).
- Giroto, E., Ceretta, C. A., Brunetto, G., Santos, D. R., Silva, L. S., Lourenzi, C. R., Lorensini, F., Vieira R. C. B. and Schmatz, R. (2010). Acúmulo e formas de cobre e zinco no solo após aplicações sucessivas de dejetos líquidos de suínos. *Revista Brasileira de Ciência do Solo*, 34, 955-965. <http://dx.doi.org/10.1590/S0100-06832010000300037>.
- Hansi, M., Weidenhamer, J. D. and Sinkkonen, A. (2014). Plant growth responses to inorganic environmental contaminants are density-dependent: experiments with copper sulfate, barley and lettuce. *Environmental Pollution*, 184, 443-448. <http://doi.org/10.1016/j.envpol.2013.09.027>.
- Herlihy, M. and McCarthy, J. (2006). Association of soil test phosphorus with phosphorus fractions and adsorption characteristics. *Nutrient Cycling in Agroecosystems*, 75, 79-90. <http://doi.org/10.1007/s10705-006-9013-2>.
- Hristovski, S., Berg, B. and Melovski, L. (2014). Limitless decomposition in leaf litter of Common beech: patterns, nutrients' and heavy metal's dynamics. *Pedobiologia*, 57, 131-138. <http://doi.org/10.1016/j.pedobi.2014.01.005>.
- Kaminski, J., Gatiboni, L. C., Rheinheimer, D. S., Martins, J. R., Santos, E. J. S. and Titsot, C. A. (2002). Estimativa da acidez potencial em solos e sua implicação no cálculo da necessidade de calcário. *Revista Brasileira de Ciência do Solo*, 26, 1107-1113. <http://doi.org/10.1590/S0100-06832002000400029>.
- Komárek, M., Čadková, E., Chrastný, V., Bordas, F. and Bollinger, J. C. (2010). Contamination of vineyard soils with fungicides: a review of environmental and toxicological aspects. *Environment International*, 36, 138-151. <http://doi.org/10.1016/j.envint.2009.10.005>.

- Lair, G. J., Gerzabek, M. H., Haberhauer, G., Jakusch, M. and Kirchmann, H. (2006). Response of the sorption behavior of Cu, Cd, and Zn to different soil management. *Journal of Plant Nutrition and Soil Science*, 169, 60-68. <http://doi.org/10.1002/jpln.200521752>.
- López-Piñeiro, A., Muñoz, A., Zamora, E. and Ramírez, M. (2013). Influence of the management regime and phenological state of the vines on the physicochemical properties and the seasonal fluctuations of the microorganisms in a vineyard soil under semi-arid conditions. *Soil and Tillage Research*, 126, 119-126. <http://doi.org/10.1016/j.still.2012.09.007>.
- Loures, A. P. S., Soares, A. A., Matos, A. T., Cecon, P. R. and Pereira, O. G. (2006). Remoção de fósforo em sistema de tratamento de esgoto doméstico, por escoamento superficial. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 10, 706-714. <http://dx.doi.org/10.1590/S1415-43662006000300025>.
- Mackie, K. A., Müller, T. and Kandeler, E. (2012). Remediation of copper in vineyards – a mini review. *Environmental Pollution*, 167, 16-26. <http://doi.org/10.1016/j.envpol.2012.03.023>.
- Marchand, L., Nsanganwimana, F., Cook, B. J., Vystavna, Y., Huneau, F., Le Coustumer, P., Lamy, J. B., Oustrière, N. and Mench, M. (2014). Trace element transfer from soil to leaves of macrophytes along the Jalle d'Eysines River, France and their potential use as contamination biomonitors. *Ecological Indicators*, 46, 425-437. <http://doi.org/10.1016/j.ecolind.2014.07.011>.
- McBride, M. B. (1994). *Environmental chemistry of soils*. New York: Oxford University Press.
- Michaud, A. M., Bravin, M. N., Galleguillos, M. and Hinsinger, P. (2007). Copper uptake and phytotoxicity as assessed in situ for durum wheat (*Triticum turgidum durum* L.) cultivated in Cu-contaminated, former vineyard soils. *Plant and Soil*, 298, 99-111. <http://doi.org/10.1007/s11104-007-9343-0>.
- Mileusnić, M., Mapani, B. S., Kamona, A. F., Ružičić, S., Mapaure, I. and Chimwamurombe, P. M. (2014). Assessment of agricultural soil contamination by potentially toxic metals dispersed from improperly disposed tailings, Kombat mine, Namibia. *Journal of Geochemical Exploration*, 144, 409-420. <http://doi.org/10.1016/j.gexplo.2014.01.009>.
- Murphy, J. and Riley, J. P. (1962). A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta*, 27, 31-36. [http://dx.doi.org/10.1016/S0003-2670\(00\)88444-5](http://dx.doi.org/10.1016/S0003-2670(00)88444-5).
- Nagajyoti, P. C., Lee, K. D. and Sreekanth, T. V. M. (2010). Heavy metals, occurrence and toxicity for plants: a review. *Environmental Chemistry Letters*, 8, 199-216. <http://doi.org/10.1007/s10311-010-0297-8>.
- Pandolfo, C., Braga, H. J., Silva Jr., V. P., Massignam, A. M., Pereira, E. S., Thomé, V. M. R. and Valci, F. V. (2002). *Atlas climatológico do Estado de Santa Catarina*. Florianópolis: Epagri. CD-Rom.
- Pérez-Esteban, J., Escolástico, C., Moliner, A. and Masaguer, A. (2013). Chemical speciation and mobilization of copper and zinc in naturally contaminated mine soils with citric and tartaric acids. *Chemosphere*, 90, 276-283. <http://doi.org/10.1016/j.chemosphere.2012.06.065>.
- Pérez-Novó, C., Pateiro-Moure, M., Osorio, F., Nóvoa-Muñoz, J. C., López-Periago, E. and Arias-Estévez, M. (2008). Influence of organic matter removal on competitive and noncompetitive adsorption of copper and zinc in acid soils. *Journal of Colloid and Interface Science*, 322, 33-40. <http://doi.org/10.1016/j.jcis.2008.03.002>.
- Pinho, A. P., Matos, A. T., Costa, L. M., Morris, L. A. and Martinez, M. A. (2006). Modelagem da retenção de herbicidas em zonas ripárias. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 10, 896-902. <http://dx.doi.org/10.1590/S1415-43662006000400017>.
- Ramos, M. C. (2006). Metals in vineyard soils of the Penedès area (NE Spain) after compost application. *Journal of Environmental Management*, 78, 209-215. <http://doi.org/10.1016/j.jenvman.2005.04.017>.
- Ruyters, S., Salaets, P., Oorts, K. and Smolders, E. (2013). Copper toxicity in soils under established vineyards in Europe: a survey. *Science of the Total Environment*, 443, 470-477. <http://doi.org/10.1016/j.scitotenv.2012.11.001>.
- Sposito, G. (1989). *The chemistry of soils*. New York: Oxford University Press.
- Stüpp, J. J., Rosa, E. F. F., Amarante, C. V. T., Mafra, A. L. and Steffens, C. A. (2013). Nutrição, sanidade, rendimento e qualidade de frutos em macieiras 'Catarina' conduzidas sob manejo integrado e orgânico. *Revista Brasileira de Fruticultura*, 35, 634-641. <https://dx.doi.org/10.1590/S0100-29452013000200036>.
- Tedesco, M. J., Gianello, C., Bissani, C. A., Bohnen, H. and Volkweiss, S. J. (1995). *Análises de solo, plantas e outros materiais*. 2. ed. Porto Alegre: Departamento de Solos, UFRGS.

- Tessier, A., Campbell, P. G. C. and Bisson, M. (1979). Sequential extraction procedure for the speciation of particulate trace metals. *Analytical Chemistry*, 51, 844-851. <http://doi.org/http://10.1021/ac50043a017>.
- Toselli, M., Schiatti, P., Ara, D., Bertacchini, A. and Quartieri, M. (2009). The accumulation of copper in soils of the Italian region Emilia-Romagna. *Plant, Soil and Environment*, 55, 74-79. <https://doi.org/10.17221/317-PSE>.
- Verdisson, S., Couderchet, M. and Vernet, G. (2001). Effects of procymidone, fludioxonil and pyrimethanil on two non-target aquatic plants. *Chemosphere*, 44, 467-474. [https://doi.org/10.1016/S0045-6535\(00\)00468-9](https://doi.org/10.1016/S0045-6535(00)00468-9).
- Wang, S. and Mulligan, C. N. (2013). Effects of three low-molecular-weight organic acids (LMWOAs) and pH on the mobilization of arsenic and heavy metals (Cu, Pb, and Zn) from mine tailings. *Environmental Geochemistry and Health*, 35, 111-118. <http://doi.org/10.1007/s10653-012-9461-3>.
- Yadav, S. K. (2010). Heavy metals toxicity in plants: an overview on the role of glutathione and phytochelatins in heavy metal stress tolerance of plants. *South African Journal of Botany*, 76, 167-179. <http://doi.org/10.1016/j.sajb.2009.10.007>.
- Yuan, S., Xi, Z., Jiang, Y., Wan, J., Wu, C., Zheng, Z. and Lu, X. (2007). Desorption of copper and cadmium from soils enhanced by organic acids. *Chemosphere*, 68, 1289-1297. <http://doi.org/10.1016/j.chemosphere.2007.01.046>.
- Zhang, X., Wang, X., Wei, D., Li, B., Ma, Y. and Huang, Z. (2013). The influence of soil solution properties on phytotoxicity of soil soluble copper in a wide range of soils. *Geoderma*, 211-212, 1-7. <http://doi.org/10.1016/j.geoderma.2013.07.001>.