



Boletín de la Sociedad Geológica Mexicana

ISSN: 1405-3322

Sociedad Geológica Mexicana, A.C.

Rivera-Sylva, Héctor E.; Frey, Eberhard; Stinnesbeck,
Wolfgang; Amezcua, Natalia; Flores Huerta, Diana
First occurrence of Parksosauridae in Mexico, from the Cerro del Pueblo
Formation (Late Cretaceous; late Campanian) at Las Águilas, Coahuila
Boletín de la Sociedad Geológica Mexicana, vol. 70, no. 3, 2018, pp. 779-785
Sociedad Geológica Mexicana, A.C.

DOI: <https://doi.org/10.18268/BSGM2018v70n3a10>

Available in: <https://www.redalyc.org/articulo.oa?id=94358881010>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Short note

First occurrence of Parksosauridae in Mexico, from the Cerro del Pueblo Formation (Late Cretaceous; late Campanian) at Las Águilas, Coahuila

Héctor E. Rivera-Sylva, Eberhard Frey, Wolfgang Stinnesbeck, Natalia Amezcua, Diana Flores Huerta

Héctor E. Rivera-Sylva*hrivera@museodeldesierto.org*

Departamento de Paleontología, Museo del Desierto, Carlos Abedrop Dávila 3745, 25022, Saltillo, Coah., México.

Eberhard Frey

Staatliches Museum für Naturkunde Karlsruhe, Geowissenschaftliche Abteilung, Erbprinzenstraße 13, D76133 Karlsruhe, Germany.

Wolfgang Stinnesbeck

Institut für Geowissenschaften, Universität Heidelberg, Im Neunheimer Feld 234-236, 69120 Heidelberg, Germany.

Natalia Amezcua**Diana Flores Huerta**

Servicio Geológico Mexicano, Blvd. Felipe Ángeles km 9.5-4, Pachuca, Hgo., México.

ABSTRACT

The Las Águilas locality near Porvenir de Jalpa, Coahuila, México, is extremely rich in fossil remains comprising both bones and trackways of several late Campanian dinosaur taxa. Here we present a tooth with an asymmetrically bulged crown and an isolated shallow amphicoelous vertebral centrum. Both are assigned by us to Parksosauridae. The material represents the first occurrence of this enigmatic family in México.

Keywords: Cerro del Pueblo Formation, Parksosauridae, Campanian, Coahuila, México.

RESUMEN

La localidad de Las Águilas cerca de Porvenir de Jalpa, Coahuila, México, es extremadamente rica en restos de fósiles dinosaurios comprendiendo huesos y huellas de varios taxa Campanianos. Aquí presentamos el registro de un diente con una corona bulbosa y asimétrica además de un centrum de una vértebra anficélica. Ambos asignados por nosotros como Parksosauridae. El material representa el primer registro de esta enigmática familia para México.

Palabras clave: Formación Cerro del Pueblo, Parksosauridae, Campaniano, Coahuila, México.

BOL. SOC. GEOL. MEX. 2018
VOL. 70 NO. 3
P. 779 – 784
<http://dx.doi.org/10.18268/BSGM2018v70n3a10>

Manuscript received: January 30, 2018.
Corrected manuscript received: February 13, 2018.
Manuscript accepted: February 18, 2018.

1. Introduction

Parksosauridae are a little understood family of small ornithischian dinosaurs, with a controversial phylogenetic relationship within Ornithopoda (Bucholtz, 2002; Boyd, 2015). As its skull and dentition appear to be primitive within Ornithopoda, much resembling hypsilophodontids (Galton, 1974a; Boyd, 2014), the family it has been grouped either as “hypsilophodontid” (Sereno, 1998), a member of the Thescelosauridae (*e.g.* Galton, 1973; Butler and Galton, 2008), or equivalent to that family (Boyd, 2015).

Parksosauridae are small-bodied, bipedal, cursorial, lightly-built dinosaurs, which are known from the Aptian to Maastrichtian of North America and Asia (Norman *et al.*, 2004; Boyd *et al.*, 2009). The skull is proportionally small with respect to the body, it is elongate with flat frontals and has lingually curved and labiolingually compressed leaf-shaped teeth (Galton, 1997). The enamel of the teeth crowns is crenulated and lacks serration. Parksosauridae is documented from Canada and the United States, while it was hitherto unknown in México. Here we describe the fossil remains to a basal ornithopod remains found during the spring of 2014, 4.2 km east of the Las Águilas locality, east of the hamlet Porvenir de Jalpa, and about 50 km west of Saltillo, Coahuila, north-eastern México, in rocks of the Cerro del Pueblo Formation (Figure 1).

Institutional Abbreviations: **CPC-** Colección Paleontológica de Coahuila, Saltillo, Coahuila, México; **CNA-** Consejo Nacional de Arqueología, México; **IEUH-** Institute of Earth Sciences, Heidelberg University, Germany; **INAH-** Instituto Nacional de Antropología e Historia; **MUDE-** Museo del Desierto, Saltillo, Coahuila, México; **SMNK-** State Museum of Natural History, Karlsruhe, Germany; **SGM-** Servicio Geológico Mexicano, Pachuca, México.

2. Materials and Methods

The geology, stratigraphy, and paleoecology of the Las Águilas section have recently been presented by Vogt *et al.* (2016) and Rivera-Sylva *et al.* (2017). The locality with remains of Parksosauridae, with the field number “Las Águilas 14” (LA 14), is a new site located about 1.5 km west of Cerro de Angostura. At LA 14, abundant surface material has weathered out of a fine-grained, green-to buff-colored siltstone.

The fossils documented here at LA14 were collected during an extensive surface survey in 2014 by a joint team from MUDE, SMNK, IEUH, and SGM. They are housed and cataloged at the Museo del Desierto (MUDE). Locality information is on file at that institution. The specimens were photographed using an Olympus E 620 with an Olympus Zuiko standard lens (14 – 42 mm, f/3.5 – 5.6), an Olympus Zuiko macro lens (35 mm, f/3.5); and a Canon EOS Rebel T2i with a Canon lens 18 – 55mm, 1:3.5 – 5.6 IS II.

The specimens reported here were collected with permission of the INAH through its Consejo Nacional de Arqueología [CNA; Oficio 401.B(4)19.2013/36/1862].

3. Systematic paleontology

Ornithischia Seeley, 1888

Cerapoda Sereno, 1986

Parksosauridae Bucholtz, 2002

Parksosauridae indet.

Referred specimens – CPC 1872 (Figure 2, A-B), a right premaxillary tooth; CPC 1866 (Figure 2, C-H), a posterior sacral vertebral centrum.

Locality – LA 14 near Las Águilas, Porvenir de Jalpa, Coahuila, northeast México (Figure 1).

Description – CPC 1872 (Figure 2, A-B) consists of a premaxillary tooth crown, slightly labi-



Figure 1 Geographic location of the Porvenir de Jalpa, Las Águilas fossil site in Coahuila, north-east México.

olingually compressed and slightly constricted at its base. Apically to this constriction, the crown is asymmetrically bulged. The bulge is twice as pronounced distally compared to mesially, and only misses the labial face of the crown. The tip of the crown is blunt. The tooth crown is slightly curved lingually and bears a small wear facet on its tip, showing a few coarse labial striations, which display a high degree of wear. The absence of the root suggests a shed tooth (Frey and Monninger, 2010). The crown height is 5 mm, with a basal maximum diameter of 3 mm.

CPC 1866 (Figure 2, C-H) is a posterior sacral vertebral centrum, interpreted as the fifth sacral, which is shallowly amphicoelous and constricted around its middle. The bone is 1.5 times longer

and slightly higher than wide. In anterior and posterior views, the centrum is rounded triangular in cranial outline and has a concave area separated by a dorso-medial ridge, with a flat dorsal margin and slightly convex lateral margins. The dorsal surface is ornamented with deep transverse striae that accommodate the peduncles of the neural arch. The floor of the neural canal is hourglass shaped in dorsal aspect. It is 48 mm long, 42 mm deep and 31 mm wide.

4. Discussion

The shape of the premaxillary tooth face, including the blunt tip, the striations on the lingual side,



Figure 2 Parksosauridae fossils from the Cerro del Pueblo Formation. A right premaxillary tooth (CPC 1872) in: (A) labial view and, (B) lingual view. Fifth sacral vertebra (CPC 1866) in: (C) left lateral view; (D) right lateral view; (E) ventral view; (F) dorsal view; (G) cranial view; (H) caudal view. Scale in A-B: 5 mm; C-H: 5 cm.

the smooth lingual surface, the constricted base, and the lingually curving blunt crown, are seen in *Thescelosaurus* and *Hypsilophodon* (Galton, 1974b; Boyd, 2014). In *Parksosaurus*, the premaxilla is missing (Parks, 1926; Sternberg, 1940), thus differing from other ornithischians with leaf-shape teeth (*i.e.*, pachycephalosaurids). Serrations are absent on both the anterior and posterior margins, which also is the case in neoornithischians *Changchunsaurus*, *Haya*, and *Jeholosaurus* (Barrett and Han, 2009; Jin *et al.*, 2010; Makovicky *et al.*, 2011).

The sacral vertebrae of *Parksosaurus*, *Thescelosaurus*, and *Dysalotosaurus* are very similar to those reported for *Hypsilophodon* (Galton, 1974b). The constricted centrum across their middle circumference and the shallow platycoelous intervertebral joints are features seen only in this group (Gilmore, 1915; Brown *et al.*, 2013). The widened centrum, in cranial view, towards the caudal face, and ventrally longitudinally concavity, identifies this vertebra as the posterior-most sacral element of the series (sa. 5) (Galton, 1974b) it much resembles vertebrae illustrated by Carr and Seitz (2015; p. 107) for

Thescelosaurus, and Galton (1974b) for *Hypsilophodon*. The neural arch is not preserved and may have fallen off during diagenesis. The striae on the centropeduncular articulation indicate that the neurocentral suture must have been open, which is suggestive for an immature individual (Brochu, 1996; Irmis, 2007), contrasting with the vertebrae of adult pachycephalosaurids, which show strong sutural fusion (Gilmore, 1924; Sues and Galton, 1987).

The only well-known basal cerapoda taxon from the late Campanian of North America, *Orodromeus*, occurs in the Two Medicine Formation (Montana). These osteological remains resemble an unnamed basal ornithopod from the Kaiparowits Formation Utah; Gates *et al.*, 2013.

5. Conclusions

The material described here represents the first evidence of Parksosauridae in México. This discovery from the northeastern Mexican state of

Coahuila extends the geographic range of Parksosauridae for about 2000 kilometers to the south of North America.

Acknowledgements

We wish to thank the following friends and colleagues for their participation in fieldwork in the Las Águilas area: Lucía Alfaro (Saltillo), Felix Bayer (Heidelberg), Christiane Birnbaum (Karlsruhe), Kristina Eck (Heidelberg), Aaron Linsler (Heidelberg), Tim Niggemeyer (Karlsruhe), Melissa Perner (Heidelberg), Ulli Karas (Siegburg), Sarah Stinnesbeck (Karlsruhe), Eva Susanne Stinnesbeck (Troisdorf), Manfred Vogt (Heidelberg), and Sven Wetzel (Heidelberg). The Museo del Desierto in Saltillo provided the infrastructure for the fieldwork of HR, EF, and WS. Rubén Guzmán-Gutierrez (Aguascalientes) is gratefully acknowledged for providing literature. The Servicio Geológico Mexicano supported field research for NA and DFH. Iván Erick Sanchez Uribe is thanked for image editing. Kenneth Carpenter and Rodolfo Coria are thanked for their reviews and useful comments. The Deutsche Forschungsgemeinschaft (DFG) covered travel and fieldwork expenses for EF (FR1314/19) and WS (STI128/24 and 44).

References

- Barrett, P.M., Han, F.-L., 2009, Cranial anatomy of *Jeholosaurus shangyuanensis* (Dinosauria: Ornithischia) from the Early Cretaceous of China: *Zootaxa*, 2072, 31–55.
- Boyd, C.A., 2014, The cranial anatomy of the neornithischian dinosaur *Thescelosaurus neglectus* (en linea): PeerJ, e779, disponible en: <<https://doi.org/10.7717/peerj.669>>.
- Boyd, C.A., 2015, The systematic relationships and biogeographic history of ornithischian dinosaurs (en linea): PeerJ, e1523, disponible en: <<https://doi.org/10.7717/peerj.1523>>.
- Boyd C.A., Brown, C.M., Scheetz, R.D., Clarke, J.A., 2009, Taxonomic revision of the basal neornithischian taxa *Thescelosaurus* and *Bugenasaura*: *Journal of Vertebrate Paleontology*, 29, 758–770.
- Brochu, C.A., 1996, Closure of neurocentral sutures during crocodilian ontogeny: implications for maturity assessment in fossil archosaurs: *Journal of Vertebrate Paleontology*, 16, 49–62.
- Brown, C.M., Evans, D.C., Ryan, M.J., Russell, A.P., 2013, New data on the diversity and abundance of small-bodied ornithopods (Dinosauria: Ornithischia) from the Belly River Group (Campanian) of Alberta: *Journal of Vertebrate Paleontology*, 33(3), 495–520.
- Bucholtz, P.W., 2002, Phylogeny and biogeography of basal Ornithischia, in Brown, D.E. (ed.) *The Mesozoic in Wyoming*: Casper, Tate Geological Museum, 18–24.
- Butler, R.J., Galton, P.M., 2008, The “dermal armour” of the ornithopod dinosaur *Hypsilophodon* from the Wealden (Early Cretaceous: Barremian) of the Isle of Wight: a reappraisal: *Cretaceous Research*, 29, 636–642.
- Carr, T.D., Seitz, M.E., 2015, *Field Guide to the Microfossils of the Hell Creek Formation*: Carthage Institute of Paleontology, 177 p.
- Frey, E., Monninger, S., 2010, Lost in action – isolated crocodilian teeth from Enspel and their interpretative value: *Palaeobiodiversity and Palaeoenvironments*, 90, 65–81.
- Galton, P.M., 1973, Redescription of the skull and mandible of *Parksosaurus* from the Late Cretaceous with comments on the

- family Hypsilophodontidae (Ornithischia): Life Sciences Contribution Royal Ontario Museum, 89, 30 p.
- Galton, P.M., 1974a, Notes on *Thescelosaurus*, a conservative ornithopod dinosaur from the Upper Cretaceous of North America, with comments on ornithopod classification: *Journal of Paleontology*, (48)5, 1048–1067.
- Galton, P.M., 1974b, The ornithischian dinosaur *Hypsilophodon* from the Wealden of the Isle of Wight: London, Bulletin of the British Museum (Natural History), Geology, 25(1), 152 p.
- Galton, P.M., 1997, Cranial anatomy of the basal hypsilophodontid dinosaur *Thescelosaurus neglectus* Gilmore (Ornithischia: Ornithopoda) from the Upper Cretaceous of North America: *Revue Paléobiologie*, 16(1), 231–258.
- Gates, T.A., Lund, E.K., Boyd, C.A., Deblieux, D.D., Titus, A.L., Evans, D.C., Getty, M.A., Kirkland, J.I., Eaton, J.G., 2013, Ornithopod Dinosaurs from the Grand Staircase-Escalante National Monument Region, Utah, and their role in paleobiogeographic and macroevolutionary studies, in Titus, A.L., Loewen, M.A. (eds.) *At the top of the Grand Staircase, the Late Cretaceous of Southern Utah*: Indiana University Press, 463–481.
- Gilmore, C.W., 1915, Osteology of *Thescelosaurus*, an ornithopodous dinosaur from the Lance Formation of Wyoming: *Proceedings of the United States National Museum*, (49) 2127, 591–616.
- Gilmore, C.W., 1924, On *Troodon validus*, an orthopodous dinosaur from the Belly River Cretaceous of Alberta, Canada: *Department of Geology Bulletin (University of Alberta)*, 1, 1–43.
- Irmis, R.B., 2007, Axial skeleton ontogeny in the parasuchia (Archosauria: Pseudosuchia) and its implications for ontogenetic determination in archosaurs: *Journal of Vertebrate Paleontology*, 27, 350–361.
- Jin L.Y., Chen, J., Zan, S.Q., Butler, R.J., Godefroit, P., 2010, Cranial anatomy of the small ornithischian dinosaur *Changchunsaurus parvus* from the quantou formation (Cretaceous: Aptian-Cenomanian) of Jilin Province, northeastern China: *Journal of Vertebrate Paleontology*, 30, 196–214.
- Makovicky, P.J., Kilbourne, B.M., Sadleir, R.W., Norell, M.A., 2011, A new basal ornithopod (Dinosauria, Ornithischia) from the Late Cretaceous of Mongolia: *Journal of Vertebrate Paleontology*, 31, 626–640.
- Norman, D.B., Sues, H.-D., Witmer, L.M., Coria, R.A., 2004, Basal Ornithopoda, in Weishampel, D.B., Dodson, P., Osmólska, H. (eds.), *The Dinosauria* (2nd Edition), Berkeley, University of California Press, 393–412.
- Parks, W.A., 1926, *Thescelosaurus warreni*, a new species of orthopodous dinosaur from the Edmonton Formation of Alberta: *University of Toronto Studies (Geological Series)*, 21, 1–42.
- Rivera-Sylva, H.E., Frey, E., Schulp, A.S., Meyer, C., Thuring, B., Stinnesbeck, W., Vanhecke, V.A.A., 2017, Late Campanian theropod trackways from Porvenir de Jalpa, Coahuila, México (en línea): *Palaeovertebrata*, 41(2)-e1, disponible en: <doi: 10.18563/pv.41.2.e1>.
- Seeley, H.G., 1888, The classification of the Dinosauria: Report of the British Association for the Advancement of Science, 1887, 698–699.
- Sereno, P., 1986, Phylogeny of the bird-hipped dinosaurs (Order Ornithischia): *National Geographic Research*, 2, 234–256.
- Sereno, P., 1998, A rationale for phylogenetic definitions, with application to the higher-level taxonomy of Dinosauria: *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 210, 41–83.

- Sternberg, C.M., 1940, *Thescelosaurus edmontonensis*, n. sp., and classification of the Hypsilophodontidae: *Journal of Paleontology*, 14(5), 481–494.
- Sues, H.-D., Galton, P.M., 1987, Anatomy and classification of the North American Pachycephalosauria (Dinosauria: Ornithischia): *Palaeontographica, Abteilung A*, 198, 1–40.
- Vogt, M., Stinnesbeck, W., Zell, P., Kober, B., Kontny, J., Herzer, N., Frey, E., Rivera-Silva, H.E., Padilla Gutierrez, J.M., Amezcua Torres, N., Flores Huerta, D., 2016, Age and depositional environment of the “dinosaur graveyard” at Las Águilas, southern Coahuila, NE México: *Palaeogeography, Palaeoclimatology, Palaeoecology*, 441(1), 758–769.