



Salud Pública de México

ISSN: 0036-3634

spm@insp.mx

Instituto Nacional de Salud Pública
México

Fleury, Agnès; Sciutto, Edda; Larralde, Carlos
Neurocysticercosis is still prevalent in Mexico
Salud Pública de México, vol. 54, núm. 6, noviembre-diciembre, 2012, pp. 632-636
Instituto Nacional de Salud Pública
Cuernavaca, México

Available in: <http://www.redalyc.org/articulo.oa?id=10625052009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Neurocysticercosis is still prevalent in Mexico

Agnès Fleury, MD, PhD,^(1,2) Edda Sciutto, PhD,⁽¹⁾ Carlos Larralde MD, PhD.⁽¹⁾

Fleury A, Sciutto E, Larralde C.
Neurocysticercosis is still prevalent in Mexico.
Salud Publica Mex 2012;54:632-636.

Abstract

In this work, we report the published cases of human and porcine cysticercosis, as well as *Taenia solium* taeniasis diagnosed in Mexico during the last 10 years. Numerical data allow us to state that this disease remains as a public health problem in our country. Whereas efficient tools have been developed for the diagnosis and prevention of cysticercosis, we strongly recommend further measures allowing the control and eventual eradication of this parasite in Mexico.

Key words: epidemiology; *Taenia solium*; neurocysticercosis; Mexico

Fleury A, Sciutto E, Larralde C.
La neurocisticercosis es aún prevalente en México.
Salud Publica Mex 2012;54:632-636.

Resumen

En este trabajo reportamos los casos publicados de cisticercosis humana y porcina, así como de teniosis por *Taenia solium* diagnosticados en México en los últimos 10 años. Los datos numéricos nos permiten sostener que esta parasitosis continúa siendo un problema de salud pública en nuestro país. Considerando que se han desarrollado herramientas eficientes para el diagnóstico y la prevención de la cisticercosis, recomendamos enfáticamente seguir promoviendo las medidas que permitan el control y posterior erradicación de esta parasitosis en México.

Palabras clave: neurocisticercosis; epidemiología; México

Taeniasis and cysticercosis are endemic in most of Latin America, Asia and Africa countries.¹ However, the burden of this endemic disease is not easy to appraise due to some characteristics of the infection, namely: 1) human infection with the adult form of the parasite (taeniasis) is mostly asymptomatic; 2) a large proportion of the humans infected by the larvae (cysticercus) are also asymptomatic; 3) taeniasis diagnosis relies on laboratory techniques not routinely made, and 4)

accurate diagnosis of human neurocysticercosis (NC) requires costly neuro-radiological studies not available to the whole population at risk. The scarcity and/or uncertainty of the epidemiological data on taeniasis and cysticercosis have raised some debate and led to unsound conclusions.

Herein, we review and summarize the published studies reporting *Taenia solium* infections in humans and pigs in Mexico during the last decade.

(1) Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México. México DF, México.

(2) Instituto Nacional de Neurología y Neurocirugía (INNN). México DF, México.

Received on: November 28, 2011 • Accepted on: July 24, 2012

Corresponding author: Agnès Fleury. Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México. Insurgentes Sur 3877, Col. La Fama. 14269 Tlalpan, México DF, México.
E-mail: afleury@biomedicas.unam.mx

Methodology

In this study, a PubMed revision of the papers published in the last ten years containing data on *Taenia solium* taeniasis and on human or porcine cysticercosis in Mexico is presented. Different combinations of the following key words were used: "*Taenia solium*, cysticercosis, epidemiology, epilepsy, Mexico". In this review we included only those publications in which: 1) the diagnostic criteria were clearly defined; 2) NC diagnosis was made between 2000 and 2010, or, (since in many publications date of diagnosis is not cited) either vesicular (viable) or colloidal (inflammatory) parasites were included between 2000 and 2010. The articles published within these years but without information for dating the diagnosis of human or porcine cysticercosis or for dating the period of inclusion of the patients and the evolution stage of the parasites were not included.

Results

Table I presents the results found in the PubMed database.²⁻¹⁹ As can be seen, several studies demonstrate the persistence of human and porcine *T. solium* cysticercosis in Mexico. We did not find any publication about the diagnosis of intestinal *T. solium* tapeworm carriers.

Discussion

There is no doubt that epidemiological transition is ongoing in Mexico, as shown by the increasing incidence of metabolic, neoplastic and degenerative diseases.²⁰ Nonetheless, this trend gives no good reason for disregarding the persistence of infectious diseases in Mexico, mainly those linked with poverty.

Data presented in table I illustrates that the *Taenia solium* life-cycle is still present in Mexico: up to 32 % of porcine cysticercosis prevalence in some communities and about 500 cases of human NC were diagnosed and reported in the last ten years. Although it is evident that these data are incomplete and do not represent the real burden of human NC and porcine cysticercosis in Mexico, since only a small proportion of the cysticercotic cases are published, these data are sufficient evidence that *Taenia solium* is still a public health problem in Mexico.

In one recent paper, mainly based on the results of official statistics on human NC morbidity, the opposite conclusion was reached.²¹ The divergence between the evidence herein shown and those published could be due to differences in the accuracy of NC diagnosis. Most of the studies shown in table I came from medi-

cal institutions in which NC diagnosis was made using modern radiological technology; in contrast, no information in this respect is given in official statistics, and probably, diagnostic is performed by procedures not equally qualified. Differences in the reliability of the data source could also be involved. According to the official statistics, approximately 400 new NC cases were reported throughout Mexico in 2004. In clear contrast, 120 new NC cases were reported in only one neurological institution (INNN) this same years.¹² It appears unlikely that a single institution concentrated 25% of all national NC cases in one year, especially considering the huge differences in sample sizes. Indeed, official statistics come from the entire population of Mexico (circa 112×10^6 inhabitants), while INNN data include only few thousands of Mexican adults. Such a disproportion points to a possible significant under-registration of NC in the nationwide sample, which occurs even though NC notification is mandatory.²² The clinical heterogeneity of NC, and the lack of access by a large part of the Mexican population to neuroradiological studies, indispensable for accurate NC diagnosis, could contribute to the nationwide NC under-diagnosis. Such hypothesis is further strengthened by the results of the two CT scan epidemiological studies performed in rural communities of central Mexico, which reported that 9 to 10 % of apparently healthy residents had cranial CT scan images compatible with NC^{15,16} (table I).

Some considerations about the appropriateness of whether considering a parasitic disease as a "public health problem" or not have to be mentioned. "Public Health" is "the science and art of preventing disease, prolonging life and promoting health through the organized efforts and informed choices of society, organizations, public and private, communities and individuals".²³ Based on this definition, a public health problem can be defined as a problem which occurs frequently and widely, causes severe disability and suffering, may be effectively dealt with, and treated by ways and means acceptable to patients, their families, and society. Undoubtedly, NC satisfies the three latter points, as it may cause severe disability and suffering,^{24,25} there are effective methods for dealing with it,^{19,26-28} and current treatments are acceptable to all involved.²⁹ The issue here is the first point: ¿What is the frequency of NC in Mexico? Data in table I does not answer the question with an acceptable level of precision, but this question can be substituted by: What is the number of NC cases which public health authorities may consider as an "acceptable" proof that NC still merits their attention? In this respect, table I confirms that the absolute number of NC cases is costly in terms of suffering for stricken

Table I
EPIDEMIOLOGICAL STUDIES ON TAENIASIS/CYSTICERCOSIS IN MEXICO

Reference	Place of the study	Type of study	Diagnosis based on (serology, radiology)	Number of subject included	Number of cases
Human studies					
Hospital cases					
Cohn-Zurita <i>et al.</i> , 2006 ²	México DF (CMN Siglo XXI)	Case-report	MRI*	NA [‡]	1
Chavarria <i>et al.</i> , 2006 ³	México DF, (INNN)	Case-series; diagnosed between 2000-2003	MRI	NA	23 ^{&}
Ortiz-Trejo <i>et al.</i> , 2006 ⁴	Zacatecas (Public general Hospital)	Case-series; diagnosed between January and September 2004	CT scan	NA	85
Alvarado-Esquivel <i>et al.</i> , 2008 ⁵	Durango (Psychiatric Hospital)	Sample of psychiatric patients	Serology, ELISA and WB	105	8 (ELISA) 1 (EITB)
Jimenez-Vazquez and Nagore, 2008 ⁶	Morelia, Michoacan (Public general Hospital)	Case-report of vesicular NC	MRI	NA	1
Jimenez-Vazquez and Nagore, 2008 ⁷	Morelia, Michoacan (Public general Hospital)	Case report of vesicular NC	MRI	NA	1
Rio de la Loza and Meza, 2008 ⁸	Hermosillo, Sonora	Case-report of colloidal parasites	MRI	NA	1
Gongora <i>et al.</i> , 2008 ⁹	México DF, (INNN)	Case-series of vesicular parasites included between 2000-2003	MRI	NA	31 ^{&}
Torres-Corzo <i>et al.</i> , 2009 ¹⁰	San-Luis Potosi (Central Hospital)	Case-series of neuro-endoscopies made between 1/2003 and 6/2006	MRI	NA	25
Suastegui <i>et al.</i> , 2009 ¹¹	10 different Mexican epileptic centers	Samples of late-onset epileptic patients (debut in 2000 or after)	CT scan	455	96
Fleury <i>et al.</i> , 2010 ¹²	México DF, (INNN)	All NC patients diagnosed at INNN in 2004	CT scan +/- MRI	4706	120
Cárdenas <i>et al.</i> , 2011 ¹³	México DF, (INNN)	Case-report	MRI	NA	1
Cárdenas <i>et al.</i> , 2012 ¹⁴	México DF, (INNN)	Case-series; diagnosed between 8/2005 and 10/2007	MRI	NA	50
Population based studies					
Fleury <i>et al.</i> , 2003 ¹⁵	Tepetzintla, Puebla	Population-based, made in 2001	CT scan	154	14
Fleury <i>et al.</i> , 2006 ¹⁶	Cuentepec, Morelos	Population-based, made in 2003	CT scan	649	59
Swine-cysticercosis studies					
Morales <i>et al.</i> , 2002 ¹⁷	Cuentepec, Morelos	Sample of pigs	Tongue inspection	1300	429
Morales <i>et al.</i> , 2006 ¹⁸	State of Morelos	Sample of pigs	Tongue inspection	1747	249
Morales <i>et al.</i> , 2008 ¹⁹	Sierra de Huautla, Morelos	Sample of pigs	Tongue inspection	562	75

* Magnetic Resonance Imaging

‡ Not appropriate

& It is possible that some patients were included in these two studies

individuals and of resources for public health institutions in Mexico. On this same matter, it is interesting to note the attitude taken by the authors of a recent paper, in which the 78 NC cases acquired inside the United States between 1954 and 2005 were reported.³⁰ The authors comment: "Several factors, including the severe, potentially fatal, nature of cysticercosis; its fecal-oral route of transmission; the considerable economic effect; the availability of a sensitive and specific serologic test for infection by adult *Taenia solium* tapeworms; and the

demonstrated ability to find a probable source of infection among contacts, all provide a compelling rationale for implementation of public health control efforts". The differences between these US authors' proposal with some Mexican voices claiming "NC is no longer a public health problem in Mexico"²¹ are striking, as the number of NC cases in Mexico is much greater than in this country.

It must also be noted that, as shown in table I, the social conditions allowing the parasite to complete its

life cycle persist in rural areas of Mexico, and porcine cysticercosis is prevalent in such areas,^{19,31} while, to our knowledge, no cases of porcine cysticercosis have been reported in the United States of America.

Finally, a recent report from the WHO Expert Consultation on Foodborne Trematode Infections and Taeniasis/Cysticercosis³² is in accord with our view, stating: "Cysticercosis is endemic in much of Latin America, with "hot spots" of the disease in Mexico (e.g. Yucatan, Guanajuato, Guerrero, Morelos, Puebla), several Central American countries, such as Guatemala, Honduras and Nicaragua, the Andean countries of Bolivia, Ecuador and Peru (e.g. Tumbes), Colombia, Venezuela and northern Brazil", and also that "Neurocysticercosis is a significant public health problem in the entire region, while porcine cysticercosis is a constraint to food security and a major cause of income loss".

NC, as other infectious diseases, is still an embarrassing health problem of Mexico since it is clearly related to the poverty prevailing in under-developed countries.

It is not gratuitous to reflect more deeply on the possible causes of the persistence of *Taenia solium* infection in Mexico. As it is well known, completion of the life cycle of *Taenia solium* is clearly related with under-development, as it requires of outdoor defecation and free roaming pigs. As reported by Flisser & Correa,²¹ global indices of Mexican national development have improved in recent years, posing the question as to why does *T. solium* transmission cycle still prevail in Mexico. This is probably due to the fact that global indices of national development are not appropriate descriptors of all sectors in Mexico, where extreme social and economic disparities subsist. As table II³³ shows, Mexican population income is quite disparate among deciles. Indeed, total current income can be divided in three parts, one of them corresponding to 10% of the population in Mexico, the richest one; the other one corresponding to 30% of the population, the middle class, and the third one to 60% of the population, the poor in Mexico. Such

an unequal distribution of the national income lasts since 2002 to 2008 at least, without a significant tendency to improve with time. The social disparity among Mexican inhabitants is also clear when comparing the income in rural community population (less than 2 500 inhabitants) versus the income of urban population (more than 2 500 inhabitants). As shown in table III,³³ mean income per person grew only by 1.4% in rural areas whilst 10.8% in urban areas between 2002 and 2008. These data clearly illustrate that, although global socio-economic indices of Mexico show a positive tendency within these years, a significant part of its population (60%) –the poor and the rural of Mexico– are not included in the mentioned improvement, and it is in these areas where *Taenia solium* life-cycle finds conditions to persist.

Conclusion

All these data and arguments indicate to the persistence of *Taenia solium* transmission in Mexico at levels which are—to our mind—still too high. Consequently, Mexican health authorities must continue to consider NC as a public health problem, and the available measures for its control have to be applied in order to prevent and control this disease.

Acknowledgement

This article was partly supported by Consejo Nacional de Ciencia y Tecnología (CONACYT), project number 2008-86527, 106154, and by CYTED program (Red 211RT0432). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Declaration of conflict of interests. The authors declare that they have no conflict of interests.

Table II
DISTRIBUTION OF THE TOTAL CURRENT INCOME
(THREE MONTHS' AVERAGE) IN DECILES OF HOUSEHOLDS
(PERCENTAGE)¹⁸

Deciles	2002	2004	2006	2008
Deciles I to VI	27.0	26.9	27.6	26.7
Deciles VII to IX	37.4	36.9	36.7	37.0
Decile X	35.6	36.2	35.7	36.3
Total	100.0	100.0	100.0	100.0

Table III
TOTAL CURRENT INCOME (THREE MONTHS' AVERAGE)
PER HOUSEHOLD AND ITS VARIATION, ACCORDING
TO COMMUNITY SIZE, 2002-2008¹⁸

Year	Size of the community			
	With 2 500 or more inhabitants	Average	With less than 2 500 inhabitants	Average
	Households	Income*	Households	Income*
2002	18 759 449	37 312	5 772 182	18 019
2004	19 809 869	38 431	5 751 578	18 171
2006	20 685 257	41 677	5 856 070	21 834
2008	21 345 884	41 341	5 386 710	18 276

* Mexican pesos

References

1. Sotelo J. Clinical manifestations, diagnosis, and treatment of neurocysticercosis. *Curr Neurol Neurosci Rep* 2011;11:529-535.
2. Cohn-Zurita F, Guinto-Balazar G, Pérez-Cerdán H. Neurocysticercosis associated with pituitary adenoma. Case report and literature review. *Cir Cir.* 2006;74:47-49.
3. Chavarria A, Fleury A, Bobes RJ, Morales J, Fragoso G, Sciutto E. A depressed peripheral cellular immune response is related to symptomatic neurocysticercosis. *Microbes Infect* 2006;8:1082-1089.
4. Ortiz-Trejo JM, Correa-Chacón AJ, Sctelo-Ham EI, Torres-Valenzuela A, Alvarado-Esquivel C. Risk factors associated with neurocysticercosis in a public hospital in Mexico. *Gac Med Mex* 2006;142:175-179.
5. Alvarado-Esquivel C, Arreola-Valenzuela MA, Rodríguez-Briones A, Alanís-Quinones OP, Estrada-Martínez S, Luevanos-Becerra C, et al. Seroprevalence of selected viral, bacterial and parasitic infections among inpatients of a public psychiatric hospital of Mexico. *Rev Inst Med Trop Sao Paulo* 2008;50:161-164.
6. Jimenez-Vazquez OH, Nagore N. Cisternal neurocysticercosis. *Br J Neurosurg* 2008;22:774-775.
7. Jimenez-Vazquez OH, Nagore N. Role of neuroendoscopy in the treatment of large viable cysticerci in the brain parenchyma. *Br J Neurosurg* 2008;22:682-683.
8. Río de la Loza LJ, Meza EL. Cysticercotic encephalitis: case report of miliary infestation in an encephalopathic fashion. *Arch Neurol* 2008;65:276-277.
9. Góngora-Rivera F, Soto-Hernández JL, Guevara P, Sotelo-Morales J. In neurocysticercosis, CSF cytokines correlate with cerebral blood flow velocities. *Neurology* 2008;71:1119-1122.
10. Torres-Corzo JG, Tapia-Pérez JH, Sánchez-Aguilar M, Della Vecchia RR, Chalita Williams JC, Cerda-Gutiérrez R. Comparison of cerebrospinal fluid obtained by ventricular endoscopy and by lumbar puncture in patients with hydrocephalus secondary to neurocysticercosis. *Surg Neurol* 2009;71:376-379.
11. Suástegui R, Gutiérrez J, Ramos R, Bouchan S, Navarrete H, Ruiz J, et al. Clinical characteristics of the late-onset epilepsy in Mexico to the beginning of the new millennium: 455 cases. *Rev Invest Clin* 2009;61:354-363.
12. Fleury A, Moreno-García J, Valdez-Aguerreberre P, de Sayve-Durán M, Becerril-Rodríguez P, Larralde C, et al. Neurocysticercosis, a persisting health problem in Mexico. *PLoS Negl Trop Dis* 2010;4:e805.
13. Cardenas G, Bahena A, Soto-Hernandez JL, Fleury A. A dramatic case of intraventricular cysticercosis. *Arch Neurol* 2011;68:828-829.
14. Cárdenas G, Valdez R, Sáenz B, Bottasso O, Fragoso G, Sciutto E, et al. Impact of *Taenia solium* neurocysticercosis upon endocrine status and its relation with immuno-inflammatory parameters. *Int J Parasitol* 2012;42:171-176.
15. Fleury A, Gómez T, Alvarez I, Meza D, Huerta M, Chavarria A, et al. High prevalence of calcified silent neurocysticercosis in a rural village of Mexico. *Neuroepidemiology* 2003;22:139-145.
16. Fleury A, Morales J, Bobes RJ, Dumas M, Yáñez O, Piña J, et al. An Epidemiological study of familial neurocysticercosis in an endemic Mexican community. *Trans R Soc Trop Med Hyg* 2006;100: 551-558.
17. Morales J, Velasco T, Tovar V, Fragoso G, Fleury A, Beltrán C, et al. Castration and pregnancy of rural pigs significantly increase the prevalence of naturally acquired *Taenia solium* cysticercosis. *Vet Parasitol* 2002;108:41-48.
18. Morales J, Martínez JJ, García-Castella J, Peña N, Maza V, Villalobos N, et al. *Taenia solium*: the complex interactions of biological, social, geographical and commercial factors, involved in the transmission dynamics of pig cysticercosis in highly endemic areas. *Ann Trop Med Parasitol* 2006;100:123-135.
19. Morales J, Martínez JJ, Manoutcharian K, Hernández M, Fleury A, Gevorkian G, et al. Inexpensive anti-cysticercosis vaccine: S3Pvac expressed in heat inactivated M13 filamentous phage proves effective against naturally acquired *Taenia solium* porcine cysticercosis. *Vaccine* 2008;26: 2899-2905.
20. Stevens G, Dias RH, Thomas KJ, Rivera JA, Carvalho N, Barquera S, et al. Characterizing the epidemiological transition in Mexico: national and subnational burden of diseases, injuries, and risk factors. *PLoS Med* 2008;5:e125.
21. Flisser A, Correa D. Neurocysticercosis may no longer be a public health problem in Mexico. *PLoS Negl Trop Dis* 2010;4: e831.
22. Secretaría de Salud. Modificación a la Norma Oficial Mexicana NOM-021-SSA2-1994, para la prevención y control del complejo teniosis/cisticercosis en el primer nivel de atención médica. Diario Oficial de la Federación, 2004. [consultado 2011 agosto]. Disponible en: <http://www.salud.gob.mx/unidades/cdi/nom/m021ssa294.html>.
23. Winslow CE A. The Untilled Fields of Public Health.Science 1920;51:23.
24. Fleury A, Carrillo-Mezo R, Flisser A, Sciutto E, Corona T. Subarachnoid basal neurocysticercosis: a focus on the most severe form of the disease. *Expert Rev Anti infect Ther* 2011;9:123-133.
25. Bhattarai R, Budke CM, Carabin H, Proaño JV, Flores-Rivera J, Corona T, et al. Quality of Life in Patients with Neurocysticercosis in Mexico. *Am J Trop Med Hyg* 2011;84:782-786.
26. Del Brutto OH, Roos KL, Coffey CS, García HH. Meta-analysis: Cysticidal drugs for neurocysticercosis: albendazole and praziquantel. *Ann Intern Med* 2006;145:43-51.
27. Assana E, Kyngdon CT, Gauci CG, Geerts S, Dorny P, De Deken R, et al. Elimination of *Taenia solium* transmission to pigs in a field trial of the TSOL18 vaccine in Cameroon. *Int J Parasitol* 2010;40:515-519.
28. Cai X, Yuan G, Zheng Y, Luo X, Zhang S, Ding J, et al. Effective production and purification of the glycosylated TSOL18 antigen, which is protective against pig cysticercosis. *Infect Immun* 2008;76:767-770.
29. Jung H, Cárdenas G, Sciutto E, Fleury A. Medical Treatment for Neurocysticercosis: Drugs, Indications and Perspectives. *Curr Top Med Chem* 2008;8:424-433.
30. Sorvillo F, Wilkins P, Shafir S, Eberhard M. Public Health Implications of Cysticercosis Acquired in The United States. *Emerg Infect Dis* 2011;17:1-6.
31. Morales J, De Aluja A, Martínez JJ, Hernández M, Rosas G, Villalobos N, et al. Recombinant S3Pvac-phage anticysticercosis vaccine; Simultaneous protection against cysticercosis and hydatid disease in rural pigs. *Vet Parasitol* 2011;176: 53-58.
32. World Health Organization. WHO Expert Consultation on Food-borne Trematode Infections and Taeniasis/Cysticercosis. 2011, Geneva, Switzerland: WHO, 2011: 59.
33. Instituto Nacional de Estadística y Geografía. Encuesta Nacional de Ingresos y Gastos de los Hogares (ENIGH). 2008 [consultado 2011 agosto]. Disponible en: <http://www.inegi.org.mx>