



Salud Pública de México

ISSN: 0036-3634

spm@insp.mx

Instituto Nacional de Salud Pública
México

Denova-Gutiérrez, Edgar; Tucker, Katherine L; Salmerón, Jorge; Flores, Mario; Barquera, Simón

Relative validity of a food frequency questionnaire to identify dietary patterns in an adult Mexican population

Salud Pública de México, vol. 58, núm. 6, noviembre-diciembre, 2016, pp. 608-616

Instituto Nacional de Salud Pública
Cuernavaca, México

Available in: <http://www.redalyc.org/articulo.oa?id=10649652016>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Relative validity of a food frequency questionnaire to identify dietary patterns in an adult Mexican population

Edgar Denova-Gutiérrez, PhD,⁽¹⁾ Katherine L Tucker, PhD,⁽²⁾ Jorge Salmerón, PhD,^(3,4)
Mario Flores, PhD,⁽⁵⁾ Simón Barquera, PhD.⁽⁵⁾

Denova-Gutiérrez E, Tucker KL, Salmerón J, Flores M, Barquera S. Relative validity of a food frequency questionnaire to identify dietary patterns in an adult Mexican population. *Salud Publica Mex* 2016;58:608-616. <http://dx.doi.org/10.21149/spm.v58i6.7842>

Denova-Gutiérrez E, Tucker KL, Salmerón J, Flores M, Barquera S. Validez relativa del cuestionario de frecuencia de alimentos para identificar patrones dietarios en adultos Mexicanos. *Salud Publica Mex* 2016;58:608-616. <http://dx.doi.org/10.21149/spm.v58i6.7842>

Abstract

Objective. To examine the validity of a semi-quantitative food frequency questionnaire (SFFQ) to identify dietary patterns in an adult Mexican population. **Materials and methods.** A 140-item SFFQ and two 24-hour dietary recalls (24DRs) were administered. Foods were categorized into 29 food groups used to derive dietary patterns via factor analysis. Pearson and intraclass correlations coefficients between dietary pattern scores identified from the SFFQ and 24DRs were assessed. **Results.** Pattern 1 was high in snacks, fast food, soft drinks, processed meats and refined grains; pattern 2 was high in fresh vegetables, fresh fruits, and dairy products; and pattern 3 was high in legumes, eggs, sweetened foods and sugars. Pearson correlation coefficients between the SFFQ and the 24DRs for these patterns were 0.66 ($P<0.001$), 0.41 ($P<0.001$) and 0.29 ($P=0.193$) respectively. **Conclusions.** Our data indicate reasonable validity of the SFFQ, using factor analysis, to derive major dietary patterns in comparison with two 24DR.

Keywords: questionnaire; survey; dietary assessment; adults; Mexico

Resumen

Objetivo. Evaluar la validez de un cuestionario de frecuencia de consumo de alimentos (CFA) para derivar patrones dietarios en población adulta mexicana. **Material y métodos.** Un CFA de 140-alimentos y dos recordatorios de 24-horas (24DRs) fueron obtenidos. Los alimentos fueron categorizados en 29 grupos para derivar los patrones dietarios mediante análisis factorial. La validez se evaluó mediante coeficientes de correlación intraclase y de Pearson (CCP) entre los puntajes de los patrones dietarios identificados con CFA y 24DRs. **Resultados.** El patrón 1 fue alto en bocadillos, carnes procesadas, comida rápida, refrescos y granos refinados; el patrón 2 estuvo caracterizado por vegetales, frutas y lácteos; y el patrón 3 se caracterizó por leguminosas, huevo, azúcares, y alimentos dulces. Los CCP entre estos patrones fueron 0.66 ($P<0.001$), 0.41 ($P<0.001$) y 0.29 ($P=0.193$) respectivamente. **Conclusión.** Los datos sugieren razonable validez del CFA, utilizando análisis factorial, para derivar patrones dietarios en comparación con el 24DRs.

Palabras clave: cuestionario; encuesta; evaluación dietética; adultos; México

- (1) Unidad de Investigación en Epidemiología Clínica, Hospital Infantil de México. Ciudad de México, México.
- (2) Department of Clinical Laboratory and Nutritional Sciences, University of Massachusetts Lowell. Lowell, MA, USA.
- (3) Centro de Investigación en Salud Poblacional, Instituto Nacional de Salud Pública. Cuernavaca, Morelos, México.
- (4) Unidad de Investigación Epidemiológica y en Servicios de Salud, Instituto Mexicano del Seguro Social. Cuernavaca, México.
- (5) Centro de Investigación en Nutrición y Salud, Instituto Nacional de Salud Pública. Cuernavaca, Morelos, México.

Received on: March 10, 2016 • Accepted on: September 12, 2016

Corresponding author: Edgar Denova-Gutiérrez. Subdirección de Investigación en Epidemiología Geriátrica, Instituto Nacional de Geriátrica. Bvd. Adolfo Ruíz Cortines 2767, San Jerónimo Lídice. 10200 Ciudad de México, México.
E-mail: edenovag@gmail.com

Nutritional epidemiology research has traditionally adopted a reductionist approach^{1,2} focusing on relationships between individual nutrients or foods and disease.³ However, that approach has some important limitations. It fails to account for interactions among nutrients^{3,4} and cannot explain the synergistic effects of recognized or unrecognized constituents of daily food consumption.^{3,5,6} Further, intercorrelation between some nutrients makes it complicated to evaluate their effects independently.⁴ Finally, single nutrient assessment may be confounded by the effect of dietary patterns, with which consumption of specific nutrients is normally associated.^{7,8} Thus, dietary pattern analysis has been proposed to account for the cumulative and complex effects of simultaneous consumption of multiple nutrients or foods on a daily basis,^{4,9,10} and to understand the role whole diets play in disease occurrence.^{11,12}

Dietary patterns can be assessed with different statistical techniques: a priori techniques using score-based approaches, like the Mediterranean diet score or healthy eating index,^{13,14} or a posteriori methods; implemented with data-driven techniques such as factor analysis, cluster analysis, or, more recently, reduced rank regression.^{3,9,15,16} Factor analysis, a data reduction method which creates linear combinations of foods or food groups to identify the principal factors behind the largest variation in food consumption, is one of the most commonly used of these a posteriori techniques. Factors can be rotated (usually orthogonally) to enhance interpretability, and factor scores are computed for each individual.^{10,17}

Assessment of dietary patterns by factor analysis involves subjective judgment in classification of food items, determination of the number of factors to retain, method of rotation, and labeling of dietary patterns.¹⁸ Furthermore, dietary patterns may vary across populations with respect to food availability, socioeconomic status, resident area, ethnic group, and culture. For those reasons, it is useful to evaluate the validity of the food frequency questionnaire (FFQ) for assessing identification of dietary patterns in a particular study population.⁷

While studies have evaluated the reproducibility and validity of dietary patterns in various adult populations,^{6,7,19-21} the validity of a semi-quantitative food frequency questionnaire (SFFQ) to derive dietary patterns in the Mexican population has not been evaluated. Therefore, we assessed the relative validity of a semi-quantitative food frequency questionnaire to identify dietary patterns in an adult Mexican population.

Materials and methods

Design and study population

The present analysis was done with data from the Mexican National Health and Nutrition Survey 2012 (Ensanut 2012, for its acronym in Spanish). The Ensanut 2012 is a probabilistic population-based survey with multi-stage stratified sampling, designed to be representative of the nation, its three main regions (North, Center, and South), and rural and urban areas. The survey design and sampling procedures have been described in detail previously.²² Briefly, the Ensanut 2012 aims to monitor health and nutrition conditions, health program coverage, and access to health services. The survey obtained information from 50 528 households, with a response rate of 87%; 46 303 interviews with adults (≥ 20 y) were conducted. We include information from a subsample of adults aged ≥ 20 y, randomly selected for a validity study of the semi-quantitative food frequency questionnaire (SFFQ) used in the Ensanut 2012. The present analysis used data from 264 adults who completed one SFFQ and two 24-hour dietary recalls (24DRs). Participants with outlier energy intake values ($n=21$) were eliminated using the standard deviation method suggested by Rosner.²³ Consequently, our final analysis included 243 participants (148 women and 95 men). This study was managed according to Declaration of Helsinki guidelines, and written informed consent was obtained from all participants. Research Ethics Committee at the National Institute of Public Health (INSP, for its acronym in Spanish) approved the study protocol (Number 13CEI1700736).

Dietary assessment

The semi-quantitative food frequency questionnaire

The SFFQ used in the Ensanut 2012 is an adapted version of the questionnaire employed in the Ensanut 2006.²⁴ This questionnaire includes consumption of 140 foods during seven days prior to the date of the interview. Frequency of food items could be characterized by set categories ranging from never to six times a day. Participants also designated the portion size of the food items ingested, using predefined categories. These data were converted to portions per day. To calculate the consumption of energy (kcal/day), the daily frequency of consumption (portions/day) of each food was multiplied by the food's energy content (using the food composition tables compiled by the INSP)²⁵

and the contributions of all foods were totaled using Microsoft Visual FoxPro 7.0. The SFFQ was managed by personnel trained in standardized data collection and entry procedures.

24-hour dietary recall

Participants in the validity study completed two 24DRs 2 days apart, distributed over all days of the week, with approximately 50% of them obtained in weekend days. Personnel trained in standardized methods collected all the required information through face-to-face interviews, using 24DR automated multiple pass method (24DR-AMPM) software originally developed by the US Department of Agriculture and adapted (24DR-AMPM software version 1.0) by a group of INSP researchers to the Mexican population.²⁶ For each 24DR, participants were asked about their food consumption during the previous day in detail. In brief, at the beginning of the interview, participants listed the food items they had consumed during the previous day, with prompts from the interviewer about different possible eating occasions. A list of foods that are often forgotten was also used to elicit recall. Subsequently, detailed information on each food item was collected (including brand name, recipe, preparation method, time, occasion and amount eaten). Following data collection, we matched the 552 unique food reported in the 24DRs to food items in the SFFQ to ensure that the food intakes quantified by each were comparable.

Food grouping

The energy intake from each food was converted to percentage of total energy intake per day and standardized by Z-score.²⁷ Foods and beverages from the FFQ and the 24DRs were categorized into 29 food groups (table I) used to derive dietary patterns via factor analysis of principal components. Details of the food groupings used to derive the dietary patterns are described elsewhere.^{28,29} Briefly, as explained in Denova and colleagues,²⁸ the basis for placing a food item in a certain food group was the similarity of nutrients. Some groups were defined according to the amount of sugar added (e.g. sweetened beverages). Other groups were defined according to their lipid profile (e.g. seeds). Finally, some food items were considered individually as a food group, because their nutrient profiles were unique, they were consumed especially frequently, or they had unique culinary use (e.g. tortillas, eggs, and orange juice).

To derive dietary patterns and to determine factor loadings for each of the 29 groups, a factor analysis of the main components was performed.³⁰ The factors were

orthogonally rotated (varimax rotation) to keep them uncorrelated and to improve their interpretation. Factors above with Eigenvalues >1.5 were retained after graphic analysis, including scree plots³¹ and consideration of interpretability. Each factor was defined by a subset of at least five food groups with absolute loadings ≥ 0.2 .²⁷⁻²⁹ Factor scores for each dietary pattern were estimated by adding the consumption of the food groups weighted by their loading, and each participant received a factor score for each identified dietary pattern.

Other participant characteristics

Sociodemographic characteristics (age, sex and socioeconomic status) were obtained with predefined questionnaires. Localities with less than 2 500 residents were considered rural, and areas with 2 500 or more residents were considered urban. A household wealth index (HWI) was created using principal components analysis with household characteristics and family assets. This index was divided into tertiles, with 1 being the lowest category.

Anthropometric measures (weight and height) were collected with validated and standardized methods.³² Body weight was measured with a previously calibrated electronic scale, with participants wearing minimal clothing and no shoes. Height was measured with a stadiometer with barefoot participants standing with their shoulders in a normal position; measurements were taken with the tape in a horizontal plane perpendicular to the vertical scale, touching the top of the head at the moment of inspiration. Body mass index (BMI) was computed as a ratio of weight in kilograms divided by the height in meters squared. The definition for normal weight was BMI ≥ 18.5 and < 25.0 , participants with BMI ≥ 25.0 - < 30.0 were classified as overweight, and those with a BMI ≥ 30.0 were categorized as obese.

Statistical analysis

We performed descriptive analysis of the main characteristics of interest by sex. Mean daily intakes, in g/day, of the 29 food groups determined from the SFFQ and from the average of the two 24DRs, were calculated. We assessed intraclass correlation coefficients (ICC) comparing daily intakes between SFFQ and an average of two 24DRs of the food groups. Pearson correlation coefficients were also used to evaluate the validity of dietary patterns derived from dietary data collected with the SFFQ and the two 24DRs. To reduce within-person variation in food intake obtained from the 24DRs, we conducted factor analysis using the average consumption for each food group across the two days.

Table I
FOOD GROUPING USED IN THE DIETARY PATTERN ANALYSIS

Main group	Basis for placing a food item	Food groups	Food items
Grains	Culinary use	Corn tortilla	Corn tortilla
		Mexican food	Pozole, memela, quesadilla, sope, taco, tamal
	Proportion of fiber	Whole grains	Whole bread, oatmeal, linseed, all bran, multi bran, multigrain
		Refined grains	White bread, wheat tortilla, rice, corn flakes, honey crunch, other cereals
	Specific nutrient profile	Pastries	Pastries
		Desserts	Cookies, cakes, doughnuts
		Snacks	Potato chips, corn chips, popcorn, crackers
Fast food		Pizza, hot dogs, hamburgers	
Vegetables	Proportion of fiber	Fresh vegetables	Cauliflower, spinach, lettuce, carrots, tomato, nopal, onion, corn, cabbage, pea, green bean, chili, hot pepper, beet, mixed vegetables
	Proportion of starch	Potatoes	Potatoes
Fruits	Proportion of fiber	Fresh fruits	Banana, prune, peach, apple, orange, grapes, strawberry, melon, watermelon, mango, tangerine, pear, papaya, pineapple, guava, prickly pear
	Frequency of consumption	Fruit juice	Fruit juice
Meats	Frequency of consumption	Eggs	Eggs
		Poultry	Chicken with or without skin
	Specific nutrient profile	Red meat	Pork, beef or lamb
		Processed meat	Sausage, bacon, ham
Dairy	Specific nutrient profile “proportion of fat”	Fish and other sea food	Canned tuna fish, sardines, fresh fish, octopus, and squid
		Low-fat dairy products	Skim or low-fat milk, low-fat yogurt
		High fat dairy products	Whole milk, chocolate milk, cream, high fat yogurt, cream cheese, other cheese, ice cream
Legumes	Frequency of consumption	Sugary dairy products	Chocolate milk, other flavored milk, danonino and yakult brand yogurt, other yogurt with added sugar
Fat	Specific nutrient profile “proportion of fat and type of fat”	Legumes	Lentils, dry beans
		Oils and nuts	Peanuts, walnuts, almonds, pistachios, vegetable oils, avocados
Sugar	Frequency of consumption “proportion of sugar”	Butter	Margarine, butter, mayonnaise, animal fats
		Sweetened food and sugars	Sugar, chocolate, candies, jam, honey, jelly
		Soft drinks	Soft drinks
		Other sweetened beverages	Other sweetened beverages
Alcohol	Relative frequency of consumption	Low calorie drinks	Low calorie carbonated beverages, other low calorie beverages
Tea and coffee	Relative frequency of consumption	Alcohol	Wine, beer, brandy, whisky, tequila, rum, pulque, other hard liquor
		Tea and coffee	Tea and coffee

All statistical analyses were performed with STATA statistical software version 13.0. A value of $p < 0.05$ was considered to be statistically significant.

Results

This analysis included data from a sample of 243 adults older than 20 years. Of these participants, 61% were women and 39% men. The majority of women (71.4%) were between 20 and 49 years old, 39.8% lived in central Mexico, and approximately 80% were overweight/obese. Of the men, 54.2% lived in urban areas, 61.7% were between 20 and 49 years of age, and 70.6% were overweight or obese (table II).

Factor analysis identified three major dietary patterns that we termed "pattern 1", "pattern 2" and "pattern 3." These dietary patterns accounted for approximately 20% of the total variance; 20.4% in the SFFQ and 19.5% in the 24DRs. The greater the loading of a given food or food group to the factor, the higher the contribution of that food or food group to that specific factor.⁷ Pattern 1, which reflected correlated intakes of foods commonly considered to be unhealthy, was loaded heavily with snacks, fast food, soft drinks, processed meat and refined grains. Pattern 2 emphasized consumption of fresh vegetables, fresh fruit, and high fat dairy products. Finally, legumes, eggs and sweetened foods and sugars contributed heavily to pattern 3.

Table II
DISTRIBUTION OF CHARACTERISTICS OF INTEREST
IN THE VALIDATION SAMPLE DATA FROM THE
MEXICAN NATIONAL HEALTH AND NUTRITION
SURVEY OF 2012. MEXICO, ENSANUT 2012

Variables	Women (n = 160)		Men (n = 104)	
	n	%	n	%
Age groups				
20 – 49 years	100	71.4	53	61.7
≥ 50 years	60	28.6	51	38.3
Region				
North	32	15.2	29	25.2
Central	61	57.5	34	40.5
South	67	27.3	41	34.3
Area				
Urban	92	54.9	69	54.2
Rural	68	45.1	35	45.8
Household wealth index				
Tertile 1	60	28.3	40	36.4
Tertile 2	62	38.0	33	36.3
Tertile 3	38	33.7	31	27.4
Body mass index*				
Normal	43	18.2	33	29.4
Overweight	67	44.6	41	55.7
Obese	50	37.2	30	14.9

* Body mass index: Normal (< 25.0 kg/m²), overweight (≥ 25 kg/m² - < 30.0 kg/m²), obese (≥ 30.0 kg/m²)

In general, the first two dietary patterns derived from the SFFQ and the 24DRs were similar. However, the third pattern was less consistent across the two sources of data. Of the 29 food groups examined with the SFFQ and the two 24DRs, eight were identified as significant contributors for the dietary pattern 1, with six of them positively associated and two inversely associated. Of these, only five food groups (corn tortilla, snacks, fast foods, soft drinks, and sweetened foods and sugars) were similar when comparing pattern 1 derived with both instruments. The dietary pattern 2 originated from the SFFQ had nine food groups, while the dietary pattern 2 derived with the two 24DRs has ten. Of them, food groups that positively matched were whole grains, fresh vegetables, fresh fruits, red meat, and high fat dairy products; whereas, soft drinks matched but negatively. Lastly, for the third dietary pattern, four out of the seven and eight food groups derived with

the SFFQ and the two 24DRs respectively, were similar (eggs, legumes, sweetened foods and sugars, and tea and coffee) (table III).

Mean daily consumption of foods or food groups (g/day) was overestimated by the SFFQ compared with the 24DRs (considered the gold standard) for corn tortilla, fast food, fresh vegetables, fresh fruit, fruit juice, fish and other sea food, soft drinks and low calorie drinks. In contrast, Mexican foods, refined grains, potatoes, red meat, processed meat, legumes and sugary dairy products appeared to be underestimated by the SFFQ. The energy-adjusted ICCs between SFFQ and the average of the 24DRs were calculated to assess the relative validity of the SFFQ (table IV). Most foods or food groups were moderately correlated. The energy adjusted ICCs ranged between 0.08 (low energy drinks) to 0.64 (soft drinks). In particular, intakes of corn tortilla, pastries, fruit juices, and soft drinks showed high (>0.50) intra-class correlation coefficients. Whereas, whole grains, fresh vegetables, poultry, low-energy drinks, oils and nuts presented low (<0.20) ICCs.

The correlation between the SFFQ and the 24DRs was 0.66 ($p < 0.001$) for dietary pattern 1 and 0.41 ($p < 0.001$) for pattern 2. However, the correlation for dietary pattern 3 was less consistent and not statistically significant ($r = 0.29$; $p = 0.19$) (data not shown).

Discussion

There has recently been increased interest in dietary pattern analysis as a method to examine diet-disease relationships, as this approach offers several advantages over single nutrient or food methodologies.¹⁷ Dietary patterns represent a combination of nutrients or foods and other dietary components that can reflect the eating habits of the population. Few epidemiological studies, however, have evaluated the validity of dietary patterns.^{6,7,19-21} The aim of the present work was to evaluate relative validity of a semi-quantitative food frequency questionnaire compared to 24DRs to identify dietary patterns in an adult Mexican population. We derived three major dietary patterns in this study of the adult Mexican population, which were qualitatively similar across the two sources of dietary data. However, the patterns derived from the SFFQ and 24DRs did differ in some aspects, possibly because of methodological dissimilarities between the two dietary assessment methods³³ and random statistical variation. However, the correlation coefficients for patterns between the SFFQ and the 24DRs ranged from 0.29 to 0.66, suggesting reasonable comparability across the two methods in typifying dietary patterns, and supporting the utility of the SFFQ in assessing dietary patterns.

Table III
FACTOR LOADING MATRIX FOR THE THREE MAJOR DIETARY PATTERNS IDENTIFIED FROM THE SFFQ
AND THE AVERAGE OF TWO 24DRs. MEXICO, ENSANUT 2012

Foods or food groups	SFFQ			24DRs		
	Dietary pattern 1	Dietary pattern 2	Dietary pattern 3	Dietary pattern 1	Dietary pattern 2	Dietary pattern 3
Corn tortilla	-0.79 ^a	–	–	-0.67	–	–
Mexican food	–	–	–	0.52	–	–
Whole grains	–	0.21	–	–	0.28	–
Refined grains	–	–	–	0.31	–	–
Pastries	–	–	–	–	–	0.36
Snacks	0.58	–	–	0.43	–	–
Fast food	0.47	-0.40	–	0.39	–	–
Fresh vegetables	–	0.42	–	–	0.31	–
Potatoes	–	–	–	–	–	–
Fresh fruits	–	0.59	–	–	0.53	–
Fruit juices	–	0.36	–	–	–	–
Eggs	–	–	0.51	–	-0.55	0.31
Poultry	0.38	–	–	–	–	0.59
Red meat	-0.23	-0.32	0.24	–	-0.35	–
Processed meats	0.34	–	0.39	–	-0.32	–
Fish and other sea food	–	–	–	–	–	–
Low fat dairy products	–	0.28	–	–	–	–
High fat dairy products	–	0.31	–	–	0.42	–
Sugary dairy products	–	–	–	–	0.44	–
Legumes	–	–	0.33	-0.56	–	0.39
Oils and nuts	–	–	–	–	-0.40	–
Butter	–	–	–	–	–	-0.32
Sweetened food and sugars	0.26	–	0.55	0.29	–	0.39
Soft drinks	0.31	-0.56	–	0.52	-0.23	–
Other sweetened beverages	–	–	–	–	–	0.54
Alcohol	–	–	-0.35	–	–	–
Tea and coffee	–	–	0.48	–	–	0.21

^aAbsolute values <0.20 were excluded for simplicity

The dietary patterns derived in the present analysis are similar to patterns identified in other studies using factor analysis methodology to dietary data from different populations. Such studies have found vegetable rich patterns, similar to our pattern 2, generally labeled as “healthy” or “prudent”,^{7,19,21,34} and “western” patterns resembling those in our dietary patterns 1 and 3.^{7,19} For example, in the Health Professionals Follow-up Study, Hu and colleagues⁷ evaluated the reproducibility and validity of dietary patterns using dietary data collected by SFFQ and dietary records. In that study, “prudent” and “western” dietary patterns were identified. The correlation coefficients in the validity analysis between each of the patterns based on SFFQ and dietary records

were 0.45–0.74, comparable with the correlation coefficients that we observed (0.41 – 0.66). A similar analysis of data from the Swedish Cohort Study¹⁹ identified three dietary patterns, including “healthy” and “western” patterns similar to our patterns 2 and 1. The correlation coefficients between the patterns derived from SFFQ and dietary records in that study were 0.50–0.59.

We observed that factor loadings for patterns from the 24DRs were more weakly correlated with factor loadings from the SFFQ than factor scores overall. This most likely reveals methodological differences between the FFQ and 24DRs. The SFFQ asks for information on usual dietary intake, in our case during the past seven days, whereas the 24DRs measures food consumed the

Table IV
DAILY MEAN INTAKES AND ENERGY-ADJUSTED INTRACLASS CORRELATION COEFFICIENTS
FOR FOOD GROUP INTAKES, ESTIMATED BY THE SFFQ AND THE AVERAGE OF TWO 24DRs
IN THE NATIONAL HEALTH AND NUTRITION SURVEY 2012. MEXICO, ENSANUT 2012

Foods or food groups	Grams/day		ICC (95%CI)	
	FFQ Mean (SE)	24DRs Mean (SE)	FFQ vs 24DRs	P-value
Corn tortilla	237 (42)	173 (23)	0.57 (0.45, 0.66)	<0.001
Mexican food	89.7 (14.9)	93.9 (13.9)	0.31 (0.16, 0.44)	<0.001
Whole grains	9.1 (3.8)	8.3 (3.6)	0.08 (-0.08, 0.21)	0.201
Refined grains	92.8 (11.6)	99.5 (14.1)	0.46 (0.32, 0.60)	<0.001
Pastries	33.5 (9.5)	32.4 (5.0)	0.50 (0.38, 0.61)	<0.001
Desserts	29.8 (2.6)	33.3 (2.0)	0.31 (0.16, 0.44)	<0.001
Snacks	4.3 (0.9)	5.7 (2.1)	0.40 (0.26, 0.52)	<0.001
Fast food	37.6 (8.4)	9.5 (3.4)	0.25 (-0.02, 0.45)	<0.001
Fresh vegetables	112. (14)	55.9 (34.0)	0.16 (-0.14, 0.38)	0.134
Potatoes	10.8 (1.6)	31.5 (38.6)	0.27 (0.12, 0.41)	<0.001
Fresh fruits	232 (24)	73.3 (10.4)	0.29 (0.15, 0.42)	<0.001
Fruit juices*	27.6 (8.6)	6.1 (4.1)	0.57 (0.46, 0.67)	<0.001
Eggs	40.9 (3.7)	35.5 (5.1)	0.36 (0.13, 0.53)	0.002
Poultry	25.5 (2.8)	39.6 (8.6)	0.10 (-0.06, 0.25)	0.107
Red meat	33.7 (4.6)	46.4 (7.9)	0.39 (0.26, 0.51)	<0.001
Processed meats	12.1 (2.8)	13.8 (3.1)	0.22 (0.07, 0.36)	0.003
Fish and other sea food	10.9 (2.7)	4.6 (1.4)	0.30 (0.05, 0.48)	0.010
Low fat dairy products*	14.9 (3.0)	23.6 (9.4)	0.40 (0.26, 0.51)	<0.001
High fat dairy products*	145 (25)	85.1 (15.3)	0.33 (0.14, 0.46)	<0.001
Sugary dairy products*	28.9 (10.1)	45.5 (12.6)	0.41 (0.27, 0.53)	<0.001
Legumes	91.3 (11.0)	103 (18)	0.23 (-0.08, 0.45)	0.196
Oils and nuts	15.2 (4.4)	10.9 (1.5)	0.15 (0.01, 0.29)	0.025
Butter	5.6 (1.0)	11.2 (1.8)	0.19 (0.03, 0.32)	0.009
Sweetened food and sugars	40.8 (8.0)	20.9 (4.7)	0.43 (0.25, 0.51)	<0.001
Soft drinks*	210 (29)	190 (29)	0.64 (0.54, 0.72)	<0.001
Other sweetened beverages*	130 (28)	91.0 (20.3)	0.47 (0.22, 0.61)	<0.001
Low-energy drinks*	23.3 (9.7)	6.8 (3.5)	0.07 (-0.08, 0.23)	0.171
Alcohol*	47.2 (16.2)	76.2 (32.6)	0.29 (0.14, 0.42)	<0.001
Tea and coffee*	148 (28)	202 (31)	0.32 (0.17, 0.44)	<0.001

* Intakes are presented in mL/day

ICC: energy-adjusted Intraclass correlation coefficients

previous day, and therefore includes a smaller range of foods than are likely to be reported by SFFQ. Additionally, as factor solutions are influenced by the correlation matrix, some discrepancies may be expected between factor loading matrixes from different dietary assessment methodologies.

There are some limitations in our study. First, dietary pattern analysis should be interpreted with caution because, although at least two major patterns (“healthy”

or “prudent” and “unhealthy or “western”) have commonly emerged in different populations,^{7,19,28,29,34} their specific composition depends on geographical, cultural, socioeconomic and ethnic status, and is influenced by methodological variation (including sampling, food grouping, number of variables used in factor analysis, number of factors and the rotation employed). Further, the three major dietary patterns identified in our study explained only 20.4% of the total variance in the SFFQ

and 19.5% in the 24DRs, suggesting the existence of other eating pattern dimensions, although remaining patterns were less interpretable in our analysis. Dietary patterns aside from the “western” and “prudent”, have been shown to be highly variable across various dietary assessment methods, and may not be reproducible across populations.⁷ In the present study, two 24DRs were used as comparison method; therefore, the estimate of intake may have been closer to true intake if we had collected more days of food consumption. However, other findings suggested the possibility of using only a small number of replicate measures for the reference method combined with a statistical adjustment to remove the effects of within-person variation.³⁵ At this respect, some authors have proposed that the greatest statistical efficiency is obtained with only two, and at the most five, duplicates per subject.^{36,37} In this context, to validate our SFFQ against two 24DRs seems to be adequate. Furthermore, in order to account for the week and weekend day variation, approximately 50% of the 24DRs were obtained in weekend days. Finally, measurement errors may have occurred. Although the 24DR is the current gold standard for evaluating food consumption, it is nevertheless susceptible to measurement error due to inaccurate recording.³³ In addition, as we did not administer a second SFFQ in the validity study, we cannot assess the reproducibility of the instrument. Future work needs to be done to evaluate the reproducibility of this SFFQ.

In conclusion, our data indicate reasonable validity of the SFFQ, using factor analysis, to derive major dietary patterns in comparison with two 24DR. These results suggest the potential use of SFFQ using factor analysis-based dietary pattern identification in epidemiological studies as an alternative dietary assessment method suitable for studying diet-disease relationships.

Declaration of conflict of interests. The authors declare that they have no conflict of interests.

References

- Messina M, Lampe JW, Birt DF, Appel LJ, Pivonka E, Berry B, et al. Reductionism and the narrowing nutrition perspective: Time for reevaluation and emphasis on food synergy. *J Am Diet Assoc* 2001;101(12):1416-1419. <http://doi.org/bgdstp>
- Hoffmann I. Transcending reductionism in nutrition research. *Am J Clin Nutr* 2003;78 (suppl. 3):S145-S165.
- Jacobs DR, Steffen LM. Nutrients, foods, and dietary patterns as exposure in research: a frame work for food synergy. *Am J Clin Nutr* 2003;78 (suppl. 3):S08S-S13S.
- Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* 2002;13(1):3-9. <http://doi.org/dzfrnv>
- Jacobson HN, Stanton JL. Pattern analysis in nutrition. *Clin Nutr* 1986;5:249-253.
- Quatromoni PA, Copenhafer DL, Demissie S, D'Agostino RB, O'Horo CE, Nam BH, et al. The internal validity of a dietary pattern analysis. The Framingham Nutrition Studies. *J Epidemiol Community Health* 2002;56(5):381-388. <http://doi.org/d3rdn>
- Hu FB, Rimm E, Smith-Warner SA, Feskanich D, Stampfer MJ, Ascherio A, et al. Reproducibility and validity of dietary patterns assessed with a food frequency questionnaire. *Am J Clin Nutr* 1999;69(2):243-249.
- Kant AK, Schatzkin A, Blook G, Ziegler RG, Nestle M. Food group intake patterns and associated nutrient profiles of the US population. *J Am Diet Assoc* 1991;91(12):1532-1537.
- Newby PK, Tucker KL. Empirically derived eating patterns using factor or cluster analysis: A review. *Nutrition Reviews* 2004;62(5):177-203. <http://doi.org/bxcscck>
- Moeller SM, Reedy J, Millen AE, Dixon LB, Newby PK, Tucker KL, et al. Dietary patterns: challenges and opportunities in dietary patterns research. *J Am Diet Assoc* 2007;107(7):1233-1239. <http://doi.org/d5prtf>
- Kant AS. Dietary patterns: biomarkers and chronic disease risk. *Appl Physiol Nutr Metab* 2010;35(2):199-206. <http://doi.org/bm3m9x>
- Slattery ML. Analysis of dietary patterns in epidemiological research. *Appl Physiol Nutr Metab* 2010;35(2):207-210. <http://doi.org/drgxb4>
- Newby PK, Hu FB, Rimm EB, Smith-Warner SA, Feskanich D, Sampson L, et al. Reproducibility and validity of the diet quality index revised as assessed by use of a food-frequency questionnaire. *Am J Clin Nutr* 2003;78(5):941-949.
- Panagiotakos DB. “A priori” versus “a posteriori” methods in dietary patterns analysis: A review in nutritional epidemiology. *Nutr Bull* 2008;33:311-315. <http://doi.org/bs5htk>
- Kant AK. Dietary patterns and health outcomes. *J Am Diet Assoc* 2004;104(4):615-635. <http://doi.org/czbdxc>
- Michels KB, Schulze MB. Can dietary patterns help us detect diet-diseases association? *Nutr Res Rev* 2005;18:241-248. <http://doi.org/fn2p3s>
- Tucker KL. Dietary patterns, approaches, and multicultural perspective. *Appl Physiol Nutr Metab* 2010;35(2):211-218. <http://doi.org/b5nx3t>
- Martínez ME, Marshall JR, Sechrest L. Invited commentary: Factor analysis and the search for objectivity. *Am J Epidemiol* 1998;148(1):17-19. <http://doi.org/brbs>
- Khani BR, Ye W, Terry P, Wolk A. Reproducibility and validity of major dietary patterns among Swedish women assessed with a food frequency questionnaire. *J Nutr* 2004;134(6):1541-1545.
- Bountziouka V, Tzavelas G, Polychronopolis E, Constantinidis TC, Panagiotakos DB. Validity of dietary patterns derived in nutrition surveys using a priori and a posteriori multivariate statistical methods. *Int J Food Sci Nutr* 2011;62(6):617-627. <http://doi.org/fd4kvx>
- Nanri A, Shimazu T, Ishihara J, Takachi R, Mizoue T, Inoue M, et al. Reproducibility and validity of dietary patterns assessed by a food frequency questionnaire used in the 5-year follow-up survey of the Japan Public Health Center-Based prospective study. *J Epidemiol* 2012;22(3):205-215. <http://doi.org/brbt>
- Romero-Martínez M, Shamah-Levy T, Franco-Núñez A, Villalpando S, Cuevas-Nasu L, Gutiérrez JR, et al. National Health and Nutrition Survey 2012: design and coverage. *Salud Publica Mex* 2013;55 (Suppl. 2):S332-S340. <http://dx.doi.org/10.21149/spm.v55s2.5132>
- Rosner B. Percentage points for a generalized ESD many-outlier procedure. *Technometrics* 1983;25(2):165-172. <http://doi.org/brbv>
- Rodríguez-Ramírez S, Mundo-Rosas V, Jiménez-Aguilar A, Shamah-Levy T. Methodology for the analysis of dietary data from the Mexican National Health and Nutrition Survey 2006. *Salud Publica Mex* 2009;51 (Suppl. 4):S523-S529. <http://doi.org/fwfwpw>
- Hernández-Ávila JE, González-Avilés L, Rosales-Mendoza E. Manual de usuario. SNUT Sistema de Evaluación de Hábitos Nutricionales y Consumo de Nutrientes. México: Instituto Nacional de Salud Pública, 2003.

26. Moshfegh AJ, Rhodes DG, Murayi T, Clemens JC, Rumpel WV, Paul DR, et al. The US Department of Agriculture Automated Multiple-Pass Method reduces bias in the collection of energy intakes. *Am J Clin Nutr* 2008;88(2):324-332.
27. Newby PK, Weismayer C, Akesson A, Tucker KL, Wolk A. Long-term stability of food patterns identified by use of factor analysis among Swedish women. *J Nutr* 2006;136(3):26-33.
28. Denova-Gutiérrez E, Castañón S, Talavera JO, Flores M, Macías N, Rodríguez-Ramírez S, et al. Dietary patterns are associated with different indexes of adiposity and obesity in an urban Mexican population. *J Nutr* 2011;141(5):921-927. <http://doi.org/cnzzxx>
29. Denova-Gutierrez E, Castañón S, Talavera JO, Gallegos-Carrillo K, Flores M, Dosamantes-Carrasco D, et al. Dietary patterns are associated with metabolic syndrome in an urban Mexican population. *J Nutr* 2010;140(10):1855-1863. <http://doi.org/bbcmv>
30. Kim JO, Muller CW. Factor analysis. Newbury Park, CA: Sage Publications, Inc. 1984.
31. Kim JO, Muller CW. Factor analysis: statistical methods and practical issues. Thousand Oaks, CA: Sage Publications Inc, 1987.
32. Lohman T, Roche A, Martorell R. Anthropometric standardization reference manual. Champaign, IL: Human Kinetics, 1988.
33. Willett WC. Nutritional Epidemiology. 2nd ed. New York: Oxford University Press, 1998. <http://doi.org/dwvp9n>
34. Ambrosini GL, O'Sullivan TA, Klerk NH, Mori TA, Beilin LJ, Oddy WH. Relative validity of adolescent dietary patterns: a comparison of a food frequency questionnaire and 3-day food record. *Br J Nutr* 2011;105(4):625-633. <http://doi.org/dwx3dr>
35. Deschamps V, de Lauzon-Guillain B, Lafay L, Borys JM, Charles MA, Romon M. Reproducibility and relative validity of a food-frequency questionnaire among French adults and adolescents. *Eur J Clin Nutr* 2009;63(2):282-391. <http://doi.org/d2m7dq>
36. Carroll RJ, Pee D, Freedman LS, Brown CC. Statistical design of calibration studies. *Am J Clin Nutr* 1997;65(4 Suppl):1187S-1189S.
37. Rosner B, Willett WC. Interval estimates for correlation coefficients corrected for within-person variation: implications for study design and hypothesis testing. *Am J Epidemiol* 1988;127(2):377-386.