



Escola Anna Nery Revista de
Enfermagem

ISSN: 1414-8145

annaneryvista@gmail.com

Universidade Federal do Rio de Janeiro
Brasil

Barbosa, Luma Natalia; Cunha dos Santos, Neuci; Maciel de Moraes, Maria Auxiliadora;
Daiana Rizzardi, Scheila; da Costa Corrêa, Eloah

Prevalence of educational practices about exclusive breastfeeding (EBF) in Cuiabá - MT
Escola Anna Nery Revista de Enfermagem, vol. 19, núm. 1, enero-marzo, 2015, pp. 147-
153

Universidade Federal do Rio de Janeiro
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=127737750020>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Prevalence of educational practices about exclusive breastfeeding (EBF) in Cuiabá - MT

Prevalência de práticas educativas acerca do aleitamento materno exclusivo (AME) em Cuiabá - MT

Prevalencia de prácticas educativas sobre amamantamiento materno exclusiva (AME) en Cuiabá - MT

Luma Natalia Barbosa¹

Neuci Cunha dos Santos²

Maria Auxiliadora Maciel de Moraes¹

Scheila Daiana Rizzardi¹

Eloah da Costa Corrêa¹

1. Federal University of Mato Grosso.
Cuiabá - MT, Brazil.

2. Oswaldo Cruz Foundation.
Cuiabá - MT, Brazil.

ABSTRACT

Objective: To describe the prevalence of educational practices in health on exclusive breastfeeding (EBF) in the city of Cuiabá between July and December 2012. **Methods:** This was a descriptive, cross-sectional study, carried out among hospitalized women. The sample consisted of 306 women in the immediate postpartum period. Data were collected through a semi-structured questionnaire. **Results:** Eight years of schooling or more (OR = 1.77); income of up to two minimum wages (OR = 1.22); planned pregnancy (OR = 1.31); start of antenatal care in the first term (OR = 1.65) and primiparous (OR = 1.21) were features that appear associated with a greater chance of receiving breastfeeding training than other women. **Conclusion:** It is necessary to rethink these practices, in order to carry out educational activities that aim for changes in the professional/user relationship, with a view to the welcoming of pregnant women.

Keywords: Breastfeeding; Health Education; Prenatal Care.

RESUMO

Objetivo: Descrever a prevalência das práticas educativas em saúde sobre o aleitamento materno exclusivo (AME) no Município de Cuiabá - MT nos meses de julho a dezembro de 2012. **Métodos:** Trata-se de um estudo descritivo transversal, realizado junto a mulheres internadas em um hospital. A amostra foi composta por 306 mulheres no período pós-parto imediato. Os dados foram coletados por meio de um questionário semiestruturado. **Resultados:** Ter oito anos ou mais de estudo (RP = 1,77); ter renda até dois salários mínimos (RP = 1,22); ter planejado a gravidez (RP = 1,31); ter iniciado o atendimento no serviço de pré-natal no primeiro trimestre (RP = 1,65) e serem primíparas (RP = 1,21) são características que aparecem associadas com uma maior chance de receber orientações sobre aleitamento materno do que as demais mulheres. **Conclusão:** Faz-se necessário repensar mudanças dessas práticas, a fim de se realizar ações educativas que busquem efetivar transformações na relação profissional/usuário, visando o acolhimento das gestantes.

Palavras-chave: Aleitamento Materno; Educação em Saúde; Cuidado Pré-Natal.

RESUMEN

Objetivo: Describir la prevalencia de prácticas educativas en salud sobre el Amamantamiento Materno Exclusivo (AME) en la ciudad de Cuiabá, entre julio y diciembre de 2012. **Métodos:** Se realizó un estudio transversal entre mujeres internadas en un hospital. La muestra fue compuesta por 306 mujeres en el posparto inmediato. Los datos fueron recolectados a través de un cuestionario semiestruturado. **Resultados:** Tener ocho o más años de escolaridad (RP = 1,77); renta de hasta dos salarios mínimos (PR = 1,22); haber planeado el embarazo (PR = 1,31); haber iniciado el atendimento de atención prenatal en el primer trimestre (PR = 1,65); y ser primípara (PR = 1,21), son características que aparecen asociadas a una mayor probabilidad de recibir orientación sobre la lactancia que otras mujeres. **Conclusión:** Es necesario repensar los cambios de estas prácticas con el fin de realizar acciones educativas que tratan de efectuar transformaciones en la relación profesional/usuario, objetivando la acogida de las embarazadas.

Palabras-clave: Lactancia Materna; Educación en Salud; Atención Prenatal.

Corresponding Author:

Luma Natalia Barbosa.
E-mail: lumanataliab@hotmail.com

Submitted on 10/25/2013.
Accepted on 09/15/2014.

DOI: 10.5935/1414-8145.20150020

INTRODUCTION

Breast milk is currently considered the ideal food for infants, especially in the first six months of life, with benefits superior to other milks. The justification is based on the fact that it is rich in fats, minerals, vitamins, enzymes and immunoglobulins. Moreover, it has nutritional advantages, which promote growth and development and influence the future academic performance of children¹.

Breastfeeding prevents gastrointestinal, respiratory and urinary tract infections; besides having a protective effect on allergies. Breast milk makes babies adapt better to other foods. In the long term, we can also refer to the importance of breastfeeding in the prevention of diabetes and lymphomas. With regard to the benefits for the mother, breastfeeding facilitates an early uterine involution due to the release of oxytocin, and is associated with a lower likelihood of her developing cancer of the breast and ovaries, as well as the recovery of her pre-gestational weight. It mainly allows the mother to feel the unique pleasure of breastfeeding. Apart from all these benefits, breast milk is the cheapest and safest method of feeding infants and, in most situations, protects mothers from a new pregnancy².

The World Health Organization and the Brazilian Ministry of Health recommend exclusive breastfeeding (EBF) until the sixth month of the child's life, defining it as follows: "only receive milk from their mother or human milk banks, and no other liquids or solids except for vitamins, minerals and medicines". After the age of six months, the inclusion of complementary foods is recommended up to two years of age, along with breastfeeding^{3,2,4}.

Despite an upward trend in breastfeeding practices in the last three decades, it is still not possible to see mothers breastfeed their children exclusively until the sixth month of the child's life⁵. In Brazil, in general, breastfeeding is growing every year, but still far from what is widely recommended⁶.

According to the 2nd Survey of the Prevalence of Breastfeeding in the state capitals and the Federal District, held in 2008, the prevalence of exclusive breastfeeding in the first six months of life is 41%. In the same research, the capital of Cuiabá deserves special mention, where only 27.1% of mothers maintained EBF for six months. When compared with other state capitals, Cuiabá ranks first in early weaning⁷.

These data point to the relevance and rationale of the present study, whose problem was implied in the understanding that a woman's decision to breastfeed may be the result of an educational activity, considering that education is a powerful instrument for the emancipation of the decision and subject.

Educational practices are health education activities, designed to develop both individual and collective capacity, in order to improve the quality of life and health, ensuring access to quality health services. Health education can be seen as a constant

process of intervention for the individual and the community to have the means to maintain or restore their state of health, which are related the organic, psychological, socioeconomic and spiritual factors⁸.

Health education should be present at all care levels, but it is believed that it is in primary care that it is more meaningful to the subjects, because it can be the base to promote preventive actions and train individuals to be aware of their citizenship, to develop decision-making power over their own health and responsibility for the health of the community they live in. But for that to happen effectively, the sharing of knowledge, wisdom and experiences is needed⁹.

It is considered to be critical to conduct educational activities during all stages of pregnancy and childbirth, but it is in the prenatal stage that women should build for themselves the reasons that help in decision making to experience childbirth in a positive way, have less risk of complications in the puerperium and succeed in breastfeeding¹⁰.

Thus, the present article aimed to describe the prevalence of educational practices in health on exclusive breastfeeding in the city of Cuiabá between July and December 2012. This study permits diagnosing the failures in the lack of educational practices in health in the town under study. This is an initial way to rethink strategies and methods to reach these women.

METHOD

This descriptive, cross-sectional, quantitative study was undertaken at the Charity Hospital St. Helena (CHSH) in the city of Cuiabá. This service was chosen because it is a Baby-Friendly Hospital (BFH) and receives about 70% of all births in the capital of Mato Grosso, accepting pregnant women from all regions of the city.

According to population estimates for 2011, Cuiabá had a female population of childbearing age (10-49 years) corresponding to 192,422 residents. In the same year, 9,956 births occurred, among them 6,955 at the hospital observed in this study. The study was conducted using a sample calculated from of population and reported birth data with a 95% confidence interval (= 2 standard deviations) and sampling error of 2%.

- Proportion to be estimated: $p = 0.5$ (no information about the proportion of the population).
- Population size: $N = 6955$.

Equation used for the sample size (with correction for finite population):

$$n = (\sigma^2 \cdot p \cdot q \cdot N) / (e^2 (N-1) + \sigma^2 \cdot p \cdot q)$$

N is the population size of the study and p is the prevalence of the trait in the population of interest.

This formula calculates the sample size for finite populations of size N , and uses the information about the prevalence rate, sampling error and desired confidence interval.

This results in an initial sample of 1839 women of childbearing age. Assuming that there is a relative homogeneity of births between the months of the year, the final sample size was obtained by dividing the total sample by 12 months. The projection is a sample of 153 women per month. We chose to perform the collection in two months (December 2012 and January 2013) and thus expand the database and the sample population.

Data were collected from women in the postpartum period, and the selection of women was defined by the following inclusion criteria: mothers who had given birth in the last 24 hours and who had the ability to breastfeed. Exclusion criteria: mothers who by known biological factors were prevented from experiencing breastfeeding (i.e. with a diagnosis of active tuberculosis, leprosy, people with HIV, herpes, simplex virus in the breast, severe debilitating diseases, maternal malnutrition and need to ingest medicines harmful to the child over a long period).

Data were collected through a semi-structured questionnaire consisting of three groups of variables that included related issues: the identification data of the woman; data on prenatal care and data on educational practices in health about EBF.

Data were organized and analyzed using measures of frequency and distribution and the prevalence ratio was calculated using Epi Info software version 6.0.

The study results are consistent with information about antenatal health care services provided to pregnant women in the second half of 2012. This projection is based on the fact that 70% of women sought antenatal care before the fourth month of pregnancy.

For this study, the recommended ethical and legal aspects were observed and regulated by of the National Health Council (NHC) Resolution 196/96, in force at the time of the study. Approval was obtained from the Research Ethics Committee of the University Hospital Júlio Muller under number 146 386. After being informed about the objectives, the method and the potential benefits of the study, each study participant signed the Instrument of Consent. The study also received approval from the director of the institution where the research was conducted.

RESULTS

The sample population was composed of 306 postpartum women who completed the questionnaire and were still admitted at the study hospital.

The study participants' average age was 27.1 years. This finding reveals an appropriate age for the conception of a baby,

since the Ministry of Health (MOH) recommends that the best age to conceive a child is more than 15 years and less than 35 years. However, we know that teen pregnancy has increased in recent years, revealing the troubling demands to attend to these clients. In our findings, the average age of the participants reveals a situation contrary to what the MOH has indicated as frequent among women who become pregnant¹¹.

The education level of the participants was mostly concentrated (55.3%) in the group that completed high school and completed or are taking higher education. However, the data correspond to a lower level than that found in the city's female population though. In 2010, 70.61% of the female population in Cuiabá aged 15 years or more held a secondary education degree or higher.

With regard to the participants' marital status, 78% lived with a partner, in a stable relationship (35%), married (26%) or had partners (17%).

With regard to skin color, approximately 65% of the women self-reported that they were mulatto; 16.4% considered themselves white and 16.4% black. The amount of natives and yellows amounted to less than 3% of the total.

Of the total participants, ($n = 100$) reported having a remunerated job, 72% of whom affirm a formal employment contract.

According to Table 1, approximately 75% of the women reported having unplanned pregnancies. However, this factor did not discourage them from seeking pregnancy monitoring, as 99.7% of the total received antenatal care.

Of all the women who answered the questionnaire, 48.9% reported having received some professional guidelines for monitoring during the antenatal period, whereas only 20.3% of women reported having participated in some of the educational practice conversation groups and lectures. It should be emphasized that providing orientations on exclusive breastfeeding corresponds to what the Ministry of Health recommends as the minimum a routine antenatal consultation should contain¹¹.

Another fact related to access to health services refers to the sector where the woman received the antenatal care. We identified a large concentration in mixed services, involving the search for care in public and private units. This information shows the deficiencies in the health service network, especially in the provision of specialized tests. Of the study participants, 54.1% received antenatal care in the public sector, but resorted to ultrasound tests in private health services.

Women are increasingly seeking antenatal care early. The search for the service occurred mostly (73.4%) in the first term of pregnancy. The average number of antenatal consultations was about nine throughout the gestational period, higher than the minimum number of visits recommended by the Ministry of Health.

Table 1. Characterization of educational practices in healthcare about breastfeeding in Cuiabá - MT in the year 2012

Did you take the antenatal test for this pregnancy?	N	%
Yes	305	99.7
No	1	0.3
Total	306	100
Month that antenatal care began?	N	%
1 st month	49	16.1
2 nd month	95	31.1
3 rd month	80	26.2
4 th month	40	13.1
5 th month	26	8.5
6 th month	10	3.3
7 th month	4	1.3
8 th month	1	0.3
Total	305	100
Attended conversation groups and/or lectures on breastfeeding during antenatal care?	N	%
Yes	62	20.3
No	243	79.7
Total	305	100
How often have you attended?	N	%
All or almost all consultations	32	51.6
Only once	16	25.8
Rarely	14	22.6
Total	62	100
Received breastfeeding training during the medical consultations, nursing and/or home visits?	N	%
Yes	149	48.9
No	151	49.5
Does not remember	5	1.6
Total	305	100
Who was the professional who addressed the topic?	N	%
Doctor	117	38.2
Nurse	92	30.1
Community health worker	10	3.3
Technical nurse	8	2.6
Other	1	0.3
Total	228*	74.5

* This alternative does not total 149 nursing mothers (those who received orientations) as more than one alternative could be marked on the questionnaire.

We emphasize the following data on the prevalence and prevalence ratio (PR) indicating an association between the practice of health education and some variables that were most significant in the study. Table 2 shows the variables with a prevalence ratio superior to one. The variables that were highlighted were: eight or more

years of study; income up to two minimum wages; planned pregnancies; start of antenatal care in the first term, and first pregnancy.

Women with eight or more years of study reported having participated in or having received guidance on breastfeeding, 1.77 times more than those with less than eight years of study.

Table 2. Prevalence and Prevalence Ratio of childrearing practices on breastfeeding in relation to some variables of the study in Cuiabá - MT in the year 2012

Variable		Educational practices in health						
		Yes		No		Total	Prevalence	RP
		N	%	n	%			
Years of study	Eight full years and/or more	57	91.94	207	85.19	264	0.22	1.77
	Less than eight years	5	8.06	36	14.81	41	0.12	
Average salary	Up to two minimum wages	39	62.90	136	56.43	175	0.22	1.22
	More than two minimum wages	23	37.10	105	43.57	128	0.18	
Planning for pregnancy	Yes	19	30.65	56	23.05	75	0.25	1.31
	No	43	69.35	187	76.95	230	0.19	
Start of prenatal care	First term	51	82.26	173	71.19	224	0.23	1.65
	Second and third term	11	17.74	70	28.81	81	0.14	
Previous pregnancies	Primipara	23	37.10	77	31.69	100	0.23	1.21
	Not primipara	39	62.90	166	68.31	205	0.19	

Have income up to two minimum wages was seemingly a feature that distinguished the pregnant women as targets of health education activities (PR = 1.22).

Women who planned their pregnancy had more guidance on breastfeeding (OR = 1.31) than those who did not.

The start of antenatal care in the first term also appears as a factor that distinguishes women who received more guidance on breastfeeding (PR = 1.65).

The primipara women were 1.21 times more likely to receive information than other women.

DISCUSSION

The average age of participants is relatively high compared with other studies conducted in Brazil. The younger maternal age may be related to the shorter duration of breastfeeding, perhaps motivated by some difficulties, such as: low educational level, lower purchasing power and, often, the fact that they are single¹².

We can consider that living with a companion can be seen as fundamental for greater adherence to breastfeeding because it promotes better understanding of the benefits, and the presence of a companion may be beneficial, especially if he encourages, supports and helps in general tasks, both house chores and in child care¹³.

The increase in female participation in the labor market in the country has been one of the social changes since the 70's. Statistics point to an increasingly high presence of women in the Brazilian labor market and shows no inverse trend. This inclusion implies a change in the behavior of women regarding breastfeeding. However, our findings did not suggest a negative influence of work on EBF, because the mothers reported working outside the home as uncommon problems regarding interference in the maintenance of breastfeeding¹².

Even before this "almost perfect" coverage related to antenatal care, the prevalence of exclusive breastfeeding in the first six months of life remains low in the capital of Mato Grosso⁷.

Health education activities, mainly conducted in groups, are resources that permit further approximation between health professionals and the population, contributing to the supply of humanized and skilled care, as they are based on dynamic and reflexive human interactions. The group work technique promotes the strengthening of the individual and group potential, the value of health, the use of available resources and the exercise of citizenship as important tools in the development of educational practices in health¹⁴.

Nevertheless, there are still difficulties the health professionals need to overcome. Welcoming seems to be a practice some professionals still do not practice. In addition, some nurses' lack of theoretical and practical knowledge in antenatal care and the lack of dialogue between medical professionals and the multi-professional team impairs holistic care for pregnant women¹⁵.

Comprehensive care can be achieved through a focus on local needs and with the interdisciplinary and intersectional resources available. Through the articulation of these factors, popular education, social control, and the promotion of humanized health care are achieved. Educational actions in the care process can be developed in a context which many professionals are inserted in, the interdisciplinary articulation that provides the best quality of health care. In the processes of health education, the need for distinguished and welcoming care is apparent, granting the professionals a more comprehensive tool for care¹⁶.

In Brazil, access to quality care, ensuring access, admission and humanization, is still the privilege of a few women, which implies an increase in problems related to reproductive health, such as the risk of maternal death from pregnancy, childbirth or puerperium¹⁷.

Health professionals' approaches to breastfeeding during antenatal care are crucial to guarantee the exercise of women's right to breastfeed their children, permitting reflection on this practice, knowledge of their rights and preparation for its management. Similarly, care given to women and children in the puerperium should be able to intervene early, welcoming and listening to the difficulties to start this practice, expectations and desires, not only in relation to breastfeeding, but other aspects of their lives, ensuring that comprehensive care is a basic assumption of health care programs for women and children¹⁸.

It is known that educational activities during the antenatal period may be lacking, as women who go through a pregnancy without complications and attend antenatal care still reach the last month showing a lack of knowledge about alterations caused by pregnancy and unprepared to experience childbirth. These results reinforce the concern with regard to the way that antenatal education is being carried out¹⁰.

Let us assume that mothers with less education in years would demand more from health professionals in terms of their doubts and questions on the subject as, in this study the mothers' high level of education was appointed as a positive influence on the duration of EBF.

Maternal education constitutes a factor that affects the motivation to breastfeed. In many developed countries, mothers with higher levels of education tend to breastfeed longer, mainly due to the possibility of greater access to information about the benefits of breastfeeding. In developing countries, underprivileged mothers, often unmarried and uneducated, begin antenatal care later and, consequently, the decision on how to feed the baby also occurs later¹².

The distinction between women who planned the pregnancy and those who did not is perhaps related to greater emotional involvement with pregnancy and the desire to become a mother, which could be an extra motivation for asking health professionals on the subject.

In our study, we found the possibility that the chances of pregnant women being reached by the health education activities increase the extent to which these women join antenatal care services early.

The wider range of guidelines available to first-time mothers may be related to the health professionals' understanding that multiparous women have had greater opportunity to access information and experiences from a previous pregnancy than mothers with their first child. Thus, first-time mothers are preferably chosen for health-education activities.

Given the above, nurses need to realize the importance of antenatal health education to effectively and continuously develop a critical awareness in pregnant women, thus causing the pregnant woman to reflect on their attitudes and gain empowerment in relation to EBF until six months and continuing until two years,

through processes that lead to emancipation, since knowledge enables women to know themselves and decide.

Freire argues that, for the teaching/learning process, the relationship between educator and student should be aggregative, conceiving that one learns from the other, through dialogue and reflections to solve everyday problems. The educator shall comply with the limitations and prior knowledge of the student and share the experiences of their reality¹⁹.

Thus, it is relevant to value the educational strategies with this goal, in view of the singularity of pregnant women's experience and its implications for successful breastfeeding. When considering health education, the subjects involved should be empowered. They should occupy a central position in the process, whose purpose is to promote their emancipation as a way to give significance to the themes addressed. Therefore, in educational practices, women should be considered as active subjects of this process, and not as *tabula rasa* or a deposit of contents of traditional education that reproduces ready and finished concepts¹⁹.

The present study is limited by the factor of seasonality, since data were collected only from the second half of the chosen year. Perhaps, if the data were collected over a full year, we would get different results that would permit generalizations about the population under study. Another factor in our study is the limit related to the location chosen, since we chose the reference hospital for births in the capital Cuiabá - MT (70% of total births), although there are other hospitals that perform less deliveries. Perhaps the inclusion of these other units would allow us to accomplish more reliable generalizations on the capital studied.

FINAL CONSIDERATIONS

Despite the 99.7% coverage rate of antenatal care and 73.4% of pregnant women starting in the first term, we note that 48.9% of postpartum women received guidelines on exclusive breastfeeding during the antenatal period. In total, 62 women (20.3%) participated in lectures/conversation groups. In relation to the professionals who addressed the EBF topic, doctors stood out with 38.2% of the total.

This reveals a major deficiency in the quality of care provided to pregnant women during antenatal monitoring, since the guidelines about pregnancy and, if so, on EBF, make up the minimum required from professionals in this stage of the pregnant woman's life in order to support their decision process.

Regarding the data on the prevalence and prevalence ratio about the association between health education and some significant variables, the following stood out: having eight or more years of study; income up to two minimum wages; planned pregnancies; start of antenatal care in the first term and first pregnancy.

These data demonstrate that there may be a gap in the care given to pregnant women because the prevalence of early weaning found in Cuiabá remains high, despite knowing that information about the benefits of exclusive breastfeeding for mother and infant are increasingly disseminated.

It is noticed that, in the prenatal period, health professionals are responsible for raising the pregnant women's awareness about the benefits of exclusive breastfeeding, as this period begins the welcoming. Such practices are guided through communication, listening, trying to understand the reality of this woman, her socioeconomic, cultural, family context. Accordingly, one should strengthen this professional-patient relationship to establish trust, so as to provide orientations on problem-free breastfeeding, showing the benefits of breastfeeding, developing the pregnant women's critical awareness on the importance of EBF.

In this context, health education can be an indispensable tool because it allows one to develop activities directed at the principles of educating, thereby facilitating the mothers' awareness on these values. For this to happen, the women need to know their own body, because this enables them to develop their autonomy and empowerment.

Without empowerment, the users are unable to make decisions favorable to their health, because they are unaware of their rights and do not have sufficient knowledge to identify the type of care received and to judge what would be most appropriate.

It is necessary to change these practices in order to rethink and reframe educational activities that seek to make changes in the professional/user relationship, aiming at the admission of pregnant women in the context of breastfeeding promotion, protection and support.

REFERENCES

1. Brito RS, Oliveira EMF. Opinião do pai sobre o aleitamento materno. *Rev Rene*. 2006;7(1):9-16.
2. Levy L, Bertolo H. Manual de Aleitamento Materno. Comitê Português para a UNICEF/Comissão Nacional Iniciativa Hospitais Amigos dos Bebês. Edição Revista de 2008. Disponível: http://www.unicef.pt/docs/manual_aleitamento.pdf
3. World Health Organization - WHO. Report of an expert consultation on the optimal duration of exclusive breastfeeding. Geneva (SUI): World Health Organization; 2001.
4. Ministério da Saúde (BR). Secretaria de Atenção à Saúde. Promovendo o Aleitamento Materno. 2ª ed. Brasília (DF): Editora do Ministério da Saúde; 2007.
5. Azeredo CM, Társis MM, Rosa TCA, Silva FF, Cecon PR, Cotta RM. Percepção de mães e profissionais de saúde sobre o aleitamento materno: encontros e desencontros. *Rev. paul. pediatri*. São Paulo, 2008 dez; 26(4):336-44.
6. Fujimori E, Minawaga AT, Laurenti D, Montero RMJM, Borges ALV et al. Duração do aleitamento materno em menores de dois anos de idade em Itupeva, São Paulo, Brasil: há diferenças entre os grupos sociais? *Rev. bras. saude mater. infant*. 2010; 10(1):39-49.
7. Venancio SI, Escuder MML, Saldiva SRDR, Giugliani ERJ. A prática do aleitamento materno nas capitais brasileiras e Distrito Federal: situação atual e avanços. *J. Pediatr*. 2010; 86(4):317-24.
8. Souza VB, Roecker S, Marcon SS. Ações educativas durante a assistência pré-natal: percepção de gestantes atendidas na rede básica de Maringá-PR. *Revista Eletrônica de Enfermagem*. 2011;13(2):199-210.
9. Feijão AR, Galvão MTG. Ações de Educação em Saúde na Atenção Primária: revelando métodos, técnicas e bases teóricas. *Rev Rene*. 2007;8(2):41-9.
10. Rios, CTF, Vieira NFC. Ações educativas no pré-natal: reflexão sobre a consulta de enfermagem como um espaço para educação em saúde. *Cienc. saude colet*. 2007; 12(2):477-86.
11. Ministério da Saúde (BR), Secretaria de Atenção à Saúde. Pre-natal e puerpério: atenção qualificada e humanizada. Brasília (DF): Ministério da Saúde; 2005.
12. Araújo OD, Cunha AL, Lustosa LR, Nery IS, Mendonça RCM, Campelo SMA et al. Aleitamento materno: fatores que levam ao desmame precoce. *Rev. bras. enferm*. 2008 ago;61(4):488-92.
13. Batista KRA, Farias MCAD, Melo WSN. Influência da Assistência de Enfermagem na Prática da Amamentação no Puerpério Imediato. *Saude em debate*. 2013;37(96): 130-8.
14. Santos AL, Radovanovic CAT, Marcon SS. Assistência Pré-natal: satisfação e expectativas. *Rev Rene*. 2010; 11(número especial):61-71.
15. Matos DS, Rodrigues MS, Rodrigues TS. Atuação do enfermeiro na assistência ao pré-natal de baixo risco na estratégia saúde da família em um município de Minas Gerais. *Enferm. rev*. 2013;16(1):18-33.
16. Ceron MI, Barbieri A, Fonseca LM, Fedosse E. Assistência pré-natal na percepção de puérperas provenientes de diferentes serviços de saúde. *Rev. CEFAC [online]*. 2013 jun;15(3): 653-62.
17. Britto, FL. Orientação e incentivo ao aleitamento materno na assistência pré-natal e puerperal: uma revisão de literatura. *Rev. Saúde Públ. Santa Cat*. 2013 jan/mar;6(1):66-80.
18. Damião JJ. Influência da escolaridade e do trabalho maternos no aleitamento materno exclusivo. *Rev. bras. epidemiol. [online]*. 2008 out;11(3):442-52.
19. Freire P. Pedagogia da Autonomia: saberes necessários à práticas educativa. 43ª ed. São Paulo: Paz e Terra; 2011.