



Revista Portuguesa de Pneumologia
ISSN: 0873-2159
sppneumologia@mail.telepac.pt
Sociedade Portuguesa de Pneumologia
Portugal


Nunes Paiva, Dulciane; Bender Assmann, Laíse; Fanfa Bordin, Diogo; Gass, Ricardo; Trevisan Jost, Renan; Bernardo-Filho, Mario; Alves Franca, Rodrigo; Machado Cardoso, Dannuey
Inspiratory muscle training with threshold or incentive spirometry: Which is the most effective?
Revista Portuguesa de Pneumologia, vol. 21, núm. 2, 2015, pp. 76-81
Sociedade Portuguesa de Pneumologia
Lisboa, Portugal

Available in: <http://www.redalyc.org/articulo.oa?id=169737161006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



ORIGINAL ARTICLE

Inspiratory muscle training with threshold or incentive spirometry: Which is the most effective?☆



Dulciane Nunes Paiva^{a,b,*}, Laíse Bender Assmann^b, Diogo Fanfa Bordin^b,
Ricardo Gass^b, Renan Trevisan Jost^c, Mario Bernardo-Filho^d, Rodrigo Alves França^e,
Dannuey Machado Cardoso^b

^a Pos Graduate Course in Health Promotion, University of Santa Cruz do Sul (UNISC), Santa Cruz do Sul, RS, Brazil

^b Physical Therapy Course, University of Santa Cruz do Sul (UNISC), Santa Cruz do Sul, RS, Brazil

^c Faculty of Medicine, Catholic University of Rio Grande do Sul, Porto Alegre, Brazil

^d Department of Biophysics and Biometry, State University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

^e University of Brasília, Brasília, DF, Brazil

Received 27 February 2014; accepted 24 May 2014

Available online 7 February 2015

KEYWORDS

Maximal inspiratory
pressure;
Inspiratory muscular
training;
Incentive spirometry

Abstract Inspiratory muscular training (IMT) increases the respiratory muscle strength, however, there is no data demonstrating its superiority over the incentive spirometry (IS) in doing so. Values of muscle strength after IMT (Threshold IMT[®]) and by the IS (Voldyne[®]) in healthy females was compared. Subjects ($n = 40$) were randomly divided into control group (CG, $n = 14$), IS group (ISG, $n = 13$) and threshold group (TG, $n = 13$). PI_{max} was measured before (pre-IMT), at 15 and 30 days of IMT. There was an increase in PI_{max} of the TG at 15 days ($p < 0.001$) and 30 days of IMT ($p < 0.001$). The same occurred with the ISG, which increased the PI_{max} at 15 days ($p < 0.001$) and 30 days of training ($p < 0.001$). After 30 days of IMT, the TG presented a PI_{max} which was significantly higher than ISG and the CG ($p = 0.045$ and $p < 0.001$, respectively). It can be concluded that IMT by threshold was more effective in increasing muscle strength than the Voldyne.

© 2014 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. All rights reserved.

Introduction

The ability to generate inspiratory pressure can be increased by the formation of a pressure threshold. This threshold can be produced by respiratory muscular training (RMT), which increases resistance to muscular fatigue, improves the

☆ Study conducted at the Laboratory Methods and Evaluation Techniques of the Physiotherapy Course of the University of Santa Cruz do Sul – Rio Grande do Sul, Brazil.

* Corresponding author.

respiratory function and increases the blood flow redistribution to the muscles.¹⁻³

The increment in respiratory muscular strength increases the lung capacity in order to offset the increased respiratory load.^{2,4} However, there are some points about the RMT which are not clear, such as the mechanisms through which this kind of training improves the performance in the maximal incremental exercise,¹ as well as the devices which result in more efficiency.

Among the most commonly used devices for inspiratory muscular training (IMT), Threshold IMT[®] stands out⁵⁻⁸: it has been widely used in the implementation of IMT in several clinical situations, with the aim of increasing exercise capacity in chronic obstructive pulmonary disease (COPD),⁶ in postoperative coronary artery bypass surgery,⁹ preoperative correction of scoliosis in children with Duchenne muscular dystrophy,¹⁰ in post-bariatric surgery¹¹ and weaning from mechanical ventilation.¹² In addition, some studies have reported the use of incentive spirometer Voldyne[®] for purposes of IMT,^{13,14} although this feature is based on the supply of non-linear pressure load. It is worth noting, however, there are not many studies that compare the effectiveness of Threshold IMT[®] compared with Voldyne[®] as a resource for RMT.¹³

Authors on this subject consider that the ideal protocol for evaluating the real effectiveness of the IMT has still not been established.^{15,16} There is a great diversity in suggested training times and in devices used for such purposes.^{7,17-19} For this reason, the present study aimed to compare the effect of IMT on inspiratory muscle strength (IMS) using the Threshold IMT[®] device and the Voldyne[®] in healthy and sedentary subjects.

Methods

This is a randomized clinical trial, comprised of sedentary and healthy subjects (females) aged between 18 and 40 years, with body mass index (BMI) of less than 25 kg/m² and normal lung function. Active smokers were excluded, as were subjects that took regular physical activity. The study was approved by the Ethics Committee of the University of Santa Cruz do Sul, RS, Brazil. All the subjects read and signed an informed consent form.

The subjects were allocated from a list of random numbers generated by computer (Random Number Generator – Pro v2.00, Segobit, USA). The randomization sequence was done by a researcher who was directly not involved in the study and the subjects were randomly divided into control group (CG), threshold group (TG) and incentive spirometry group (ISG). Afterwards, the lung volumes were measured by a spirometer (EasyOne[®], model 2001, Zurich, Switzerland) for the analysis of forced vital capacity (FVC), forced expiratory volume in the first second (FEV₁) and peak expiratory flow (PEF).²⁰ The best curve was compared with the predicted values in the literature.²¹ The anthropometric characteristics of the subjects were also assessed.

The PI_{max} was evaluated with a digital pressure manometer (MDI[®] model MVD300, Porto Alegre, Brazil) having been obtained with the individual breathing in from residual volume (RV) to total lung capacity and effort was made

against the occluded valve, with the measurements carried out in the sitting position.²² The figure considered for data analysis was the highest value among the three maneuvers, if it did not differ more than 10% from the second highest value in descending order.²² To evaluate the effect of the IMT, the PI_{max} was measured at pre-training (pre-IMT), after 15 days and after 30 days.

The IMT was performed with the Threshold IMT[®] (Respironics New Jersey, Inc., USA) and with an incentive spirometry (IS) with a volume-oriented device (Voldyne 5000[®], Sherwood Medical, St. Louis, USA). CG performed breathing exercises twice.²³ Blinding was strictly maintained; the team that carried out the review was not informed about which type of device was being used by the participants not were the participants informed about the device or the form of training used by the other group.

Inspiratory muscle training protocol with Threshold IMT[®]

The TG participants were instructed to perform IMT with the Threshold[®] twice a day, three times a week on non-consecutive days during the 30-day period. There was a series of six repetitions with an initial resistive load of 40% of the PI_{max}. Evaluations of the PI_{max} were not carried out in the first week of IMT to ensure the subjects were familiarized with the established protocol. After this period the PI_{max} measurements were performed weekly with the aim of increasing the resistive inspiratory pressure so as to achieve the training effect. The incremental load was 10% of the PI_{max} per week or enough to reach the maximum resistive value of the Threshold IMT[®], which is 41 cmH₂O.^{4,13}

Inspiratory muscle training protocol with Voldyne[®]

The subjects of the ISG were instructed to perform maximal inspiration from RV to TLC with Voldyne[®] in the sitting position.²⁴ The expiration was performed at functional residual capacity in order to avoid possible hyperventilation; there were intervals of 60s between respiratory maneuvers.²⁵ All subjects received verbal commands as to exactly when to start a new inspiration. The Voldyne[®] was used for 20 min (20 repetitions or one repetition per minute). The total time of this program was 30 days, twice a day, three times per week on nonconsecutive days.

Breathing exercises protocol

The breathing exercise performed by GC was performed 02 times per day with 03 sets of 20 repetitions with a pause of 02 min between each set, 03 times a week for 30 non-consecutive days. In this breathing exercise the individual performs diaphragmatic inspiration from functional residual capacity to maximum inspiratory lung volume with 02 consecutive breaks, while maintaining a ratio of 02 to 01 breaths.²³




Figure 1 Effects of inspiratory muscular training in groups studied. Comparison between pre-IMT at 15 days in threshold group – TG (* $p < 0.001$) and incentive spirometry group – ISG (* $p < 0.001$), and at 30 days in TG (** $p < 0.001$) and ISG (** $p < 0.001$). Comparison between the three groups in pre-IMT ($p = 0.494$), 15 days ($##p < 0.001$) and 30 days ($###p < 0.001$) conditions. Difference between the TG and ISG at 15 days (* $p = 0.328$) and at 30 days (** $p = 0.045$). CG: control group; TG: threshold group; ISG: incentive spirometry group. ANOVA two-way test followed by Bonferroni multiple comparison. Significance was accepted at $p < 0.05$.

Statistical analysis

The SPSS (version 20.0, USA) was used for the analysis of the results which were presented as mean and standard deviation and the normal distribution observed by Shapiro–Wilk test. The data referring to anthropometric variables and to pulmonary function were compared between groups by one-way ANOVA followed by Tukey post hoc. The behavior of the PI_{max} in the different groups with two-way ANOVA followed by Bonferroni multiple comparison and the interaction of the FEV_1 with the PI_{max} by ANCOVA were also observed. Statistical significance of $p < 0.05$ was used.

Results

The subjects were distributed into CG ($n = 14$), TG ($n = 13$) and ISG ($n = 13$) and their characteristics (baseline) are presented in Table 1. The sample being studied was found to be homogeneous, except for the FEV_1 , however, all subjects had values which were close to those predicted.

In relation to IMT, there was a significant increase in the PI_{max} in both TG and ISG at 15 days ($p < 0.001$ and $p < 0.001$, respectively) and at 30 days ($p < 0.001$ and $p < 0.001$, respectively) compared to the pre-IMT. When comparing the effect of IMT between ISG and TG, it was observed that it was only at 30 days was there a higher gain of the PI_{max} by the subjects of the TG in comparison to the subjects of the ISG ($p = 0.045$) (Fig. 1). There was no increase in PI_{max} at 15 ($p = 0.988$) or 30 days ($p = 0.942$) for CG compared to the pre-IMT. In addition, no interaction was observed between the FEV_1 ($p = 0.208$) and the PI_{max} .

Discussion

Our findings showed an increase in the PI_{max} caused training by Voldyne® and Threshold IMT®, although the latter device was more effective in increasing the IMS after 30 days of IMT. Traditionally the Threshold IMT® is the device used for respiratory muscle training as it provides linear pressure load with the possibility of gradual increments.²⁶

Due to the visual feedback the IS with a volume-oriented device encourages the following; the performance of the subject, the realization of sustained maximal inspiration and hyperpnea aiming to gain pulmonary volume.²⁷ It is known that maximum inspiration causes the increase of the transpulmonary pressure and the increase of the pulmonary volume. Furthermore, resting at the end of inspiration keeps up the increase of the transpulmonary pressure and ensures the alveolar stability.²⁸

The literature provides for the use of specific inspiratory incentive guidelines in accordance with the norms of the American Association for Respiratory Care; its use is prioritized as part of prophylactic regimens routine and therapeutic respiratory care.²⁹ However, it should be noted that such devices are rarely used for IMT, which means a lack of empirical standardization.

This research showed that the IMT instituted through the Threshold IMT® caused an increase in the PI_{max} after 15 and 30 days of training. Our findings are confirmed by other studies which suggest the effectiveness of Threshold IMT® in promoting gains in the IMS.^{6,30} Volianitis et al. used IMT for 11 weeks in females rowing athletes and noted an increase in the PI_{max} and in performance during exercise. They have demonstrated that the applicability of the Threshold IMT® is useful not only in clinical approaches but also in improving physical performance.³¹ Likewise, Gosselink et al. have highlighted that the effects of IMT includes delay in respiratory muscle fatigue, redistribution of the blood flow to the respiratory and locomotor muscles and reduction of the perception of discomfort of the respiratory muscles. However, the mechanisms associated with such effects are not very clear.³²

Regarding to the use of Voldyne® was found an increase of PI_{max} in the end of 15 and 30 days of training. Reinforcing our findings, some studies show that the use of this type of IS increases the IMS due to the increased recruitment of the motor units, resulting in greater respiratory muscle strength.^{33,34} However, it should be noted that in clinical practice, this feature is not used for IMT, but as a resource for obtaining pulmonary volume gain.

The IS enables the mobilization of large pulmonary volumes and it is also responsible for increasing the intra-alveolar pressure at the end of inspiration. This condition improves breathing capacity, and challenges the patient with the visual stimulus generated by the device.^{35–38} The increase of the intra-alveolar pressure is proportional to the contractile force of the respiratory muscles, including the diaphragm and accessory muscles of the respiration, justifying the fact that to achieve the TLC, intense muscle activity should occur.³⁹

It is worth noting that in order to perform breathing exercises with an IS, the subjects mobilize large tidal volume associated with a low respiratory rate, tending to increase

Table 1 Baseline characteristics of the participants from groups studied.

	Groups			p-Value
	Control (n = 14)	Threshold (n = 13)	Incentive spirometry (n = 13)	
<i>Antropometrics</i>				
Age (years)	23.86 ± 3.32	22.92 ± 2.06	21.92 ± 1.44	0.101
Weight (kg)	58.07 ± 9.13	60.15 ± 8.65	58.69 ± 7.76	0.813
Height (m)	1.66 ± 0.06	1.68 ± 0.06	1.68 ± 0.08	0.860
BMI (kg/m²)	20.29 ± 2.36	20.85 ± 1.72	20.31 ± 2.05	0.736
<i>Lung function</i>				
FVC (%pred)	95.14 ± 8.95	102.77 ± 11.86	93.77 ± 12.34	0.096
FEV ₁ (%pred)	97.50 ± 9.82	104.77 ± 11.24	94.31 ± 10.33	0.048†
PEF (%pred)	94.14 ± 16.62	102.23 ± 17.60	90.69 ± 18.79	0.146
<i>Respiratory muscular strength</i>				
PI _{max} (%pred)	102.00 ± 6.13	101.62 ± 8.46	100.62 ± 6.94	0.878

BMI: body mass index; FVC: forced vital capacity; FEV₁: forced expiratory volume in the first second; PEF: peak expiratory flow.

[†] Significant difference between threshold group and incentive spirometry group ($p = 0.042$).

ANOVA test with Tukey post hoc. Significance was accepted at $p < 0.05$.

muscle strength due to increased inhalation/exhalation ratio.²⁷

Putting together our findings, the results of this investigation demonstrate the relevance of the Threshold IMT[®] in producing training effect on the inspiratory muscles, as the gain in the muscle strength is proportional to the applied load. According to Komi, changes in the muscular strength during short periods of training with incremental load appear to be due to the improvement of intra- and inter-muscular neural adjustment.⁴⁰ These mechanisms allow better synchronization in the frequency of neuromuscular firing, resulting in greater activation of the motor units.^{41,42}

It is important to consider that the subjects of the CG did not significantly change the PI_{max} with time. This fact demonstrates that the breathing exercise alone is not effective in increasing the IMS. A study of asthmatic children also found that a IMT program using the Threshold[®] IMT significantly increased PI_{max}, which did not happen with the group that only followed a therapeutic exercise program and conventional education for children with asthma.⁴³ This demonstrates that breathing exercises alone are not sufficient to produce significant gain in muscle strength and when applied in isolation do not impose a resistive load on the inspiratory muscles.

There are some limitations to our investigation that need to be considered. We used a volitional test of generation of pressure (PI_{max}) instead of an electrically evoked potential test for the assessment of muscle strength, only healthy subjects were tested and it is not possible to correlate this experimental situation directly with clinical situations involving patients. Only healthy female subjects were available; there was no male representation. Care should also be taken in comparing the effectiveness of these devices, since the training protocols and intensities of training may differ. It is recognized that the physiological effects of the training are related to its intensity. The differences between the training groups are probably related to the intensity and protocols, and the difference seen at 30 days, with

better outcomes in the threshold group could also reflect the growing intensity of the training in this group.

In conclusion, our results showed that the Voldyne[®] device, as well as promoting its known benefits in increasing lung volume, at the same time promotes a similar increased muscle strength to Threshold IMT[®] device, which is designed specifically for the purpose of TMI. However, after 30 days of training increased PI_{max} caused by Threshold IMT[®] became more evident.

Ethical disclosures

Protection of human and animal subjects. The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors have obtained the written informed consent of the patients or subjects mentioned in the article. The corresponding author is in possession of this document.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

1. McConnell AK, Romer LM. Respiratory muscle training in healthy humans: resolving the controversy. *Int J Sports Med.* 2004;25:284–93.

2. Huang CH, Martin AD, Davenport PW. Effects of inspiratory strength training on the detection of inspiratory loads. *Appl Psychophysiol Biofeedback*. 2009;34:17–26.
3. Witt JD, Guenette JA, Rupert JL, McKenzie DC, Sheel AW. Inspiratory muscle training attenuates the human respiratory muscle metaboreflex. *J Physiol*. 2007;584:1019–28.
4. Enright SJ, Unnithan VB. Effect of inspiratory muscle training intensities on pulmonary function and work capacity in people who are healthy: a randomized controlled trial. *Phys Ther*. 2011;91894–905.
5. West CR, Taylor BJ, Campbell IG, Romer LM. Effects of inspiratory muscle training on exercise responses in Paralympic athletes with cervical spinal cord injury. *Scand J Med Sci Sports*. 2013, <http://dx.doi.org/10.1111/sms.12070>.
6. Tout R, Tayara L, Halimi M. The effects of respiratory muscle training on improvement of the internal and external thoraco-pulmonary respiratory mechanism in COPD patients. *Ann Phys Rehabil Med*. 2013;56:193–211.
7. Cavalheri V, Camillo CA, Pitta F, Alves LA, Probst VS, Brunetto AF. Influência do posicionamento de membros superiores sobre os efeitos do treinamento muscular inspiratório de curta duração e alta intensidade em indivíduos jovens saudáveis. *Fisiot Pesq*. 2008;15:367–73.
8. Lin SJ, McElfresh J, Hall B, Bloom R, Farrell K. Inspiratory muscle training in patients with heart failure: a systematic review. *Cardiopulm Phys Ther J*. 2012;23:29–36.
9. Matheus GB, Dragosavac D, Trevisan P, da Costa CE, Lopes MM, Ribeiro GC. Treinamento muscular melhora o volume corrente e a capacidade vital no pós-operatório de revascularização do miocárdio. *Rev Bras Cir Cardiovasc*. 2012;27.
10. Takaso D, Nakazawa T, Imura T, Takahira N, Itoman M, Takahashi K, et al. Surgical management of severe scoliosis with high-risk pulmonary dysfunction in Duchenne muscular dystrophy. *Int Orthop*. 2010;34:401–6.
11. Casali CC, Pereira AP, Martinez JA, de Souza HC, Gastaldi AC. Effects of inspiratory muscle training on muscular and pulmonary function after bariatric surgery in obese patients. *Obes Surg*. 2011;21:1389–94.
12. Cader SA, Vale RG, Castro JC, Bacelar SC, Biehl C, Gomes MC, et al. Inspiratory muscle training improves maximal inspiratory pressure and may assist weaning in older intubated patients: a randomised trial. *J Physiother*. 2010;56:171–7.
13. Fonseca MdA, Cader SA, Dantas EHM, Bacelar SC, Silva EB, Leal SMD. Respiratory muscle training programs: impact on the functional autonomy of the elderly. *Rev Assoc Med Bras*. 2010;56:642–8.
14. Verges S, Renggli AS, Notter DA, Spengler CM. Effects of different respiratory muscle training regimes on fatigue-related variables during volitional hyperpnoea. *Respir Physiol Neurobiol*. 2009;169:282–90.
15. Reid WD, Samra B. Respiratory muscle training for patients with chronic obstructive pulmonary disease. *Phys Ther*. 1995;75:996–1005.
16. Hsiao YL, Chang CC, Chen CM. Profile of hospitalized elderly patients treated for falling. *Int J Gerontol*. 2012;6:42–5.
17. Aznar-Lain S, Webster AL, Canete S, San Juan AF, Lopez Mojares LM, Perez M, et al. Effects of inspiratory muscle training on exercise capacity and spontaneous physical activity in elderly subjects: a randomized controlled pilot trial. *Int J Sports Med*. 2007;28:1025–9.
18. Nosedá A, Carpioux JP, Vandeput W, Prigogine T, Schmerber J. Resistive inspiratory muscle training and exercise performance in COPD patients. A comparative study with conventional breathing retraining. *Bull Eur Physiopathol Respir*. 1987;23:457–63.
19. Charusisin N, Gosselink R, Decramer M, McConnell A, Saey D, Maltais F, et al. Inspiratory muscle training protocol for patients with chronic obstructive pulmonary disease (IMTCO study): a multicentre randomised controlled trial. *BMJ Open*. 2013;3.
20. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J*. 2005;26:319–38.
21. Knudson RJ, Lebowitz MD, Holberg CJ, Burrows B. Changes in the normal maximal expiratory flow-volume curve with growth and aging. *Am Rev Respir Dis*. 1983;127:725–34.
22. American Thoracic Society/European Respiratory S. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med*. 2002;166:518–624.
23. Ide MR, Caromano FA, Dip MAVB, Guerino MR. Exercícios respiratórios na expansibilidade torácica de idosos: exercícios aquáticos e solo. *Fisioter Mov*. 2007;20:33–40.
24. Weindler J, Kiefer RT. The efficacy of postoperative incentive spirometry is influenced by the device-specific imposed work of breathing. *Chest*. 2001;119:1858–64.
25. Machado MdGR. Bases da fisioterapia respiratória: terapia intensiva e reabilitação. Rio de Janeiro: Guanabara Koogan; 2008.
26. Savci S, Degirmenci B, Saglam M, Arkan H, Inal-Ince D, Turan HN, et al. Short-term effects of inspiratory muscle training in coronary artery bypass graft surgery: a randomized controlled trial. *Scand Cardiovasc J*. 2011;45:286–93.
27. Parreira VF, Tomich GM, Britto RR, Sampaio RF. Assessment of tidal volume and thoracoabdominal motion using volume and flow-oriented incentive spirometers in healthy subjects. *Braz J Med Biol Res*. 2005;38:1105–12.
28. Oliveira M, Santos C, Oliveira C, Ribas D. Efeitos da técnica expansiva e incentivador respiratório na força da musculatura respiratória em idosos institucionalizados. *Fisioter Mov*. 2013;26:133–40.
29. Restrepo RD, Wettstein R, Wittnebel L, Tracy M. Incentive spirometry: 2011. *Respir Care*. 2011;56:1600–4.
30. Zanon C, Rodrigues C, Mariano D, Suzan A, Boaventura L, Galvão F. Efeitos do treinamento muscular inspiratório em universitários tabagistas e não tabagistas. *Fisiot Pesqui*. 2012;19:147–52.
31. Volianitis S, McConnell AK, Koutedakis Y, McNaughton L, Backx K, Jones DA. Inspiratory muscle training improves rowing performance. *Med Sci Sports Exerc*. 2001;33:803–9.
32. Gosselink R, De Vos J, van den Heuvel SP, Segers J, Decramer M, Kwakkel G. Impact of inspiratory muscle training in patients with COPD: what is the evidence? *Eur Respir J*. 2011;37:416–25.
33. Shapira N, Zabatin SM, Ahmed S, Murphy DM, Sullivan D, Lemole GM. Determinants of pulmonary function in patients undergoing coronary bypass operations. *Ann Thorac Surg*. 1990;50:268–73.
34. Romanini W, Muller AP, Carvalho KAT, Olandoski M, Faria-Neto JR, Mendes FL, et al. Os efeitos da pressão positiva intermitente e do incentivador respiratório no pós-operatório de revascularização miocárdica. *Arq Bras Cardiol*. 2007;89:94–9.
35. Muller AP, Olandoski M, Macedo R, Costantini C, Guarita-Souza LC. Comparative study between intermittent (Muller Respirometer) and continuous positive airway pressure in the postoperative period of coronary artery bypass grafting. *Arq Bras Cardiol*. 2006;86:232–9.
36. Tomich G, França D, Diório A, Britto R, Sampaio R, Parreira V. Breathing pattern, thoracoabdominal motion and muscular activity during three breathing exercises. *Braz J Med Biol Res*. 2007;40:17–409.
37. Yamaguti WP, Sakamoto ET, Panazzolo D, Peixoto CaC, Cerri GG, Albuquerque AL. Diaphragmatic mobility in healthy subjects during incentive spirometry with a flow-oriented device and with a volume-oriented device. *J Bras Pneumol*. 2010;36:738–45.
38. Santos TV, Ruas G, Sande de Souza LA, Volpe MS. Influence of forward leaning and incentive spirometry on inspired volumes and inspiratory electromyographic activity during

- breathing exercises in healthy subjects. *J Electromyogr Kinesiol.* 2012;22:961–7.
39. Wattie J. Incentive spirometry following coronary artery bypass surgery. *Physiotherapy.* 1998;84:508–14.
40. Komi PV. Training of muscle strength and power: interaction of neuromotoric, hypertrophic, and mechanical factors. *Int J Sports Med.* 1986;7:10–5.
41. Staron RS, Karapondo DL, Kraemer WJ, Fry AC, Gordon SE, Falkel JE, et al. Skeletal muscle adaptations during early phase of heavy-resistance training in men and women. *J Appl Physiol.* 1994;76:1247–55.
42. Jurimae J, Abernethy PJ, Blake K, McEniery MT. Changes in the myosin heavy chain isoform profile of the triceps brachii muscle following 12 weeks of resistance training. *Eur J Appl Physiol Occup Physiol.* 1996;74:287–92.
43. Lima EV, Lima WL, Nobre A, dos Santos AM, Brito LM, Costa MoR. Inspiratory muscle training and respiratory exercises in children with asthma. *J Bras Pneumol.* 2008;34:552–8.