



Revista Portuguesa de Pneumologia

ISSN: 0873-2159

sppneumologia@mail.telepac.pt

Sociedade Portuguesa de Pneumologia
Portugal

Sá-Sousa, A.; Amaral, R.; Morais-Almeida, M.; Araújo, L.; Azevedo, L.F.; Bugalho-Almeida, A.; Bousquet, J.; Fonseca, J.A.

Asthma control in the Portuguese National Asthma Survey

Revista Portuguesa de Pneumologia, vol. 21, núm. 4, julio-agosto, 2015, pp. 209-213

Sociedade Portuguesa de Pneumologia
Lisboa, Portugal

Available in: <http://www.redalyc.org/articulo.oa?id=169741610008>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



BRIEF COMMUNICATION

Asthma control in the Portuguese National Asthma Survey



A. Sá-Sousa^a, R. Amaral^a, M. Morais-Almeida^{b,c}, L. Araújo^{d,e,f}, L.F. Azevedo^{a,e},
A. Bugalho-Almeida^{b,g}, J. Bousquet^{h,i}, J.A. Fonseca^{a,c,e,f,*}

^a Center for Research in Health Technologies and Information Systems – CINTESIS, Universidade do Porto, Porto, Portugal

^b Allergy and Clinical Immunology Department, Hospital CUF-Descobertas, Lisboa, Portugal

^c Sociedade Portuguesa de Alergologia e Imunologia Clínica, Lisbon, Portugal

^d Immunology Department, Faculdade de Medicina da Universidade do Porto, Porto, Portugal

^e Health Information and Decision Sciences Department – CIDES, Faculdade de Medicina, Universidade do Porto, Porto, Portugal

^f Allergy Unit, Instituto CUF Porto e Hospital CUF Porto, Porto, Portugal

^g Comissão de Acompanhamento do Programa Nacional de Controlo da Asma, Lisbon, Portugal

^h Hôpital Arnaud de Villeneuve, Centre Hospitalier Universitaire Montpellier, Montpellier, France

ⁱ Centre de recherche en Epidémiologie et Santé des Populations – CESP Inserm U1018, Villejuif, France

Received 2 May 2014; accepted 19 August 2014

Available online 3 March 2015

KEYWORDS

Asthma;
Disease management;
Cross-sectional
studies;
Quality of life;
Control perception

Abstract

Introduction: We aimed (1) to measure asthma control using a structure-questionnaire and patient self-perception of asthma-control in the Portuguese National Asthma Survey (INAsma) and (2) to study the relationship between asthma control and asthma-related quality of life.

Methods: We analyze data of asthma patients from a cross-sectional, nationwide telephone interview study – INAsma. Controlled asthma was defined as CARAT global score >24 or CARAT lower airways score ≥ 16 . Mini-AQLQ was used to measure quality of life.

Results: Two hundred and seven (56.9% [95%CI: 51.8–62.0]) of the 364 patients had controlled asthma. Most patients with non-controlled asthma (88%) perceived their disease as controlled. Patients with controlled asthma presented higher mini-AQLQ scores (median, P25–P75; 6.6, 6.0–6.9) than those with non-controlled asthma (4.9, 3.7–5.7) ($p < 0.001$) and a significant positive correlation between CARAT and mini-AQLQ scores was observed ($r = 0.706$; $p < 0.001$).

Conclusion: More than half of the Portuguese patients presented controlled asthma and showed significantly better asthma-related quality of life. Almost 9 out of 10 patients with non-controlled disease have poor perception of their asthma control, which may hinder them from seeking better asthma control.

© 2014 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: jfonseca@med.up.pt (J.A. Fonseca).

Introduction

Proper asthma therapy and education reduces the socioecomic burden of asthma and improves patients' quality of life.¹ However, the proportion of patients achieving asthma control remains low worldwide without substantial improvement in recent years.^{2–6}

The Portuguese National Program for Asthma Control conducted by the Directorate-General of Health between 2005 and 2010, aimed to increase the proportion of asthma patients with controlled disease, and to decrease the personal and community burden of the disease.

At the end of this program, in 2010, we conducted the first Portuguese National Asthma Survey – *Inquérito Nacional sobre Asma* (INAsma), which consisted of two phases. In the first phase, aiming to evaluate asthma prevalence, we estimated that 7% of the Portuguese population had current

asthma.⁷ In the second phase we focused on asthma patients addressing disease control.

We aim to measure asthma control using a structure-questionnaire and patient self-perception of asthma-control in the Portuguese National Asthma Survey (INAsma) and to study the relation of asthma control with asthma-related quality of life (ARQoL).

Methods

The INAsma was a cross-sectional, nationwide, telephone interview study. Sample size calculations details have been previously reported.⁷ All the participants identified with possible asthma in the first phase of the survey were eligible for participation in the second phase. Detailed information is provided in the Supplementary Material. The study was approved by the Hospital Ethics Committee of *Hospital de São João (Porto, Portugal)*.

Table 1 Socio-demographic characteristics of the participants with classification for asthma control by asthma control status and patient perception of control and asthma related quality of life.

| | Total (n = 364) | Controlled asthma | | Patient perception of control | | ARQoL cut-off value | |
|--------------------------------------|--------------------|-------------------|--------------|-------------------------------|--------------------------------|---------------------|-------------------|
| | | Yes (n = 207) | No (n = 157) | Controlled (n = 338) | Non- controlled (n = 21) | Above (n = 188) | Below (n = 89) |
| Sex, n (%) | | | | | | | |
| Male | 155 (42.6) | 111 (71.6) | 44 (28.4) | 146 (96.1) | 6 (3.9) | 79 (81.4) | 18 (18.6) |
| Female | 209 (57.4) | 96 (45.9) | 113 (54.1) | 192 (92.8) | 15 (7.2) | 109 (60.6) | 71 (39.4) |
| Age groups, n (%) | | | | | | | |
| <18 years old | 87 (23.9) | 59 (67.8) | 28 (32.2) | 82 (96.0) | 4 (4.7) | 0 (0.0) | 0 (0.0) |
| 18–64 years old | 177 (48.6) | 112 (63.3) | 65 (36.7) | 168 (96.0) | 7 (4.0) | 132 (74.6) | 45 (25.4) |
| >64 years old | 100 (27.5) | 36 (6.0) | 64 (64.0) | 88 (89.8) | 10 (10.2) | 56 (56.0) | 44 (44.0) |
| Education level ^a , n (%) | | | | | | | |
| <9 years | 240 (65.9) | 120 (50.0) | 120 (50.0) | 220 (92.4) | 18 (7.6) | 92 (57.5) | 68 (42.5) |
| 9–12 years | 83 (22.8) | 57 (68.7) | 26 (31.3) | 80 (98.8) | 1 (1.2) | 60 (78.9) | 16 (21.1) |
| >12 years | 41 (11.3) | 30 (73.2) | 11 (26.8) | 38 (95.0) | 2 (5.0) | 36 (87.8) | 5 (12.2) |
| SES ^b , n (%) | | | | | | | |
| Low | 59 (16.3) | 18 (30.5) | 41 (69.5) | 51 (87.9) | 7 (5.9) | 29 (49.2) | 30 (50.8) |
| Medium low | 222 (61.2) | 127 (57.2) | 95 (42.8) | 211 (95.9) | 9 (4.1) | 116 (69.5) | 51 (30.5) |
| Medium high | 49 (13.5) | 35 (71.4) | 14 (28.6) | 44 (91.7) | 4 (8.3) | 27 (84.8) | 5 (15.6) |
| High | 33 (9.1) | 27 (81.8) | 6 (18.2) | 31 (96.9) | 1 (3.1) | 15 (83.3) | 3 (16.7) |
| Smoking status, n (%) | | | | | | | |
| Non-smoker | 274 (75.3) | 153 (55.8) | 121 (44.2) | 255 (93.8) | 17 (6.3) | 121 (64.0) | 68 (36.0) |
| Ex-smoker | 53 (14.6) | 33 (62.3) | 20 (37.7) | 46 (92.0) | 4 (8.0) | 39 (73.6) | 14 (26.4) |
| Current smoker | 37 (10.2) | 21 (56.8) | 16 (43.2) | 37 (100.0) | 0 (0.0) | 28 (80.0) | 7 (20.0) |
| Controlled asthma, n (%) | | | | | | | |
| Yes | 207 (56.9) | – | – | 201 (98.5) | 3 (1.5) | 137 (92.6) | 11 (7.4) |
| No | 157 (43.1) | – | – | 137 (88.4) | 18 (85.7) | 51 (39.5) | 78 (60.5) |
| Patient perception of control, n (%) | | | | | | | |
| Controlled | 338 (94.2) | 201 (59.5) | 137 (40.5) | – | – | 181 (70.7) | 75 (29.3) |
| Not controlled | 21 (5.8) | 3 (14.3) | 18 (85.7) | – | – | 3 (17.6) | 14 (82.4) |

Asthma Related Quality of Life (ARQoL) cut-off value of 5.4.

^a A total of 60 (2.7%) participants were preschoolers (n = 20) – Data retrieved from 1st phase of INAsma.

^b Socioeconomic Status was categorized in high (A social class), medium high (B social class), medium low (C social classes) and low (D social class) based on occupation and school education of the person who contributes more for the household income – Data retrieved from 1st phase of INAsma.




Figure 1 Relationship between CARAT and mini-AQLQ scores ($n=274$); (A) ARQoL by asthma control groups; Scatterplots show the correlation between (B) CARAT global score and ARQoL; (C) CARAT superior airways subscore and ARQoL; (D) CARAT inferior airways subscore and ARQoL. In A the horizontal line represents the Mini-AQLQ cut-off of 5.4¹¹ NCA: non-controlled asthma; CA: Controlled asthma. In (B, C and D) the vertical lines represent the Mini-AQLQ cut-off of 5.4¹¹ and the horizontal lines represent the CARAT cut-offs; (B) global score cut-off of 24; (C) superior airways subscore cut-off of 8 and (D) the CARAT inferior subscore cut-off of 16. The scatterplots are divided into four quadrants, the inferior left corresponds to participants with non-controlled disease and more impairments and the upper right to participants with controlled disease and less impairments.

Current asthma was defined as self-report of asthma and one of the following: at least one medical appointment due to asthma in the last 12 months; current use of asthma medication; asthma symptoms in the last 12 months (wheeze, cough, breathlessness or sputum production without a cold or respiratory infection).

Asthma control was defined based on Control of Allergic Rhinitis and Asthma Test (CARAT) scores.^{8,9} Asthma was considered controlled for CARAT global score above 24 or CARAT lower airways score of 16 or above. Self-perception of asthma control was evaluated by the answer to the question "Do you believe your asthma was well controlled in the last 4 weeks?". Additional questions were included assessing socio-educational variables.

ARQoL was measured by mini-Asthma Quality of Life Questionnaire (mini-AQLQ)¹⁰ with a cut-off value of 5.4 set according to a previous study in northern Portugal.¹¹

Statistical analyses were performed using IBM SPSS Statistics v21 (2012 SPSS Inc., IBM Company, Chicago, US). Categorical variables were described with absolute frequencies, proportions and 95% Confidence Interval (95% CI)

and continuous variables were described with measures of central tendency and dispersion as appropriate. Correlation between CARAT and mini-AQLQ scores was tested with Pearson's Correlation Coefficient, with a p -value of <0.05 considered as statistically significant.

Results

Of the 707 participants identified with possible asthma in the Phase 1, 559 agreed to participate in Phase 2, from which 405 had current asthma as described in the online supplement (Fig. S1). Sufficient data to define asthma control were available for 364 patients. Participants' characteristics are summarized in Table 1.

According to the patients' answers to CARAT, the most frequent symptoms were related to upper airways with 24.9% ($n=90$) and 21.4% ($n=78$) of the patients answering "Always", to sneezing and itchy nose, respectively (Fig. S2). The less common symptom reported was nocturnal awakening with 69.5% ($n=253$) of the patients answering "Never".

Asthma was controlled in 56.9% [95%CI, 51.8–62.0] ($n=207$) of the patients (Table 1). Regarding patient self-perception of control, 94% ($n=338$) reported having their asthma controlled (Table 1). Among the non-controlled patients 88.4% ($n=137$) perceived their asthma as controlled (Table 1, Fig. S3).

Controlled patients had higher mini-AQLQ scores than the non-controlled (median, P25–P75; 6.6, 6.0–6.9 and 4.9, 3.7–5.7, respectively; $p<0.001$) (Fig. 1A). A significant positive correlation between CARAT and mini-AQLQ scores was observed ($r=0.706$; $p<0.001$, Fig. 1B).

Discussion

We report the first nationwide results on asthma control in Portugal with more than half (57%) of the patients presenting controlled asthma. A significant positive correlation was found between CARAT and mini-AQLQ, indicating that patients with controlled asthma had significant better ARQoL comparing to patients with non-controlled asthma. However, most patients (88%) with non-controlled asthma perceived their asthma as controlled.

The proportion of asthma control was in accordance with recent studies in other countries.⁴ In 2010, the European National Health and Wellness Survey⁵ estimated a global proportion of non-controlled asthma of 53.5% across five European countries. Similarly, the International Asthma Patient Insight Research (INSPIRE) study⁴ found a worldwide proportion of non-controlled asthma of 51%. In Portugal, a study of asthma patients from one Allergy unit concluded that 57% had partially or complete asthma control when assessed with ACT.¹²

A strong positive correlation was observed between asthma control assessed by CARAT and ARQoL assessed by Mini-AQLQ. This correlation was similar to previous reports using other asthma control questionnaires^{11,13,14} but had not been reported with CARAT questionnaire.

Most of the participants with non-controlled asthma perceived their disease as controlled. This is very similar to the data from the Asthma Insights and Reality in Europe survey.³ Also in the INSPIRE study,⁴ patient perception of asthma control was strikingly different from that based on symptoms assessment. These data suggest that patients, especially those with poor control, continue to have very low expectations regarding asthma outcomes. Clinical studies are warranted on the relation between asthma control and patient-self perception of control.

The main strength of the present study is that INAsma is the first Portuguese study to include patients from the general population from all municipalities and all age groups.⁷ Some limitations are inherent to a landline telephone-based interview study, especially with the recent shift from landline telephones to cellphones. Other limitation is the potential for confusion as the definition of control may have been influenced by the rhinitis subscore. However, we compared the results of patients with and without associated rhinitis and there were no differences in the proportion of patients with controlled asthma, in perceptions of asthma control and in the correlation between asthma control and ARQoL (data not shown). Moreover, the CARAT superior airways subscore seems not to have affected

the correlation between CARAT total score and miniARQLQ (Fig. 1C).

In conclusion, more than half of the Portuguese asthma patients have their disease controlled and these have significantly better asthma-related quality of life. Poor perception of control seems to be an obstacle to achieving better disease control as almost 9 out of 10 patients with non-controlled disease had poor perception of their asthma control. The objective assessment of asthma control using validated tools may help patients to increase their expectations regarding asthma outcomes.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflicts of interest

The authors have no conflicts of interest to declare.

Acknowledgements

The Portuguese National Asthma Survey – *Inquérito Nacional sobre Asma* (INAsma) was conducted by the Center for Health Technology and Services Research (CINTESIS) in collaboration with *Sociedade Portuguesa de Pneumologia*, by appointment of the Portuguese Health Directorate.

The work presented in this short communication was partially funded by FEDER funds through COMPETE (*Programa Operacional Factores de Competitividade*) and by National funds through a FCT (*Fundação para a Ciência e a Tecnologia*) project with reference “PTDC/SAU-SAP/119192/2010”.

Appendix A. Supplementary data

Supplementary material associated with this article can be found in the online version available at <http://doi.org/10.1016/j.rppneu.2014.08.001>.

References

- Bateman ED, Bousquet J, Keetch ML, Busse WW, Clark TJ, Pederesen SE. The correlation between asthma control and health status: the GOAL study. *Eur Respir J*. 2007;29:56–62.
- Gold LS, Yeung K, Smith N, Allen-Ramey FC, Nathan RA, Sullivan SD. Asthma control, cost and race: results from a national survey. *J Asthma*. 2013;50:783–90.
- Rabe K, Vermeire P, Soriano J, Maier W. Clinical management of asthma in 1999: the Asthma Insights and Reality in Europe (AIRE) study. *Eur Respir J*. 2000;16:802–7.
- Partridge MR, van der Molen T, Myrseth S-E, Busse WW. Attitudes and actions of asthma patients on regular maintenance therapy: the INSPIRE study. *BMC Pulm Med*. 2006;6:13.

5. Demoly P, Annunziata K, Gubba E, Adamek L. Repeated cross-sectional survey of patient-reported asthma control in Europe in the past 5 years. *Eur Respir Rev.* 2012;21:66–74.
6. Slejko JF, Ghushchyan VH, Sucher B, Globe DR, Lin S-L, Globe G, et al. Asthma control in the United States, 2008–2010: indicators of poor asthma control. *J Allergy Clin Immunol.* 2013;133:1579–87.
7. Sa-Sousa A, Morais-Almeida M, Azevedo LF, Carvalho R, Jacinto T, Todo-Bom A, et al. Prevalence of asthma in Portugal – The Portuguese National Asthma Survey. *Clin Transl Allergy.* 2012;2:15.
8. Fonseca JA, Nogueira-Silva L, Morais-Almeida M, Azevedo L, Sa-Sousa A, Branco-Ferreira M, et al. Validation of a questionnaire (CARAT10) to assess rhinitis and asthma in patients with asthma. *Allergy.* 2010;65:1042–8.
9. Fonseca JA, Nogueira-Silva L, Morais-Almeida M, Sa-Sousa A, Azevedo LF, Ferreira J, et al. Control of Allergic Rhinitis and Asthma Test (CARAT) can be used to assess individual patients over time. *Clin Transl Allergy.* 2012;2:16.
10. Juniper EF, Guyatt GH, Cox FM, Ferrie PJ, King DR. Development and validation of the Mini Asthma Quality of Life Questionnaire. *Eur Respir J.* 1999;14:32–8.
11. Correia de Sousa J, Pina A, Cruz A, Quelhas A, Almada-Lobo F, Cabrita J, et al. Asthma control, quality of life, and the role of patient enablement: a cross-sectional observational study. *Prim Care Respir J.* 2013;22:181–7.
12. Leblanc A, Botelho C, Coimbra A, da Silva J, de Castro E, Cernadas J. Assessment of asthma control: clinical, functional and inflammatory aspects. *Eur Ann Allergy Clin Immunol.* 2013;45:90–6.
13. Schatz M, Mosen DM, Kosinski M, Vollmer WM, Magid DJ, O'Connor E, et al. The relationship between asthma-specific quality of life and asthma control. *J Asthma.* 2007;44:391–5.
14. Siroux V, Boudier A, Bousquet J, Vignoud L, Gormand F, Just J, et al. Asthma control assessed in the EGEA epidemiological survey and health-related quality of life. *Respir Med.* 2012;106:820–8.