



Revista Portuguesa de Pneumologia  
ISSN: 0873-2159  
sppneumologia@mail.telepac.pt  
Sociedade Portuguesa de Pneumologia  
Portugal

Santos, V.; Magalhães, A.; Sucena, M.; Fernandes, G.  
The role of endobronchial ultrasound-guided transbronchial needle aspiration in the  
diagnosis of lymphoproliferative disorders  
Revista Portuguesa de Pneumologia, vol. 23, núm. 6, 2017, pp. 362-363  
Sociedade Portuguesa de Pneumologia  
Lisboa, Portugal

Available in: <http://www.redalyc.org/articulo.oa?id=169754669012>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

M. Costa e Silva<sup>a</sup>, R. Rolo<sup>b</sup>

<sup>a</sup> Serviço de Pneumologia, Centro Hospitalar de Gaia/Espinho, Vila Nova de Gaia, Portugal

<sup>b</sup> Serviço de Pneumologia, Hospital de Braga, Braga, Portugal

1 June 2017 5 September 2017 15 October 2017

<https://doi.org/10.1016/j.rppnen.2017.10.003>  
2173-5115/

© 2017 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## The role of endobronchial ultrasound-guided transbronchial needle aspiration in the diagnosis of lymphoproliferative disorders



Dear Editor,

Endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) is a minimally invasive bronchoscopic technique included in the initial diagnostic workup of mediastinal and hilar lymphadenopathy and masses of unknown etiology.<sup>1,2</sup>

The value of EBUS-TBNA in the diagnosis of lymphoproliferative disorders (LPD) is not well established.<sup>1-8</sup> LPD diagnosis requires lymph node sampling for pathologic and flow cytometry analysis. Are the samples obtained by EBUS-TBNA good enough to provide the diagnosis of LPD?

The usefulness of EBUS-TBNA in the diagnosis of LPD in patients with lymph nodes or mediastinal masses and clinical suspicion of LPD was evaluated.

One hundred and eighty nine patients underwent EBUS-TBNA to investigate the etiology of MLN between 2009 and 2012. The suspicion of LPD was based on past history of lymphoma or clinical manifestations and associated with unexplained lymph nodes. EBUS-TBNA biopsies were performed under general anesthesia. The ecoendoscope used was the Olympus-BF-TYPE-UC-180F, with a convex transducer (7.5MHz) and integrated Doppler mode. A dedicated

22-gauge needle (NA-2015X-4022-Olympus) was used to perform TBNA. The ultrasound processor was Olympus-EU-ME1. At least 3 or 4 punctures were made in each target station without rapid on-site cytology. Immunohistochemistry and flow cytometry were requested on all samples. The sample was considered adequate if lymphocytes, antracosis or a specific diagnosis was identified. A true positive diagnosis was made if there was a positive biopsy consistent with lymphoma by EBUS-TBNA or on subsequent follow-up by another method within 6 months. A true negative diagnosis was considered with a negative biopsy and stability or regression of the lymph nodes over a period of at least 6 months.

6.9% (13) patients had clinical suspicion of LPD. 61.5% were male, with mean age of  $58.4 \pm 15.9$  years. Nine patients had enlarged mediastinal lymph nodes and four mediastinal masses.

The EBUS-TBNA results were: three LPD, two granulomatous disease, four reactive lymph nodes and four inadequate samples. Among patients with non-diagnostic results (8), 37.5% (3) had LPD: one patient repeated EBUS-TBNA, one underwent a mediastinoscopy and one patient was diagnosed with LPD by peripheral lymph node biopsy. The diagnosis of reactive lymph node was confirmed in three patients after clinical and radiological monitoring (Table 1).

The diagnosis of LPD was established in six patients. The sensitivity, negative predictive value and diagnostic accuracy of EBUS-TBNA in the diagnosis of LPD were 50, 70 and 76.9%, respectively.

In the literature, the yield of EBUS-TBNA in patients with lymphoma is not well defined with variable results;

**Table 1** Endobronchial ultrasound needle aspiration results and method of final diagnosis in 13 patients with mediastinal lymph node with suspicion of lymphoma.

Patient no.	EBUS-TBNA diagnosis	Other procedure	Definitive diagnosis
# 1	Hodgkin lymphoma	-	Hodgkin lymphoma
# 2	B-cell lymphoma	-	Non-Hodgkin lymphoma
# 3	Hodgkin lymphoma	-	Hodgkin lymphoma
# 4	Reactive lymph node	Peripheral lymph node biopsy	Non-Hodgkin lymphoma
# 5	Inadequate sample	Mediastinoscopy	Hodgkin lymphoma
# 6	Reactive lymph node	Repeat EBUS-TBNA	Non-Hodgkin lymphoma
# 7	Inadequate sample	Repeat EBUS-TBNA	Reactive lymph node
# 8	Inadequate sample	Repeat EBUS-TBNA	Reactive lymph node
# 9	Granulomatous disease	-	Tuberculosis
# 10	Granulomatous disease	-	Sarcoidosis
# 11	Inadequate sample	Clinical/Radiological monitoring - resolution	Reactive lymph node
# 12	Reactive lymph node	Clinical/Radiological monitoring - resolution	Reactive lymph node
# 13	Reactive lymph node	Clinical/Radiological monitoring - resolution	Reactive lymph node

in the few studies published they vary between 22%<sup>1</sup> and 91%.<sup>2</sup>

In this retrospective case study analysis, the prevalence of LPD was low (46.2%), with a low pretest probability in some cases, which could have hampered sensitivity.

According the Chest guideline about EBUS-TBNA, trials evaluating needle size were reviewed and both 21 and 22 gauge needles are considered adequate.<sup>6</sup> The role of 19 gauge needles is yet to be defined, but it is likely that larger needles will provide superior sample quality and improve the yield and sensitivity of EBUS-TBNA in patients with mediastinal adenopathy secondary to lymphoma.

No data exist regarding the number of needle passes required to obtain a sufficient diagnostic yield for lymphoma.<sup>6,8</sup> In the absence of rapid on-site evaluation (ROSE), usually 4 passes were done, the latter was used for immunophenotyping.<sup>4,7,8</sup>

The present study demonstrated that EBUS-TBNA has an important role in the diagnostic workup of mediastinal involvement of LPD. If available, it should be included in the initial evaluation because mediastinoscopy or thoracoscopy pose higher risk to the patient and increased cost. Considering the low negative predictive value of EBUS-TBNA for the diagnosis of lymphoma, patients with inconclusive results must be submitted to further diagnostic investigation.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## References

- Iqbal S, DePew ZS, Kurtin PJ, Sykes AM, Johnson GB, Edell ES, et al. Endobronchial ultrasound and lymphoproliferative disorders: a retrospective study. *Ann Thorac Surg.* 2012;94:1830–4.
- Kennedy MP, Jimenez CA, Bruzzi JF, Mhatre AD, Lei X, Giles FJ, et al. Endobronchial ultrasound-guided transbronchial needle aspiration in the diagnosis of lymphoma. *Thorax.* 2008;63:360–5.
- Steinfors DP, Conron M, Tsui A, Pasricha SR, Renwick WE, Antippa P, et al. Endobronchial ultrasound-guided transbronchial needle aspiration for the evaluation of suspected lymphoma. *J Thorac Oncol.* 2010;5:804–9.
- Marshall CB, Jacob B, Patel S, Sneige N, Jimenez CA, Morice RC, et al. The utility of endobronchial ultrasound-guided transbronchial needle aspiration biopsy in the diagnosis of mediastinal lymphoproliferative disorders. *Cancer Cytopathol.* 2011;119:118–26.
- Senturk A, Babaoglu E, Kilic H, Hezer H, Dogan HT, Hasanoglu HC, et al. Endobronchial ultrasound-guided transbronchial needle aspiration in the diagnosis of lymphoma. *Asian Pac J Cancer Prev.* 2014;15:4169–73.
- Wahidi MM, Herth F, Yasufuku K, Shepherd RW, Yarmus L, Chawla M, et al. Technical aspects of endobronchial ultrasound-guided transbronchial needle aspiration: Chest guideline and expert panel report. *Chest.* 2016;149:816–35.
- Grosu HB, Iliesiu M, Caraway NP, Medeiros LJ, Lei X, Jimenez CA, et al. Endobronchial ultrasound-guided transbronchial needle aspiration for the diagnosis and subtyping of lymphoma. *Ann Am Thorac Soc.* 2015;12:1336–44.
- Moonim MT, Breen R, Fields PA, Santis G. Diagnosis and subtyping of de novo and relapsed mediastinal lymphomas by endobronchial ultrasound needle aspiration. *Am J Respir Crit Care Med.* 2013;188:1216–23.

V. Santos<sup>a,\*</sup>, A. Magalhães<sup>a</sup>, M. Sucena<sup>a</sup>, G. Fernandes<sup>a,b</sup>

<sup>a</sup> *Pulmonology Department, Centro Hospitalar de São João, Porto, Portugal*

<sup>b</sup> *Faculty of Medicine of Porto University, Porto, Portugal*

\*Corresponding author.

E-mail address: [vferreirads@gmail.com](mailto:vferreirads@gmail.com) (V. Santos).

<https://doi.org/10.1016/j.rppnen.2017.07.007>  
2173-5115/

© 2017 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Beneficial effects of pulmonary rehabilitation in adult asthma



Dear Editor,

It is well established that exercise improves physical fitness without any adverse effects on asthma control.<sup>1,2</sup> Asthma patients present lower levels of physical fitness and cardiopulmonary conditioning than their peers.<sup>1</sup> Low physical activity levels are associated with more symptoms, a higher risk of exacerbations and a lower health related quality of life in asthma patients.<sup>2</sup>

Physical exercise improves cardio-pulmonary fitness, muscle trophism, muscle vascularization, and muscle ability to extract/use oxygen, resulting in a lowering of respiratory rate during activity.<sup>1</sup> With a lower respiratory rate during activity, bronchoconstriction during exercise becomes less likely.<sup>1,3</sup> Recent studies have also identified that regular

exercise resulted in a reduction of airway inflammation, which in turn improved bronchioles patency.<sup>4</sup>

Although pulmonary rehabilitation programs have a well established role in chronic obstructive pulmonary disease treatment,<sup>5,6</sup> studies about pulmonary rehabilitation effects on asthma patients are sparse.

A retrospective study was performed in order to assess the effect of a supervised pulmonary rehabilitation program on asthma symptoms (dyspnea), functional capacity and health related quality of life in adult asthma patients.

The intervention consisted of a pulmonary rehabilitation program with medical supervision. All asthma patients admitted to the pulmonary rehabilitation program between 2010 January and 2016 July were included.

The rehabilitation program lasted 8–12 weeks and was held twice a week. Each session had a duration of 1 h and included a warm up period with breathing pattern re-education exercises, aerobic training (bicycle or