



Archivos de Medicina Veterinaria

ISSN: 0301-732X

archmv@uach.cl

Universidad Austral de Chile

Chile

Muller, M; Ilardi, P; Avendaño-Herrera, R

Efficacy of a commercial disinfectant against *Vibrio ordalii*, *Vibrio anguillarum*, *Francisella* sp. and Infectious Pancreatic Necrosis Virus (IPNV) pathogens of Atlantic salmon (*Salmo salar*) farmed in Chile

Archivos de Medicina Veterinaria, vol. 43, núm. 1, 2011, pp. 73-78

Universidad Austral de Chile

Valdivia, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=173019392010>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Efficacy of a commercial disinfectant against *Vibrio ordalii*, *Vibrio anguillarum*, *Francisella* sp. and Infectious Pancreatic Necrosis Virus (IPNV) pathogens of Atlantic salmon (*Salmo salar*) farmed in Chile

Eficacia de un desinfectante sobre *Vibrio ordalii*, *Vibrio anguillarum*, *Francisella* sp. y Virus de la Necrosis Pancreática Infecciosa (IPNV), patógenos de salmón del Atlántico (*Salmo salar*) cultivado en Chile

M Muller^a, P Ilardi^b, R Avendaño-Herrera^{c*}

^aAQUAFARMA, División de Acuicultura, Veterquímica S.A., Puerto Montt, Chile.

^bLaboratorio de Investigación y Desarrollo-Veterquímica, Cerrillos, Santiago, Chile.

^cUniversidad Andrés Bello, Facultad de Ciencias Biológicas, Departamento de Ciencias Biológicas, Viña del Mar, Chile.

RESUMEN

En el presente trabajo se evaluó la eficacia *in vitro* del desinfectante Duplalin[®], una combinación sinérgica de glutaraldehído y sales de amonio cuaternario de cuarta generación, contra 4 patógenos de peces prevalentes de la salmonicultura chilena. Los resultados muestran que todas las concentraciones ensayadas (diluciones entre 1:200 a 1:400) fueron eficaces sobre los aislados de *Vibrio ordalii* y *Vibrio anguillarum* post-30 s de exposición, detectando niveles de reducción igual a 1.8×10^6 UFC/ ml. Concentraciones superiores de Duplalin[®] (dilución 1:50) y un tiempo de exposición no menor a 5 min. fueron necesarios para eliminar completamente al patógeno intracelular *Francisella* sp. Cuando el desinfectante fue ensayado contra el Virus de la Necrosis Pancreática Infecciosa (IPNV), se detectó que la dilución 1:400 tiene un efecto significativo después de 2 minutos sin importar los títulos de IPNV testeados (mayor concentración evaluada $10^{7.6}$ TCID₅₀/ ml). Duplalin[®] se evaluó en condiciones masivas contra los miembros de la familia *Vibrionaceae*. En comparación a los controles (sin adición desinfectante), la dilución 1:400 de Duplalin[®] eliminó completamente *V. ordalii* y *V. anguillarum* después de 15 minutos de tratamiento, tanto en el agua de cultivo como en la superficie de mallas usadas en el cultivo del salmón. Así, el análisis microbiológico del agua de los controles mostró concentraciones de $1.4 \pm 0.3 \times 10^6$ UFC/ ml, mientras en el caso de las mallas $7.6 \pm 3.2 \times 10^5$ UFC/ ml¹. En resumen, los antecedentes obtenidos indican que el uso del desinfectante Duplalin[®] es efectivo contra *V. ordalii*, *V. anguillarum* y IPNV en bajas concentraciones y cortos periodos de exposición (dilución 1:400 por 15 min.), mientras que para el patógeno intracelular se requiere una concentración mayor.

Key words: disinfectant, Chilean fish pathogens, Atlantic salmon.

Palabras clave: desinfectante, patógenos de peces, salmón del Atlántico.

INTRODUCTION

Fish diseases of major importance in Chilean salmon farming includes infectious pancreatic necrosis (IPN) and *Piscirickettsia salmonis* (Fryer and Hedrick 2003), which cause significant economic losses due to mortalities of Atlantic salmon, *Salmo salar*, fry and post-smolt in seawater. In addition, an increase in disease detection of emergent pathogens as *Vibrio ordalii* (Colquhoun *et al* 2004), *Streptococcus phocae* (Gibello *et al* 2005, Romalde *et al* 2008), *Francisella* (Birkbeck *et al* 2007) and, lately, *Vibrio anguillarum* (Avendaño-Herrera *et al* 2007) has been observed in the last decade. To date, economic losses caused by these bacterial pathogens, except *V. ordalii*, have decreased; however, it remains one of the

most important pathogen in the salmon industry in Chile because no commercial vaccines are available. In general, most treatments proposed for the outbreaks are based on the administration of drugs through feed. Considering that a selective effect of antimicrobial use on the emergence of resistant fish bacteria has been documented in several reports (Smith *et al* 1994, Alderman and Hastings 1998), alternative treatments to the use of drugs are necessary.

In salmonids aquaculture facilities, disinfectants are vital tools for effective farm biosecurity, used to reduce pathogenic microorganisms on biological surfaces by rather non-specific actions. They include iodophores used for eggs and equipment disinfectants, salts, organic chlorocompounds (e.g. chloramines-T), aldehydes (e.g. formalin), hydrogen peroxide and quaternary ammonium compounds (e.g. benzalkonium chloride) (Burka *et al* 1997). The majority of these chemicals have also been used to inactivate bacterial and virus pathogens on contaminated rearing equipment, seawater pipes, air hoses, tanks, and nets as well as the hands and feet of the working staff. Besides its disinfecting

properties, few studies have been designed to assess the quaternary ammonium compounds, particularly a synergistic blend of superquats and glutaraldehyde (Duplalm[®], Veterquímica S.A., Chile), which is commercialized as a disinfectant of broad spectrum, presenting fast action and low toxicity. For these reasons, the aims of this study were to determine the efficacy of Duplalm[®] in treating various fish pathogens under *in vitro* conditions and large scale and validate the treatment regimes for controlling these diseases in salmon production.

MATERIAL AND METHODS

This study was done in two experimental stages: I) *In vitro* susceptibility of the extracellular bacteria *V. ordalii* and *V. anguillarum*, intracellular pathogens as *Francisella* sp. and IPNV. II) Large scale assays conducted at the hatchery of the Laboratory of Research and Development at Veterquímica. The experiments 1 and 2 of the stage I was carried out in triplicate for each bacterial strain, while the experiment 3 considered the use of IPNV.

STAGE I: *IN VITRO* ASSAYS IN LABORATORY

EXPERIMENT 1

The effectiveness of different Duplalm[®] treatments against *V. ordalii* and *V. anguillarum* was assessed following the procedures of Avendaño-Herrera *et al* (2006). The *V. ordalii* strains used in this work, Au2, Au3 and PF180, were isolated in Chile during 2003-2005 from clinically infected Atlantic salmon (Silva-Rubio *et al* 2008^a). In the case of *V. anguillarum*, three strains isolated from Atlantic salmon coded as PF2, PF7 and PF8 and belonging to serotype O3 of this fish pathogen (Silva-Rubio *et al* 2008^b) were employed. The reference strains of *Vibrio ordalii* ATCC 33509^T and *Vibrio anguillarum* ATCC 43307 from the American Type Culture Collection were used for comparative purposes. The experimental assays were conducted in sterile 6-well Multidishes (Nunc[™] Surface) containing 10 ml of natural sterilized (filtered and autoclaved) seawater. Inocula were prepared from bacteria grown on tryptone soy agar (Oxoid) supplemented with 1% NaCl (Winkler) plates (TSA-1), washed in 0.85% sterile saline solution (SS), resuspended and diluted in SS, and then each well was seeded with 500 µl of the bacterial suspension to achieve an initial *V. ordalii* and *V. anguillarum* concentration of 5×10^6 cells ml⁻¹ as determined by direct microscopy count (equivalent to 1.8×10^6 CFU x ml⁻¹, respectively). After a stabilization period of 30 min. at 19 ± 1 °C, susceptibility of each fish pathogens and bacterial strain to different Duplalm[®] concentrations was examined in triplicate, adding this chemical compound to 1:200; 1:300 and 1:400 dilutions. Duplalm[®] was evaluated during 30, 60, 120 and 180 s at 20 °C. Controls without the addition of the disinfectant

were run simultaneously in exactly the same conditions as described above.

To determine the cultivable cells in the seawater, samples of 0.5 ml were taken aseptically, after each treatment. All samples were serial-diluted in SS and 0.1 ml of each dilution was plated in duplicate on TSA-1 plates for the count of total heterotrophic marine bacteria. After incubation, biochemical identification tests (API20E; Biomérieux) and slide agglutination tests (Silva-Rubio *et al* 2008^{a, b}) were carried out to determine the presence of the inoculated *V. ordalii* and *V. anguillarum* strains in each Multidish.

EXPERIMENT 2

The *Francisella* sp. strain used in this work was isolated in Chile in 2006 from spleen of clinically infected Atlantic salmon. To determine the effect of different Duplalm[®] concentrations on *Francisella* sp, bactericidal assays were conducted according to a previously described protocol of Avendaño-Herrera *et al* (2006) with minor modification.

Pure colonies of *Francisella* sp. cultivated at 20 °C were picked up from cysteine heart agar (Difco[™]) plates containing 5% sheep blood (Olsen *et al* 2006), adjusted in sterile freshwater to contain 3×10^8 CFU x ml⁻¹. Duplalm[®] was evaluated during 2, 5 and 15 min treatment periods for the control of *Francisella* sp. using diluted disinfectant to a 1:50; 1:200 and 1:400 concentrations. *Francisella* sp. controls were run simultaneously without the addition of the disinfectant. The experiment was also carried out in triplicate.

Treatments were considered effective if a pure culture of grey white, slightly mucoid convex colonies of *Francisella* sp. was absent when the samples which were taken aseptically at the end of the incubation period were spread onto media.

EXPERIMENT 3

Infectious pancreatic necrosis virus (IPNV) strain 483, of the serotype Sp, originated from an outbreak in *Salmo salar* cultured in Chile in 2005, was used for the assay. The virulent IPNV isolate was previously identified from tissue according to Espinoza and Kuznar (2002). The virus was subjected to two rounds of plaque purification on Chinook salmon embryo cells (CHSE-214) before propagation in the same fish cell line to obtain a stock (Johansen and Sommer 2001). The infected cell cultures were kept frozen at -80 °C until they were used. A small aliquot of the material was frozen separately and was used for the determination of TCID₅₀ (50% tissue culture infective dose) by end-point dilution (Reed and Muench 1938). The experimental assays were conducted in sterile 6-well Multidishes (Nunc[™] Surface) containing 10 ml of the CHSE-214 with IPNV at titres of $10^{5.6}$; $10^{6.6}$ and $10^{7.6}$ TCID₅₀ ml⁻¹. Susceptibility of IPNV to Duplalm[®] was examined in triplicate, with

the addition of this chemical compound at final dilution of 1:400, and then incubated for 2, 5 and 15 min at $19 \pm 1^\circ\text{C}$. The action of the disinfectant was stopped using a specific neutralizer (polysorbate 80 (3%), lecithin 3 g/L, sodium thiosulphate 5 g/L, L-histidine 1 g/L and saponin 30 g/L), which was validated for all disinfectants before use in the tests. Negative controls, consisted of a harvested cell culture medium from uninfected cell cultures exposed to the same Duplalin[®] concentration but not inoculated with IPNV. The infected cell cultures but not treated with the disinfectant were included as positive controls. All controls were run simultaneously in the exact same conditions as described above.

To test the efficacy of superquats and glutaraldehyde compounds to inactivate IPNV, 0.5 ml aliquots of the virus solution were taken aseptically and transferred to plates containing the CHSE-214 cells. Each sample was passaged three times every seven days and incubated at 18°C . The cytopathic effect was observed by inverted microscopy for up to 7, 14 and 21 days post-infection.

Moreover, attempt was made to check if the inhibition caused by Duplalin[®] was produced only by the disinfectant or contributed to the persisted active residues in the media. The IPNV isolate 483 was cultured as previously described and concentrated by successive ultrafiltrated steps until $10^{1.5}$ TCID₅₀/ml was obtained. Two aliquots of 5 ml each was taken, one of them being used as control (no disinfectant added) and the other aliquot treated with Duplalin[®] at final dilution of 1:400, followed by incubation for 2 min at 18°C . Each solution was transferred to a Centricon ultrafiltration device with 10-kDa molecular mass cutoff (Millipore), and supernatants were removed by centrifugation ($3500 \times g$ for 60 min at 4°C). IPNV were resuspended in phosphate buffered saline (pH 7.4) and dialysis was repeated until a concentration of residual Duplalin[®] reached 1:10000 dilutions. The control sample (without disinfectant) was treated in the same way. Finally, 100 µl of the treatment and control were left in a 24-well plate containing CHSE-214 cells and mixed. Serial ten-fold dilutions (from 10^{-1} to 10^{-5}) were made in quadruplicate and then incubated at 18°C for two weeks. Each plate was observed by inverted microscopy for the appearance of cytopathic effect for 7 and 14 days.

STAGE II: LARGE SCALE ASSAY

To mimic a “near real-life” the assays for the evaluation of Duplalin[®] was performed in buckets containing 10 L of $0.45 \mu\text{m}$ filtered seawater. Four buckets were inoculated with the strains of *V. anguillarum* Au2 and *V. ordalii* PF2 at a final concentration of 1.8×10^6 CFU \times ml⁻¹ of a 1:1 mix concentration. Each bucket was seeded with the bacterial suspension prepared as described for experiment 1 and homogenized with a fish net for 60 s. Each fish net was maintained inside the buckets. Following a 30 min acclimatization period, addition of Duplalin[®] at

a concentration of 1:400 was performed to two buckets, while the remaining inoculated buckets without the disinfectant were considered as positive controls. The negative control corresponded to a new bucket with only seawater, no bacteria nor chemical disinfectant added.

In order to evaluate the effect of Duplalin[®] on the microorganisms attached to the surface of the fishing net (net used in salmon cages) at the end of incubation (24 h), the thread mesh of each experimental set was sampled. For this, ten pieces of net were carefully cut into 1 cm^2 pieces, washed repeatedly with SS and put into Falcon tubes containing 10 ml of SS. The bacteria that were attached to the pieces of mesh were removed using an Ultrasonics Homogenizer for 15 seconds.

Bacteriological sampling of water was carried out at 15 min, 4 h and 24 h post-treated with Duplalin[®]. Samples were serially diluted in SS and 0.1 ml of each dilution was plated on TSA-1 for the total count of *Vibrionaceae*. To increase the sensitivity of the technique, samples of 1 ml were taken directly from the water of each bucket and seeded onto the same bacteriological medium. The plates were incubated at 20°C for one week. After incubation, selective counts of CFU (according to the morphotypes grown on TSA-1 plates) and confirmation by serological identification tests (i.e. slide agglutination assays) according to Silva-Rubio *et al* (2008^{a, b}) were carried out to determine the presence of each bacterium inoculated in seawater.

RESULTS AND DISCUSSION

The addition of Duplalin[®] at concentrations among 1:200 to 1:400 to seawater killed completely *V. ordalii* and *V. anguillarum* in all *in vitro* treatments, regardless of the period tested and strains used. In contrast, there was no loss of viability in the un-disinfectant controls with values of $1.6 \pm 0.56 \times 10^6$ CFU \times ml⁻¹ for *V. anguillarum*, and $1.4 \pm 0.97 \times 10^6$ CFU \times ml⁻¹ for *V. ordalii* (similar to initial inoculate of each pathogen).

The addition of Duplalin[®] to freshwater killed completely *Francisella* sp. at a concentration of 1:50 within 5 and 15 min. (table 1). When the concentration of the disinfectant was decreased, a higher degree of resistance to Duplalin[®] was observed. Interestingly, in freshwater with a dilution of 1:400, *Francisella* sp. cells were recovered regardless the incubation time. Similar cultivability pattern to *Francisella* sp. cells was observed in the negative control groups (without disinfectant). In spite of the limited number of the isolates tested (only one) it seems that *Francisella* sp., an intracellular facultative microorganism, had the highest resistance to Duplalin[®] than the extracellular pathogens such as *V. ordalii* and *V. anguillarum*. In a recent study, Verner-Jeffreys *et al* (2009) reported that two Gram-positive bacteria (*Lactococcus garviae* and *Carnobacterium piscicola*) were apparently more resistant to four biocide products than other two

Table 1. *In vitro* bactericidal effect of Duplalin® on *Francisella* sp. at 20 °C.Efecto bactericida *in vitro* de Duplalin® en *Francisella* sp. a 20 °C.

Duplalin® dilutions	Exposure time (min)		
	2	5	15
Untreated-control	+++ ^a	+++	+++
1:50	+	–	–
1:200	+++	++	–
1:400	+++	+++	++

^a Data correspond to the results of the *Francisella* sp. isolate in triplicate.

– Effective (no CFU after exposure).

Growth of *Francisella* sp.: +, < 10²; ++, 10³ and +++, > 10⁵.

Gram-negative pathogens. In general, in the case of the Gram-positive bacteria it is suggested that the presence of some cell structures (i.e. capsular material) may confer resistance to antibacterial compounds (i.e. activity of serum). Considering that *Francisella* sp. replicates and grows within the membrane-bound cytoplasmatic vacuoles

of infected host cells (Olsen *et al* 2006) the resistance shown by the isolate to Duplalin® might be explained by the low concentration that can be attained by this disinfectant compound. Until now, the molecular base of this resistance is unknown. Further studies are needed to elucidate the composition and chemical structure and/or the role of some enzyme produced by *Francisella* in its protection.

The survival capacity of the IPNV isolate 483 in the CHSE-214 cells after the Duplalin® treatments is shown in table 2. The concentration of the disinfectant had a significant effect with 2, 5 and 15 min. of exposure, regardless of the IPNV titres employed (until a concentration of 10¹⁵ TCID₅₀ ml⁻¹). As expected, the results of cytopathic effect test showed that IPNV produced degenerative changes in the CHSE cell line without disinfectant passed three times. On the other hand, it is important to denote that the action of active residues was efficiently stopped with the addition of the neutralizer as well as by the dialysis procedures. Therefore, no toxicity of the disinfectant and solvent either detected over the cells. In fact, after incubation (IPNV with and/or without Duplalin®) only total cytopathic effect was detected in the CHSE-214 plates without the disinfectant, therefore, IPNV was re-isolated at 10⁻⁵ serial dilutions during two weeks.

Table 2. Effect of Duplalin® at final dilution of 1:400 on *in vitro* inactivation of Infectious Pancreatic Necrosis Virus.Efecto de Duplalin® en dilución final de 1:400 sobre la inactivación *in vitro* del Virus de la Necrosis Pancreática.

Titre viral (TCID ₅₀ / ml)	Exposure time (min)	Passage 1 ^a	Passage 2	Passage 3
10 ^{5.6}	2	–	–	–
	5	–	–	–
	15	–	–	–
	Untreated-control	+	+	+
10 ^{6.6}	2	–	–	–
	5	–	–	–
	15	–	–	–
	Untreated-control	+	+	+
10 ^{7.6}	2	–	–	–
	5	–	–	–
	15	–	–	–
	Untreated-control	+	+	+
10 ¹⁵	2	–	–	–
	Untreated-control	+	+	+

^a Three replicates per passage.

– Effective (no cytopathic effect after exposure).

+ Positive Sample for IPNV with an extensive cytopathic effect of approximately 90% was observed at 7 days.

In general, most of the disinfection studies in IPNV have been carried out using other chemical disinfectants such as iodine, chlorine and formalin. Both iodine and chlorine inactivated IPNV, although the efficacy depended on the pH, virus concentration and the presence of organic matter in water (Elliott and Amend 1978). Studies of Desautles and MacKelvie (1975) reported that a 30 min exposure to chlorine at a concentration of 40 ppm effectively inactivates a concentration of $10^{7.5}$ TCID₅₀/ml, while using 35 ppm iodine for the same time is appropriate to eradicate a concentration of $10^{6.6}$ TCID₅₀/ml. These authors also denoted that although formalin exhibited some *virucidal* activity, it was not a good choice for the inactivation of IPNV (MacKelvie and Desautles 1975).

With the aim to assess if Duplalin[®] can be a suitable candidate for the treatment of seawater and surface of tanks, nets and/or rearing equipment, bioassays were performed in buckets with seawater containing a mesh to stop fish and seeded with two different fish pathogens. In this study, members of the genus *Vibrio* were selected because they constitute part of the microbiota of marine fish as well as part of the normal microbiota of the aquatic environment, and therefore present a constant threat for any susceptible host (Austin and Austin 2007). The culture of *V. ordalii* and *V. anguillarum* in the buckets treated with 1:400 disinfectant concentrations had a significant effect on the survival of the microbiota in the seawater, with no viable bacteria being detectable from 15 min post-treatment. In contrast, controls without the addition of Duplalin[®] maintained $1.37 \pm 0.29 \times 10^6$ CFU/ml during the experiment. Similarly, there was no evidence of *Vibrionaceae* culture adhered on the fishing net compared to untreated Duplalin[®] tries. It is important to note that survival for an aquatic bacterium in seawater outside the fish host may depend on biofilm formation. *V. anguillarum* is normally found attached to surfaces, as opposed to a free-swimming form, which is likely to provide an adaptive or survival advantage for bacteria in the aquatic environment (Costerton *et al* 1987, 1995). In each of these environments, the bacterium likely utilizes its external membrane to protect its intracellular contents from damaging agents or conditions (Wang *et al* 2003). When attempt of resuscitation of the non-culturable bacteria with the addition of fresh medium was employed, no cells from the mesh samples were recovered onto TSA-1 (data not shown), indicating that all microorganisms were completely killed.

On the other hand, cultivable bacteria remained constant in the positive controls (without the disinfectant), regardless of the period sampled, and showing an average value of $1.4 \pm 0.3 \times 10^6$ CFU/ml. The bacteria adherence on the mesh without exposure to the disinfectant was about 55.3% ($7.6 \pm 3.2 \times 10^5$ CFU/ml) lower in concentration than those containing in the seawater after 24 h. Selective counts of *V. anguillarum* on TSA-1 sampled from the buckets with no treatment represented an average of 49.7% of the total microbiota counted. As expected, seawater without

the mixed bacteria and disinfectant remained with a low load of bacteria (6.3×10^5 CFU \times ml⁻¹) through the entire period of the study.

Although Duplalin[®] is one of the most used disinfectants in the Chilean salmon industry, it has been legally registered and recently validated to be used against Infectious Salmon Anaemia Virus (ISA) virus and its effective use may be well established in this study; we cannot discard that organic matter present in the water column or dirty conditions in the water could interfere on the efficiency of the disinfectant. Studies are being carried out to test this condition using the Verner-Jeffreys *et al* (2009) protocol.

It can be concluded that, apart from the fact that in this work we used a methodology that is different to the test standards used for the evaluation of bactericidal and virucidal activity of disinfectants and antiseptics for use in the veterinary field published recently by Verner-Jeffreys *et al* (2009), our results demonstrate that the use of Duplalin[®] in aquaculture is effective in low concentration and short time of exposure (15 min. at a concentration of 1:400 dilutions), with the exception of the intracellular pathogen used in this work. It is important to highlight that Verner-Jeffreys *et al* (2009) tested disinfectants of different nature than Duplalin[®] (i.e. hydrogen peroxide, peracetic acid, acidic iodophore and Chloramine T) and suggested that when studying other disinfectants (i.e. superquats and glutaraldehyde), the procedure can be adopted or modified to be used in countries outside Europe. Therefore, the use of Duplalin[®] may be appropriate as a general preventive disinfection method against *V. ordalii*, *V. anguillarum* and IPNV for treating water culture, surface of tanks, nets and/or rearing equipment, but all these equipments would need to rinse off before they could be used in contact with fish.

SUMMARY

The efficacy of the disinfectant Duplalin[®], a synergistic blend of superquats and glutaraldehyde, was analysed *in vitro* against 4 fish pathogens. All concentrations tested (1:200 to 1:400 dilutions) were efficacious on killing *Vibrio ordalii* and *Vibrio anguillarum* in seawater after 30 s, being the level of reduction equal to 1.8×10^6 CFU/ml. Higher concentration of Duplalin[®] (1:50 dilutions) and time of exposure (at least 5 min) is needed to kill completely *Francisella* sp, an intracellular freshwater pathogen. When Infectious Pancreatic Necrosis Virus (IPNV) was treated with 1:400 disinfectant dilution, this concentration had a significant effect after 2 minutes, regardless of the IPNV titres employed (concentration greater than $10^{7.6}$ TCID₅₀/ml). Duplalin[®] was tested in large scale against *Vibrionaceae* members. In comparison to the controls (without the disinfectant), 1:400 dilutions of Duplalin[®] totally killed *V. ordalii* and *V. anguillarum* in seawater as well as on the surface of the fishing net (used in the cages of cultured salmon) after 15 min. Cultivable bacteria remained constant in the buckets without the disinfectant ($1.4 \pm 0.3 \times 10^6$ CFU/ml), regardless of the period sampled. In the case of the adherence on the fishing net, bacteria not exposed to the disinfectant were detected at a concentration of $7.6 \pm 3.2 \times 10^5$ CFU/ml. These data indicate that the use of Duplalin[®] against *V. ordalii*, *V. anguillarum* and IPNV is effective in low concentration and short time of exposure (15 min at a concentration of 1:400 dilutions), while the intracellular pathogen requires higher concentration.

ACKNOWLEDGEMENTS

This work was supported in part by Grant IPC 019 from the Program Bicentenario Ciencia y Tecnología, CONICYT-Chile and also by Grant DI-01-10/R from the Universidad Andrés Bello.

REFERENCES

- Alderman DJ, TS Hastings. 1998. Antibiotic use in aquaculture: development of antibiotic resistance-potential consumer health risks. *Int J Food Sci Technol* 33, 139-155.
- Austin B, A Austin. 2007. *Bacterial fish pathogens: Diseases of farmed and wild fish*. Praxis Publishing Ltd, Chichester, UK.
- Avendaño-Herrera R, B Magariños, R Irgang, AE Toranzo. 2006. Use of hydrogen peroxide against the fish pathogen *Tenacibaculum maritimum* and its effect on infected turbot (*Scophthalmus maximus*). *Aquaculture* 257, 104-110.
- Avendaño-Herrera R, A Silva-Rubio, AE Toranzo. 2007. First report of *Vibrio anguillarum* in Chilean salmon farms. *AFS/ Fish Health Newsletter* 35, 8-10.
- Birkbeck TH, M Bordevik, MK Frøystad, M.K, A Baklien. 2007. Identification of *Francisella* sp. from Atlantic salmon, *Salmo salar* L., in Chile. *J Fish Dis* 30, 505-507.
- Burka JF, KL Hammell, TE Horsberg, GR Johnson, DJ Rainnie, DJ Speare. 1997. Drugs in salmonid aquaculture - a review. *J Vet Pharmacol Therap* 20, 333-349.
- Colquhoun DJ, IL Aase, C Wallace, Á Baklien, K Gravningen. 2004. First description of *Vibrio ordalii* from Chile. *Bull Eur Ass Fish Pathol* 24, 185-188.
- Costerton JW, K-J Cheng, GG Geesey, TI Ladd, JC Nickel, M Dasgupta, TJ Marrie. 1987. Bacterial biofilms in nature and disease. *Annu Rev Microbiol* 41, 435-464.
- Costerton JW, Z Lewandowski, DE Caldwell, DR Korber, HM Lappin-Scott. 1995. Microbial biofilms. *Annu Rev Microbiol* 49, 711-745.
- Elliott DG, DF Amend. 1978. Efficiency of certain disinfectants against infectious pancreatic necrosis virus. *J Fish Biol* 12, 277-286.
- Espinoza JC, J Kuznar. 2002. Rapid simultaneous detection and quantitation of infectious pancreatic necrosis virus (IPNV). *J Virol Meth* 105, 81-85.
- Fryer JL, RP Hedrick. 2003. *Piscirickettsia salmonis*: a Gram-negative intracellular bacterial pathogen of fish. *J Fish Dis* 26, 251-262.
- Gibello A, A Mata, M Blanco, A Casamayor, L Domínguez, JF Fernández-Garayzábal. 2005. First identification of *Streptococcus phocae* isolated from Atlantic salmon (*Salmo salar*). *J Clin Microbiol* 43, 526-527.
- Johansen LH, AI Sommer. 2001. Infectious pancreatic necrosis virus infection in Atlantic salmon *Salmo salar* postsmolts affects the outcome of secondary infections with infectious salmon anaemia virus or *Vibrio salmonicida*. *Dis Aquat Org* 47, 109-117.
- Mackelvie RM, D Desautels. 1975. Fish viruses-survival and inactivation of infectious pancreatic necrosis virus. *J Fish Res Board Can* 32, 1269-1273.
- Olsen AB, J Mikalsen, M Rode, A Alfjorden, E Hoel, K Straum-Lie, R Haldorsen, DJ Colquhoun. 2006. A novel systemic granulomatous inflammatory disease in farmed Atlantic cod, *Gadus morhua* L., associated with a bacterium belonging to the genus *Francisella*. *J Fish Dis* 29, 307-311.
- Reed LJ, H Muench. 1938. A simple method for estimating fifty percent endpoints. *Amer J Hyg* 27, 493-497.
- Romalde J, C Ravelo, I Valdés, B Magariños, E de la Fuente, C San Martín, R Avendaño-Herrera, AE Toranzo. 2008. Characterization of *Streptococcus phocae*, an emerging pathogen for salmonid culture. *Vet Microbiol* 130, 198-207.
- Silva-Rubio A, C Acevedo, B Magariños, B Jaureguiberry, AE Toranzo, R Avendaño-Herrera. 2008^a. Antigenic and molecular characterization of *Vibrio ordalii* strains isolated from Atlantic salmon (*Salmo salar*) in Chile. *Dis Aquat Org* 79, 27-35.
- Silva-Rubio A, R Avendaño-Herrera, B Jaureguiberry, AE Toranzo, B Magariños. 2008^b. First description of serotype O3 in *Vibrio anguillarum* strains isolated from salmonids in Chile. *J Fish Dis* 31, 235-239.
- Smith P, MP Hiney, OB Samuelsen. 1994. Bacterial resistance to antimicrobial agents used in fish farming: a critical evaluation of method and meaning. *Annu Rev Fish Dis* 4, 273-313.
- Verner-Jeffreys DW, CL Joiner, NJ Bagwell, RA Reese, A Husby, PF Dixon. 2009. development of bactericidal and virucidal testing standards for aquaculture disinfectants. *Aquaculture* 286, 190-197.
- Wang S-Y, J Lauritz, Jana, J Jass, DL Milton. 2003. Role for the major outer-membrane protein from *Vibrio anguillarum* in bile resistance and biofilm formation. *Microbiology* 149, 1061-1071.