



Latin American Journal of Aquatic
Research

E-ISSN: 0718-560X

lajar@ucv.cl

Pontificia Universidad Católica de
Valparaíso
Chile

De los Santos - Romero, Rodolfo B.; García-Guerrero, Marcelo; Vega-Villasante,
Fernando; Nolasco-Soria, Héctor

Effect of dietary chitin on digestive enzyme activity, growth and survival of
Macrobrachium tenellum juvenile prawns

Latin American Journal of Aquatic Research, vol. 45, núm. 1, marzo, 2017, pp. 130-138
Pontificia Universidad Católica de Valparaíso
Valparaíso, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=175050001012>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Research Article

Effect of dietary chitin on digestive enzyme activity, growth and survival of *Macrobrachium tenellum* juvenile prawns

**Rodolfo B. De los Santos-Romero^{1,5}, Marcelo García-Guerrero²
Fernando Vega-Villasante³ & Héctor Nolasco-Soria⁴**

¹Instituto Tecnológico del Valle de Oaxaca, Santa Cruz Xoxocotlan, Oaxaca, México

²CIIDIR-Oaxaca, Instituto Politécnico Nacional, Santa Cruz Xoxocotlan, Oaxaca, México

³Centro de Investigaciones Costeras, Universidad de Guadalajara
Puerto Vallarta, Jalisco, México

⁴Centro de Investigaciones Biológicas del Noroeste, La Paz, Baja California Sur, México

⁵Doctoral Student, CIIDIR-Oaxaca, Instituto Politécnico Nacional
Santa Cruz Xoxocotlan, Oaxaca, México

Corresponding author: Héctor Nolasco-Soria (hnolasco04@cibnor.mx)

ABSTRACT. *Macrobrachium tenellum* is a freshwater prawn native from Mexico to Peru, with potential for cultivation. Currently, it is extensively exploited by local fisheries. To understand its chitin requirements, a formulated diet with chitin added (5, 10 15, 20, and 25%) was given to juveniles for 60 days. Growth, survival, and enzyme activity (trypsin, chymotrypsin, lipase, amylase, and chitinase) were measured every 15 days. Highest final weight and specific growth rate resulted from diets with 20% chitin. Poorest performance resulted from diets with 5 and 10% chitin. Chitin in the diet did not have a significant effect on survival. Statistically significant differences in enzyme activity were not found in any treatment.

Keywords: chitin, diet, nutrition, *Macrobrachium tenellum*, aquaculture.

INTRODUCTION

Within the American species of the genus *Macrobrachium* (Bate, 1868) only few are economically important. Some species have potential for cultivation: *M. carcinus* (Linnaeus, 1758), *M. americanum* (Bate, 1868), *M. digueti* (Bouvier, 1895), *M. acanthurus* (Wiegmann, 1836), *M. amazonicus* (Heller, 1962), and *M. tenellum* (Smith, 1871). *M. tenellum*, the long arm river prawn, has been considered a candidate for cultivation because it grows quickly, tolerates high densities and wide fluctuations in water conditions, and adapts well to captivity (Ponce-Palafox *et al.*, 2002; Espinosa-Chaurand *et al.*, 2011; García-Guerrero *et al.*, 2013). This species is still common in its natural environment, which is necessary for recruitment.

Currently, there are few works focused on *M. tenellum* cultivation, particularly feeding and nutrition. Previous studies include García-Ulloa *et al.* (2004) who determined the effects of protein diets on growth, replacing fish meal with soybean meal and Espinosa-Chaurand *et al.* (2012) who determined the effect of

levels of protein in diets on growth and survival of juveniles. Additional studies of feeding and nutrition are required to establish the type and balance of feed ingredients to produce good results under cultivation. Espinosa-Chaurand (2013) describes enzyme activity in this species. However, enzymes that act on chitin are not well studied yet. This information is required because *Macrobrachium* prawns consume crustacean exoskeletons, which provide ingredients for new exoskeletons after molting (Kumar *et al.*, 2006; Zhang *et al.*, 2014). Chitin digestive enzyme in this prawn has not been studied (García-Guerrero *et al.*, 2013).

Most of the works on the effect of chitin in the diet of crustaceans have been developed with penaeid species (Chang *et al.*, 1988; Shiao & Yu, 1998; Terwilliger, 1999; Chang *et al.*, 2001; Casillas-Hernández *et al.*, 2006; Chih-Hui & Chen-Chun, 2012). For *Macrobrachium* prawns, Kumár *et al.* (2006) compared the effect of natural chitin and purified chitin on the growth of *Macrobrachium rosenbergii* (De Man, 1879).

Chitin in diets of prawns advance growth (Kumar *et al.*, 2006), promotes antimicrobial activity in diets (Somjit *et al.*, 2005), and supports molting (Zhang *et al.*, 2014). In our study, we measured the effect of different levels of chitin in the diet for *Macrobrachium tenellum* juveniles on growth and digestive enzyme activity, with special emphasis on chitinases.

MATERIALS AND METHODS

Experimental design

A batch of *Macrobrachium tenellum* juveniles were collected in the Chacahua Lagoon (15°58'20"N, 97°32'28"W) in Oaxaca, Mexico. They were transported in a 250 L tank with permanent aeration to laboratory facilities and then, acclimated during two weeks at 25°C. During this time, they were fed with marine shrimp commercial pelletized food (40% crude protein, Camaronina 40® Agribands Purina, Mexico).

In order to evaluate chitin effect on growth, a completely randomized experimental design of five treatments was realized. Every treatment had a different amount of chitin, included in a isoproteic-isolipidic diet. Different chitin amounts were 5, 10, 15, 20 and 25% of the food. An experimental control (no chitin) was also included. The experimental diet was prepared in agreement with Cortes-Jacinto *et al.* (2003) and Espinosa-Charaund *et al.* (2012) with the following ingredients: *Cyprinodontid* fish meal (50.05%), soybean meal in bulk (25.03%), Maseca® commercial corn flour (17.80%), and San Antonio® commercial wheat flour (7.12%) and vitamin and mineral supplements (Fervinac with selenium®, Laboratorios Brovel, Mexico). All ingredients were blended and mixed with a blender. Chitin was prepared by pulverizing dried shrimp carcasses (*Litopenaeus* sp.). The carcasses were cleaned by rinse and then, dried at sunlight. When completely dry, they were pulverized with the blender and added to the mixture. Then, the mixture was sieved through a 500 µm mesh and mixed again with the same blender. At this step, cornstarch were added as agglutinant. The resultant mixture was passed through a meat grinder in order to obtain 2 mm diameter pellets. Those pellets were dried in a convection oven at 30°C during 12 h. Table 1 shows the proximate analysis of the diet, to establish the values for gross energy, the conversion factors suggested by Anh *et al.* (2009) were applied: 4.2 for carbohydrates; 5.56 for proteins; and 9.54 for lipids.

Every treatment had six replicates and every replicate consisted on a dark non-translucid plastic container (50 L). All replicas of the same treatment were placed together at different levels. In the container

Table 1. Proximate analysis of tested diet. Analytical method: ¹Differences in moist and dry weight, ²AOAC, 2012 (micro-Kjendahl method), ³AOAC, 2012 (Soxhlet method), ⁴AOAC, 2012 (Weende method), ⁵AOAC, 2012 (oven at 550°C).

	g 100 g ⁻¹ of dry food Average value ± SD
Humidity (%) ¹	4.783 ± 0.37
Crude protein ²	44.137 ± 0.19
Total lipids ³	6.517 ± 0.20
Crude fiber ⁴	1.557 ± 0.09
Ash ⁵	9.700 ± 0.38
Nitrogen-free extract	38.090 ± 0.77
Gross energy (kcal g ⁻¹)	4.741

at the lowest level no specimens were placed. A water pump was placed with a hose that throw water to the highest tray. Every tray had an outcome flowing to its lower next. This way, a cascade is produced and water is constantly recirculating through all containers of the same treatment. In every treatment, water temperature was maintained always at 28°C, with submersible 300 W heaters (RENA Aquatic Supply; US) installed in the lowest non-specimen tray of every treatment. Oxygen concentration was daily measured with a Hanna® HI98186 oximeter (Hanna Instrumenta Inc; Italy) and maintained always above 5 mg O₂ L⁻¹. The photoperiod was 12 h light -12 h dark. In each container, a 0.5 m² of plastic mesh was placed inside the water to provide shade and shelter. Ten organisms (0.2 ± 0.04 g) were placed in every container. A week before the first day of the experiment, total daily food was provided at 7% of the prawn biomass. After one week, the ration was adjusted to 10% of their body weight for all treatments. They were fed every day at 18:00 h. Survival was registered daily. The trial lasted 60 days. At days 0, 15, 30, 45, and 60, all prawns were individually weighed (±0.001 g).

The following parameters were obtained: average individual weight gain (g): AIWG = final weight-initial weight; gain weight per day (g day⁻¹): GWPD = (final weight-initial weight) t⁻¹; gain weight in percentage (%): GW = 100 × (final weight - initial weight)/initial weight; and specific growth rate (%): SGR = [(ln final weight - ln initial weight) t⁻¹] × 100. Also the feed conversion ratio was calculated as: FCR = supplied food (g)/weight gain (g); and feed efficiency ratio: FER = weight gain (g)/supplied food (g). SGR and FCR were calculated in agreement with Cortes-Jacinto *et al.* (2003); Gitte & Indulkar (2005) were followed for GW and GWP; Vega-Villasante *et al.* (2011) for AIWG and SR; and Hasan *et al.* (2012) for FER.

Enzyme activity

For measuring enzymatic activity, two prawns from each replica (eight by treatment) were randomly selected on days 0, 30 and 60. These were weighed as fresh weight and kept at -20°C until analysis. Only the hepatopancreas and intestine were analyzed. By the size of juvenile prawns the anterior section of head, all appendages, and the exoskeleton were discarded during dissection. The remaining tissue after dissection were individually weighed and separately homogenized with cold distilled water (4°C) in a v/w proportion of four mL water for g fresh organ tissue. Raw extracts were separated by centrifugation at 14,000 rpm for 10 min at 4°C , and clarified crude extract was kept at -40°C until analyzed for soluble proteins and enzyme activity. All measurements were made in quadruplicate for all enzymatic analysis, a control sample (blank) was also included, where the enzyme reagent was added after the reaction ceased. Protein concentrations in the enzyme raw extracts were quantified by the Bradford method (1976). In glass tubes (100×15 mm), 8 μL of crude extract, 792 μL distilled water, and 200 μL Bradford reagent were mixed and gently vortexed. Absorbance was measured at 595 nm. Bovine serum albumin (05470, Sigma-Aldrich, St. Louis, MO) was used as the protein standard. Lipase activity was determined as described by Versaw *et al.* (1989), using β -naphthyl caprylate as the substrate. Lipase activity was expressed as lipase units mg^{-1} protein (one lipase unit was the quantity of enzymes required for an increase of 0.01 absorbance units at 540 nm min^{-1}). Amylase activity was measured, as described by Vega-Villasante *et al.* (1993), using 1% starch in 50 mM Tris-HCl at pH 7.5, as the substrate. This activity was expressed as amylase units/mg protein (one amylase unit was defined as the quantity of enzyme to increase absorbance units by 0.01 at 550 nm min^{-1}). Trypsin activity was determined, using BAPNA as the substrate (García-Carreño & Haard, 1993), adapted to a 96-well microplate by adding 10 μL crude extract, 160 μL 60 mM Tris-HCl at pH 8.0, 10 μL 192 mM CaCl_2 at pH 8.0, and 10 μL 9.6 mM BAPNA dissolved in DMSO, in each well to start the reaction. Chymotrypsin activity was determined, using 9.6 mM SAAPNA dissolved in DMSO. For both enzymes, absorbance at 414 nm was recorded every 15 s for 30 min. At the end of the assay, a linear coefficient was calculated to determine the increase of absorbance per second. Enzyme activity was calculated using molar extinction coefficient of p-nitroaniline (8800). Chitinase activity was determined by mixing 20 μL of crude extract, 50 μL 60 mM Tris-HCl at pH 8, and 530 μL substrate (C3020, Sigma-Aldrich). The reagent mixture was shaken with a vortex at 120 rpm (at 45° angle) for incubation during 2 h. The

reaction was stopped by centrifugation after 5 min at 14,000 rpm for 10 min at 4°C . The supernatant was separated immediately and absorbance was measured at 570 nm (one chitinase unit was defined as the quantity of enzymes required for an increase of 0.001 absorbance units at 570 nm min^{-1}).

Statistical analysis

Survival rate (%), average individual weight gain (g), gain weight per day (g day^{-1}), specific growth rate (%), feed conversion ratio, feed efficiency ratio and enzyme activity (U mg^{-1} Prot Sol) were determined and then compared by one way ANOVA test. Normality tests were performed (Kolmogorov-Smirnov test, $\alpha = 0.05$). Significant differences between treatment were determined by the Duncan multiple range test ($P > 0.05$). Minitab 17 statistical package (Minitab, College Park, PA) was utilized for statistical analyses.

RESULTS

During the 60 day trials, the water in all experimental units were maintained at 28.3°C , pH 7.5, and 6.8 $\text{mg O}_2 \text{ L}^{-1}$. After 60 days, only the growth parameter of absolute weight was statistically significant (Fig. 1).

The range of each growth parameter was: GWPD (0.0048 to $0.0085 \text{ g day}^{-1}$); GW (147.1 to 286.5%); SGR (1.42 to 1.88%); FCR (3.86 to 2.91); and FER (2.72 to 3.90). Additionally, Table 2 shows growth (weight) per treatment.

The inclusion of 20% in one diet produced the best average individual weight gain ($0.508 \pm 0.101 \text{ g}$) and the best gain weight per day ($0.0085 \pm 0.0017 \text{ g day}^{-1}$). Treatment with 15% and 25% chitin were less successful. The best growth rate occurred by day 45, afterward declining. The least growth occurred in the 10% and 5% chitin treatment and in the control treatment (Fig. 2). There was no statistically significant difference in survival between treatments, ranging from 72.2% (treatment 5) to 83.3% (treatments 1, 2, 3, and control). For FCR, the best treatments were the 20% chitin (2.91 ± 0.39) and 25% chitin (3.50 ± 0.55), with the highest feed conversions around day 45.

For soluble protein concentrations, there were no statistically significant differences between treatments. Chitinase activity was the only enzymatic activity with statistically significant differences between treatments. Other enzymes had similar trends in the final phase of the culture (Fig. 3). Like specific growth data (see Fig. 2), chitinase activity was highest at day 45, declining thereafter to the end of the trials (Fig. 4), differing from lipase, amylase, trypsin, and chymotrypsin enzyme activity.




Figure 1. Average individual weight gain (g) of *Macrobrachium tenellum*: effect of chitin level in diet. Same letters represent no differences between treatments ($P = 0.05$). Bars represent amount of chitin in each treatment. Whisker lines are SE from the mean.

DISCUSSION

The water physical and chemical conditions were always maintained as required by *Macrobrachium tenellum* in culture, as stated in previous studies (García-Ulloa *et al.*, 2008; Espinosa-Chaurand *et al.*, 2011; Vega-Villasante *et al.*, 2011). For this species, the best growth used to be observed at a water temperature range from 28-32°C in agreement with Espinosa-Chaurand *et al.* (2011). Schiff & Hendrickx (1997) and Vega-Villasante *et al.* (2011) report that *M. tenellum* in culture grows very well with 5.0 mg O₂ L⁻¹. For pH, Hernández *et al.* (2007) and Vega-Villasante *et al.* (2011) suggest pH from 7.5 to 8.5 for *M. tenellum* cultivation conditions.

Two trends on survival rate were observed in present work. Prawns fed chitin at 0%, 5%, 10%, and 15% had an average survival of 83.3%. Prawns fed 20% and 25% chitin had lower survival (75% and 72.2%, respectively). Lower content of chitin apparently had no direct effect on survival, what is in agreement with previous studies. Kumar *et al.* (2006) and Shiao & Yu (1998) studied chitin and shrimp head meal, respectively, on survival of *M. rosenbergii* and *Penaeus monodon* (Fabricius, 1798). Kumar *et al.* (2006) find a survival of 40-80% while Shiao & Yu (1998) reported 74-84%, without significant differences or correlation between the amount of chitin and survival. These results are similar to present work although a precise comparison cannot be executed.

Previous studies with present species includes the works of Espinosa-Chaurand *et al.* (2012), who fed *M. tenellum* crude protein in diets (20, 25, 30, 35, and 40%) and maintained 95 to 100% survival, without statistical differences between treatments. In contrast, Ponce-

Palafox *et al.* (2006) reported 50% survival after prawns were fed a diet containing 25% of protein. The most common cause of death during cultivating prawns is from cannibalism. Common causes for this behavior are lack of shelters or large variations in size among individuals in the same tank (Nair *et al.*, 1999; García-Guerrero & Apun-Molina, 2008) observe that adding chitin to the diet reduced cannibalism in *M. rosenbergii*. It is important to state that the only chitin source was from diet, since every time one specimen moult, the exuviae was immediately removed from the tank. Survival of 75 and 72% in diets containing 20 and 25% chitin, respectively, had the lowest survival, despite having the highest growth. There were no significant differences in survival among treatments and it is suggested that different diets do not influence survival since chitin variation effect could be mostly on growth rate.

Present results suggest that the adding of chitin in the diet has a statistically significant effect on *M. tenellum* growth. The highest average individual weight gain (AIWG = g) was obtained with the 20% treatment. Chitin in the diet also increases chitinase activity, which was evident at day 30, when the highest weight gain per day (GWPD = g day⁻¹) occurred. The GWPD are in agreement with Ceccaldi (1989) and Zhang *et al.* (2014) who state that chitinase is essential for digestion and molting. However, molt frequency were not determined in present study so its direct relation with growth could not be determined even considering that in present study, a decrease in chitinase activity occurred when growth rate decreased. As Lemos *et al.* (2000) find, different growth stages engenders different feeding behavior and different digestion tract functional structures, thus, different digestion capabilities. This contributes to differences in enzyme activity, as observed in shrimp by Chang *et al.* (1988). Casillas-Hernández *et al.* (2002) observe that enzymatic, hormonal, and environmental issues are involved with molting, causing an indirect effect on growth rate. Chitin at 15 and 25% in the diet contribute to molting, as reported by Chang (1995). In present study, treatments containing 20, 25, and 15% chitin produced better weight gain in prawns. Prawns fed diets containing 10, 5, and 0% chitin had poorer weight gain. Statistical differences ($P = 0.046$) suggest that small variations in the amount of chitin may not affect weight gain in a measurable way, consequently, no differences in specific growth rates, increase in weight; feed conversion, and feed conversion efficiency were found. Kumar *et al.* (2006) report a similar trend in growth rate, using diets with different chitin content for *M. rosenbergii*. They report that diets containing 5% natural chitin (22% shrimp head meal), 10% natural

Table 2. Parameters of growth in weight for each treatment of chitin added to diet. AIWG: average individual weight gain, GWPD: gain weight per day, GW: gain weight in percentage, SGR: specific growth rate, FCR: feed conversion ratio and FER: feed efficiency ratio. Values with the same letter do not show statistically significant differences ($P = 0.05$), using Duncan multiple range test.

Treatment	AIWG (g)	GWPD (g day ⁻¹)	GW (%)	SGR (%)	FCR	FER
0% chitin	0.294 ± 0.025 ^b	0.0049 ± 0.00041 ^b	151.3 ± 15.1 ^a	1.48 ± 0.09 ^a	3.53 ± 0.31 ^a	0.296 ± 0.029 ^a
5% chitin	0.291 ± 0.033 ^b	0.0048 ± 0.00055 ^b	150.8 ± 17.0 ^a	1.45 ± 0.11 ^a	3.77 ± 0.51 ^a	0.295 ± 0.047 ^a
10% chitin	0.301 ± 0.036 ^b	0.0050 ± 0.00060 ^b	147.1 ± 17.4 ^a	1.42 ± 0.11 ^a	3.86 ± 0.40 ^a	0.272 ± 0.026 ^a
15% chitin	0.383 ± 0.038 ^{ab}	0.0064 ± 0.00063 ^{ab}	201.7 ± 23.2 ^a	1.74 ± 0.12 ^a	3.53 ± 0.30 ^a	0.293 ± 0.022 ^a
20% chitin	0.508 ± 0.101 ^a	0.0085 ± 0.0017 ^a	286.5 ± 72.6 ^a	1.88 ± 0.20 ^a	2.91 ± 0.39 ^a	0.390 ± 0.071 ^a
25% chitin	0.417 ± 0.073 ^{ab}	0.0070 ± 0.0012 ^{ab}	271.9 ± 66.2 ^a	1.78 ± 0.22 ^a	3.50 ± 0.55 ^a	0.329 ± 0.060 ^a




Figure 2. Gain weight per day (g day⁻¹) of *Macrobrachium tenellum*: effect of chitin amount in diet, at four intervals in trials. Chitin amount ranges from 0% to 25%.

chitin (44% shrimp head meal), and 5% purified chitin produce good weight gain and low feed conversion ratio. Shiau & Yu (1998) report a high weight gain in giant tiger shrimp (*Penaeus monodon*) fed a diet containing 22% chitin (equivalent to 5% purified chitin). They state that chitin in diets that is >22%, reduces weight gain. Fox (1993) mentions that natural chitin from shrimp head meal has high nutritional value for *P. monodon*, compared to purified chitin. In his study, there were not significant differences in growth where diets contained 0, 4, 8, 12, or 16% natural chitin. It seems that similar percentages of chitin can produce different results in different species. Prawns may take better advantage of dietary chitin compared to shrimp. Ceccaldi (1989) states that there is no difference in the use of chitin or other carbohydrates between shrimp and prawn digestive tracts; however, there are differences in tissue and biochemical activity (Cruz-Suarez, 2002). More research is required before conclusions on this topic.

In *M. tenellum*, Espinosa-Chaurand (2013) states that different concentrations of digestive enzymes in each phase of the life cycle are related to protein amount in diets and with circadian rhythm, since alterations of these variables affects proteolytic activity of chymotrypsin, trypsin, lipase, and amylase. Since we used identical protein diets, enzyme activity did not differ significantly between treatments, however, lowest activity was observed for all enzymes during the middle of the trial (day 30). Lower physiological activity was observed by Espinosa-Chaurand (2013). This depends on fitness, metabolic rate, and nutrient requirements of specific stages (Vega-Villasante *et al.*, 1999). Casillas-Hernández *et al.* (2006) report a relationship between feeding and enzyme activity, caused by demands of each developmental stage and production of biomass. Alexandre *et al.* (2014) and Cordova-Murrueta *et al.* (2003) also consider variations in enzyme activity as responses to growth stage, molting, digestion, and nutritional condition. These factors, together, determine enzyme activity independent of dietary ingredients. Other factors also influence enzyme activity. Since enzyme activity was not significantly different between treatments, growth also depends on other factors, such as availability and expenditure of energy, as in the case of *Penaeus monodon* (Chen & Lin, 1989).

It was also found an inverse relationship between enzymatic activity and digestion of proteins, lipids, and carbohydrates and the amount of chitin in the diet. Kumar *et al.* (2006) observe a decrease in digestibility with an increase in dietary chitin. They used purified chitin, which counteracts hydrolytic activity of other enzymes, probably because chitin was present in excess. Chitin content at 20% increases chitinase, compared to diets with less chitin ($P < 0.01$). Similar to their findings, we observed significant differences in enzyme activity between different stages ($P < 0.01$), chitinase activity peaked at day 30 and lower at the beginning and end of the trial. We also observed that




Figure 3. Effect amount of chitin in the diet on the digestive enzyme activity in *Macrobrachium tenellum*. a) Amylase, b) lipase, c) trypsin, and d) chymotrypsin. Samples taken on days 1, 30, and 60. Treatments are chitin amounts in diet (0%, 5%, 10%, 15%, 20%, and 25%). Whisker lines are SE from the mean.




Figure 4. Chitinase activity in *Macrobrachium tenellum*: effect of the amount of chitin in the diet. Treatments are chitin amounts in diet (0%, 5%, 10%, 15%, 20%, and 25%). Sampling times: days 1, 30, and 60. Whisker lines are SE from the mean.

from days 30 to 60, when chitinase activity increases, activity of all other digestive enzymes decreased. Additionally, when chitinase activity declines, lipase and amylase activity increased, supporting the hypo-

thesis of an inverse relation between chitinase and other digestive enzymes.

Another relationship was a decline in growth rate at high chitin content in the diet. The likely cause is an inability to absorb or metabolize big amounts of glucosamine if all dietary chitin was digested. This was observed in *M. rosenbergii* (Kumar *et al.*, 2006) and *P. monodon* (Fox, 1993). These authors suggest that this response could be ameliorated with a diet rich in lipids, proteins, and minerals. In fish diets, containing chitin, Lindsay (1984) and Danulat (1986) report that chitinase activity could be increased with dietary protein. However, in crustaceans, this did not occur with diets containing purified chitin, but this chitinase activity has occurred in the basal region of the peritrophic membrane in the gut (Ceccaldi, 1989; Alexandre *et al.*, 2014). It is likely that the same mechanism occurs in *M. tenellum* because chitin is re-absorbed when chitinase dissolves the old exoskeleton into a more soluble form in the integument, which is common in decapods (Dall *et al.*, 1990; Chang, 1995; Shechter *et al.*, 2007; Stillman *et al.*, 2008).

In addition to chitin level in the treatments, variations in chitinase activity may be influenced by prawn physiological reactions to water conditions, such as temperature and dissolved oxygen, as reported by Alexandre *et al.* (2014) concerning enzyme activity in the whiteleg shrimp *Litopenaeus vannamei*. In *Macrobrachium nipponense* (De Haan, 1849), Zhang *et al.* (2014) found variations in chitinase activity by diet induced, but not from size, developmental stage, hierarchical position, molting stage or acclimation time.

In summary, diets containing 20% natural chitin produced the best growth of *M. tenellum*. Further research should focus in the study of how some variables such as temperature or photoperiod in the culture may affect chitin digestion causing differences in enzymatic activity, particularly chitinases. In addition, further research is needed in the relationship between the molting cycle and chitin levels in the diet to determine the impact of the chitin amount and quality on the molt cycle and growth.

ACKNOWLEDGEMENTS

We thank Patricia Hinojosa Baltazar and Manuel Trasviña Castro of CIBNOR for technical assistance. Ira Fogel of CIBNOR provided editorial services. Funding was provided by Instituto Politécnico Nacional (CCA and PIFI 0002014). SIP-IPN and COFAA-IPN also provided financial support. R.B.S.R. thanks Instituto Tecnológico del Valle de Oaxaca for permission to work full time on this project.

REFERENCES

- Alexandre, D., R.A. Ozório, R.B. Derner, D.M. Fracalossi, G.B. Oliveira, R.I. Samuels & C.P. Silva. 2014. Spatial distribution of digestive proteinases in the midgut of the Pacific white shrimp (*Litopenaeus vannamei*) indicates the existence of endo-ectoperitrophic circulation in Crustacea. *Comp. Biochem. Physiol. B*, 172-173: 90-95.
- Anh, N.T.N., T.T.T. Hien, W. Mathieu, N.V. Hoa & P. Sorgeloos. 2009. Effect of fishmeal replacement with *Artemia* biomass as a protein source in practical diets for the giant freshwater prawn *Macrobrachium rosenbergii*. *Aquacult. Res.*, 40: 669-680.
- Bradford, M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248-254.
- Casillas-Hernández, R., F. Magallón, G. Portillo, O. Carrillo, H. Nolasco & F. Vega-Villasante. 2002. La actividad de proteasas, amilasas y lipasas durante los estadios de muda del camarón azul *Litopenaeus stylirostris*. *Rev. Invest. Mar.*, 23: 35-40.
- Casillas-Hernández, R., H. Nolasco-Soria, F. Lares-Villa, T. García-Galeano, O. Carrillo-Farnés & F. Vega-Villasante. 2006. Ritmo circadiano de la actividad enzimática digestiva del camarón blanco *Litopenaeus vannamei* y su efecto en el horario de alimentación. *Rev. Lat. Rec. Nat.*, 2(2): 55-64.
- Ceccaldi, H.J. 1989. Anatomy and physiology of digestive tract of crustacean decapods reared in aquaculture. *Advances in Tropical Aquaculture AQUACOP IFREMER, Actes de Colloque Tahiti*, 9: 243-259.
- Chang, E.S. 1995. Physiological and biochemical changes during the molt cycle in decapod crustaceans: an overview. *J. Exp. Mar. Biol. Ecol.*, 193: 1-14.
- Chang, E.S., S.A. Chang & E.P. Mulder. 2001. Hormones in the lives of crustaceans: an overview. *Am. Zool.*, 41: 1090-1097.
- Chang, S.M., S.M. Rankin & L.L. Keeley. 1988. Characterization of the molt stages in *Penaeus vannamei*: setogenesis and hemolymph levels of total protein, ecdysteroids, and glucose. *Biol. Bull.*, 175: 185-192.
- Chen, H.Y. & H.F. Lin. 1989. The effects of exogenous enzymes on the growth of early postlarval *Penaeus monodon*. In: R. Hiran & I. Hanyu (eds.). *Proceedings of the Second Asian Fisheries Forum, Tokyo*, pp. 17-22.
- Chih-Hui, L. & K. Chen-Chun. 2012. Effects of shrimp waste meal on growth performance and chitinase activity in juvenile cobia (*Rachycentron canadum*). *Aquacult. Res.*, 44: 1190-1195.
- Córdova-Murueta, J.H., F.L. García-Carreño & M.A. Navarrete-del-Toro. 2003. Digestive enzymes present in crustacean feces as a tool for biochemical, physiological, and ecological studies. *J. Exp. Mar. Biol. Ecol.*, 297: 43-56.
- Cortés-Jacinto, E., H. Villarreal-Colmenares, R. Civera-Cerecedo & R. Martínez-Córdova. 2003. Effect of dietary protein level on growth and survival of juvenile freshwater crayfish *Cherax quadricarinatus* (Decapoda: Parastacidae). *Aquacult. Nutr.*, 9: 207-213.
- Cruz-Suarez, E. 2002. Digestión en camarón y su relación con formulación y fabricación de alimentos balanceados. IV. Enzimas digestivas y estudios sobre digestibilidad para organismos acuáticos. VI Simposium Internacional de Nutrición Acuicola. *Proceedings of the VI International Symposium of Aquatic Nutrition*, 3-6 México, pp. 207-232.
- Dall, W., B.J. Hill, P.C. Rothlisberg & D.J. Staples. 1990. The biology of the Penaeidae. *Adv. Mar. Biol.*, 27: 1-489.

- Danulat, E. 1986. The effects of various diets on chitinase and b-glucosidase activities and the condition of cod, *Godus morhua*. J. Fish Biol., 28: 191-197.
- Espinosa-Chaurand, L.D. 2013. La actividad enzimática digestiva y su aplicación nutricional en el langostino *Macrobrachium tenellum* (Smith, 1871). Doctoral Dissertation. Universidad de Guadalajara, Mexico, 76 pp.
- Espinosa-Chaurand, L.D., C. Flores-Zepeda, H. Nolasco-Soria, O. Carrillo-Farnés & F. Vega-Villasante. 2012. Efecto del nivel proteico de la dieta sobre el desarrollo de juveniles de *Macrobrachium tenellum*. Revista MVZ Córdoba, 17(3): 3140-3146.
- Espinosa-Chaurand, L.D., M. Vargas-Ceballos, M. Guzmán-Arroyo, H. Nolasco-Soria, O. Carrillo-Farnés, O. Chong-Carrillo & F. Vega-Villasante. 2011. Biology and culture of *Macrobrachium tenellum*: state of the art. Hidrobiológica, 21(2): 99-117.
- Fox, C.J. 1993. The effect of dietary chitin on the growth, survival and chitinase levels in the digestive glands of juvenile *Penaeus monodon* (Fab.). Aquaculture, 109: 39-49.
- García-Carreño, F.L. & N.F. Haard. 1993. Characterization of proteinase classes in langostilla (*Pleuroncodes planipes*) and crayfish (*Pacifastacus astacus*) extracts. J. Food Biochem., 17: 97-113.
- García-Guerrero, M. & J.P. Apun-Molina. 2008. Density and shelter influence the adaptation to wild juvenile caque prawns *Macrobrachium americanum* to culture conditions. North Am. J. Aquacult., 70: 343-346.
- García-Ulloa, M., H. Rodríguez & T. Ogura. 2004. Calidad de huevecillos de dos especies de langostino (Palaemonidae) del género *Macrobrachium* (*M. rosenbergii* De Man, 1879 y *M. tenellum* Smith, 1871) variando la dieta de los reproductores: índices morfo-métricos. Avances Invest. Agropec., 8(2): 17-27.
- García-Guerrero, M.U., F. Becerril-Morales, F. Vega-Villasante & L.D. Espinosa-Chaurand. 2013. The genus *Macrobrachium* prawns with economic and commercial importance in Latin America: current knowledge, ecological role and conservation. Lat. Am. J. Aquat. Res., 41(4): 651-675.
- García-Ulloa, M., L.A. López-Aceves, T. Ponce-Palafox, H. Rodríguez-González & J.L. Arredondo-Figueroa. 2008. Growth of fresh-water prawn *Macrobrachium tenellum* (Smith, 1871) juveniles fed isoproteic diets substituting fish meal by soya bean meal. Braz. Arch. Biol. Technol., 51: 57-65.
- Gitte, M.J. & S.T. Indulkar. 2005. Evaluation of marine fish meat incorporated diets on growth and survival of post-larvae of *Macrobrachium rosenbergii* (de Man). Asian Fish. Sci., 18: 323-334.
- Hasan, B.M.A., B. Guha & S. Datta. 2012. Optimization of feeding efficiency for cost effective production of *Penaeus monodon* Fabricius in semi-intensive pond culture system. J. Aquacult. Res. Dev. 3: 149. doi: 10.4172/2155-9546.100149.
- Hernández, L., G. Murugan, G. Ruiz-Campos & A. Maeda-Martínez. 2007. Freshwater shrimp of the genus *Macrobrachium* (Decapoda: Palaemonidae) from the Baja California Peninsula, México. J. Crustacean Biol., 27: 51-69.
- Kumar, P., N.P. Sahu, N. Saharan, A.K. Reddy & S. Kumar. 2006. Effect of dietary source and level of chitin on growth and survival of post-larvae *Macrobrachium rosenbergii*. J. Appl. Ichthyol., 22: 363-368.
- Lemos, D., J.M. Ezquerro & F.L. García-Carreño. 2000. Protein digestion in penaeid shrimp: digestive proteinases, proteinase inhibitors and feed digestibility. Aquaculture, 86: 89-105.
- Lindsay, G.J.H. 1984. Distribution and function of digestive tract chitinolytic enzymes in fish. J. Fish Biol., 24: 529-536.
- Nair, K.K.C., M. Branislav, H. Rosenthal, K. Jayalakshmi & J. Nost. 1999. Experimental studies on the cannibalistic habit of *Macrobrachium rosenbergii* (de Man). Proceedings of the Fourth Indian Fisheries Forum, Kochi, pp. 227-232.
- Official Methods of Analysis of AOAC International (AOAC). 2012. AOAC International, Gaithersburg, Maryland [www.eoma.aoac.org].
- Ponce-Palafox, J.T., F.C. Arana-Magallón, H. Cabanillas-Beltrán & H. Esparza-Leal. 2002. Biological bases and techniques for the culture of freshwater prawns American Pacific native *Macrobrachium tenellum* (Smith, 1871) and *M. americanum* (Bate, 1968). Proceedings of the I Iberoamerican Virtual Congress of Aquaculture, pp. 534-546.
- Ponce-Palafox, J.T., J.M. García-Ulloa, J.L. Arredondo-Figueroa, D. Hernández-Ocampo, J. Díaz-Álvarez, G. Aldama-Rojas & H. Esparza-Leal. 2006. The cultivation of freshwater prawns *Macrobrachium tenellum* in earthen ponds. Proceedings of the IV Iberoamerican Virtual Congress of Aquaculture, pp. 655-660.
- Shechter, A., M. Tom, Y. Yudkovski, S. Weil, S.A. Chang, E.E. Chang, V. Chalifa-Caspi, A. Berman & A. Sagi. 2007. Search for hepatopancreatic ecdysteroid-responsive genes during the crayfish molt cycle: from a single gene to multigenicity. J. Exp. Biol., 210: 3525-3537.
- Schiff, H. & M.E. Hendrickx. 1997. An introductory survey of ecology and sensory receptors of Tropical Eastern Pacific crustaceans. Ital. J. Zool., 64: 13-30.
- Shiau, Y.S. & Y.P. Yu. 1998. Chitin but not chitosan supplementation enhances growth of grass shrimp, *Penaeus monodon*. J. Nutr., 128: 908-912.

- Somjit, K., K. Hara, M. Yoshimura, N. Matsuo, O. Kongpun & Y. Nozahi. 2005. Effect of shrimp chitin and shrimp chitin hydrolysate on the state of water and dehydration-induced denaturation of lizard fish myofibrillar protein. *Food Sci. Technol. Res. J.*, 11: 106-114.
- Stillman, J.H., J.K. Colbourne, C.E. Lee, N.H. Patel, M.R. Phillips, D.W. Towle, B.D. Eads, G.W. Gelembuik, R.P. Henry & E.A. Johnson. 2008. Recent advances in crustacean genomics. *Integr. Comp. Biol.*, 48: 852-868.
- Terwilliger, N.B. 1999. Hemolymph proteins and molting in crustaceans and insects. *Am. Zool.*, 39: 589-599.
- Vega-Villasante, F., H. Nolasco & R. Civera. 1993. The digestive enzymes of the Pacific brown shrimp *Penaeus californiensis* I. Properties of amylase activity in the digestive tract. *Comp. Biochem. Physiol. B*, 106: 547-550.
- Vega-Villasante, F., E.A. Martínez-López, M.C. Cortés-Lara, L.D. Espinosa-Chaurand & H. Nolasco-Soria. 2011. Growth and survival of *Macrobrachium tenellum* in experimental culture summer and autumn in the Tropical Mexican Pacific coast. *Trop. Subtrop. Agroecosyst.*, 14: 581-588.
- Vega-Villasante, F., I. Fernández, R.M. Preciado, M. Oliva, D. Tovar & H. Nolasco. 1999. The activity of digestive enzymes during the molting stages on the arched swimming *Callinectes arcuatus* Orway, 1863 (Crustacea: Decapoda: Portunidae). *Bull. Mar. Sci.*, 65: 1-9.
- Versaw, W.K., S.L. Cuppett, D.D. Winters & L.E. Williams. 1989. An improved colorimetric assay for bacterial lipase in nonfat dry milk. *J. Food Sci.*, 54: 1557-1558.
- Zhang, S., S. Jiang, Y. Xiong, H. Fu, S. Sun, H. Qiao, W. Zhang, F. Jiang, S. Jin & Y. Gong. 2014. Six chitinases from oriental river prawn *Macrobrachium nipponense*: cDNA characterization, classification and mRNA expression during post-embryonic development and moulting cycle. *Comp. Biochem. Physiol. B*, 167: 30-40.

Received: 31 March 2016; Accepted: 11 October 2016