



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e Tecnologia  
Avícolas  
Brasil

Fiorentin, L; Vieira, ND; Barioni Júnior, W

Use of lytic bacteriophages to reduce *Salmonella* Enteritidis in experimentally contaminated chicken cuts

Revista Brasileira de Ciência Avícola, vol. 7, núm. 4, outubro-diciembre, 2005, pp. 255-260

Fundação APINCO de Ciência e Tecnologia Avícolas  
Campinas, SP, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179713986010>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Use of Lytic Bacteriophages to Reduce *Salmonella* Enteritidis in Experimentally Contaminated Chicken Cuts

### ■ Author(s)

Fiorentin L  
Vieira ND  
Barioni Júnior W

Embrapa Suínos e Aves

### ■ Mail Address

Laurimar Fiorentin  
Embrapa Suínos e Aves  
BR 153 km 110 - CP 21 - Vila Tamanderá  
89.700 000 - Concórdia, SC, Brazil  
Phone: +55 (49) 442-8555  
Fax: +55 (49) 442-8559

E-mail: laurimar@cnpas.embrapa.br

### ■ Keywords

Bacteriophages, contamination, *Salmonella*, poultry, skin.

### ABSTRACT

Reducing *Salmonella* contamination in poultry is of major importance to prevent the introduction of this microorganism into the food chain. *Salmonellae* may spread during storage time (shelf life) whenever pre-harvest control fails or post-harvest contamination occurs. Therefore, preventive measures should also be used in the post-harvest level of poultry production in order to control salmonellae. Chicken skin samples were experimentally contaminated by immersing whole legs (thighs and drumsticks) in a suspension containing  $10^6$  colony forming units per milliliter (CFU/mL) of *Salmonella* Enteritidis phage type 4 (SE PT4) at the slaughter day. One day later, samples from one group were immersed in a suspension pool containing  $10^9$  CFU/mL of each of three wild salmonella-lytic bacteriophages previously isolated from feces of free-range chickens. *Salmonella* counting was performed at three-day intervals in the chicken legs stored at 5°C and showed a significant reduction ( $P < 0.05$ ) of SE PT4 in bacteriophage-treated cuts on days 3, 6 and 9 post-treatment. These findings suggest that the use of bacteriophages may reduce SE PT4 in chicken skin. Further studies are encouraged and might demonstrate the potential of this approach as an efficient and safe technique to be routinely used for *Salmonella* control in chicken products.

### INTRODUCTION

Pre-harvest *Salmonella* elimination might play a significant role in the prevention of pathogen introduction into the food chain and consequently in the reduction of food poisoning in humans (Seo *et al.*, 2000). It is more likely that the pre-harvest control of salmonellae is effective if a multi-factorial program is implemented. Good agricultural practices such as hazard analysis and critical control point (HACCP) (Nayak *et al.*, 2003), vaccination (Zhang-Barber *et al.*, 1999; Yamane *et al.*, 2000), probiotics, prebiotics and symbiotics (Van Immerseel *et al.*, 2002) have been used in poultry production as preventive measures if infection by *Salmonella* is likely to occur. In spite of that, no measure has been shown to be 100% effective and salmonellae remain as major contaminants to poultry (Persson & Jendteg, 1992; Mead *et al.*, 1999). The potential of phage therapy for *Salmonella* control has been recently assessed in the pre-harvest level of poultry production (Fiorentinn *et al.*, 2005; Sklar *et al.*, 2001).

Decontamination of chicken skin has been of major interest in applied research. Some chemical and physical procedures have been tested, such as carcass washing with aqueous solution of lactic acid (Xiong *et al.*, 1998), use of calcium hypochlorite, hydrogen peroxide and radiation (Nassar *et al.*, 1997), air chilling (Allen *et al.*, 2000), chlorine and sodium triphosphate (Xiong *et al.*, 1998; Whyte *et al.*, 2001), as well as the use



of natural products such as grapefruit seed extract (Xiong *et al.*, 1998). However, it has not been developed an efficient method that is completely devoid of chemicals and that might not cause food safety concerns.

In a previous study, we isolated and characterized *Salmonella*-lytic bacteriophages (Fiorentin *et al.*, 2004). Some of these bacteriophages have been administered *in vivo* to SE PT4-infected broilers and there was a reduction in the colony forming units of SE PT4 per gram of cecal contents by 3.5 orders of magnitude (Log CFU/g) (Fiorentin *et al.*, 2005). Other authors have also successfully reported reductions in *Salmonella* counts by using bacteriophages in chicken internal organs and feces (Toro *et al.*, 2005), skin (Goode *et al.*, 2003) or poultry products (Whichard *et al.*, 2003). The positive results led to the hypothesis that the bacteriophage isolated previously in our laboratory might also be efficient in reducing SE PT4 in chicken skin. Some advantages of administering phages onto the skin of poultry carcasses are that phages would not recycle in the host and therefore selection of resistant strains would be avoided. Besides, methods of biological control pose fewer risks to the consumer compared to chemical methods.

In the present study, chicken thighs and drumsticks were contaminated with SE PT4 and later treated with a panel of salmonellae-lytic bacteriophages isolated from free-range chickens.

## MATERIAL AND METHODS

### Chicken thighs and drumsticks

The study evaluated three groups as described in Table 1. The groups were comprised of 25 whole legs (thighs and drumsticks) with mean weight between 300 and 350 grams. The legs were collected during slaughter of a *Salmonella*-negative flock previously monitored using drag swabs (Waltman *et al.*, 1998). Chicken cuts (or parts) were experimentally contaminated by immersion in a suspension of phosphate buffered saline (PBS pH7.2) containing  $10^6$  CFU/mL of SE PT4 and allowed to dry for a few minutes. The samples of the three groups were transferred to sterile plastic bags, sealed and kept at 5°C. Afterwards, five samples per treatment were randomly taken at each three days for *Salmonella* and bacteriophage counts.

### Contamination with *Salmonella*

SE PT4 isolate P125589 was kindly provided by Dr

Paul Barrow (ARFC Institute for Animal Health, Houghton Laboratory, Cambridge, England). It was originally isolated by Dr B. Rowe (Central Public Health Laboratory, London, UK) from a case of human food poisoning (Barrow & Lovell, 1991).

A fresh colony of SE PT4 was inoculated into 10mL of nutrient broth (NB, 1g/L beef extract, 2g/L yeast extract, 5g/L peptone, 5g/L sodium chloride, pH 7.0) and incubated overnight at 37°C under shaking (200rpm). The culture was frozen at -80°C and one aliquot was used for enumeration of viable cells, by counting colonies grown from tenfold dilutions streaked onto nutrient agar and incubated for 24h at 37°C. After counting, the original culture was diluted with sterile buffered saline (1 L) to produce a solution containing  $10^6$  CFU/mL that was used to contaminate the chicken cuts.

**Table 1** – Treatments used to assess the effect of bacteriophages on the reduction of *Salmonella* Enteritidis phage type 4 contamination in broiler skin samples.

Group	Cuts	Treatment
1	25	Non-contaminated and non-treated
2	25	Contaminated by immersion on a suspension containing $10^6$ CFU/mL of <i>Salmonella</i> Enteritidis phage type 4 at slaughter day
3	25	Contaminated as in group 2, treated one day later by immersion in a suspension containing $10^9$ PFU/mL of a mixture of bacteriophages CNPSA1, CNPSA3 and CNPSA4

### Treatment with bacteriophages

Bacteriophages CNPSA 1, CNPSA3 and CNPSA4 were isolated from feces of free-range chickens in Brazil and characterized as described elsewhere (Fiorentin *et al.*, 2004). Since resistance to bacteriophages may emerge in growing populations of bacteria, we decided to use a pool of three different viruses. Frozen bacteriophage stocks were amplified on overlay cultures of SE PT4 prepared with Nutrient Broth (NB) containing 0.7% agarose. Enough bacteriophage particles were then used to inoculate one liter of SM buffer (5.8g/L NaCl, 2.0g/L MgSO<sub>4</sub>·7H<sub>2</sub>O, 5.0mL/L of a 5% solution of gelatin, 50mL/L of 1M Tris-HCl pH 7.5) to a concentration of  $10^9$  plaque forming units per milliliter (PFU/mL), which results in a multiplicity of infection of 1,000 (MOI: 1,000), i.e., a thousand PFU of bacteriophages per CFU of SE PT4 was used to contaminate the chicken parts. Bacteriophage titers were determined using tenfold dilutions of the virus preparation mixed to SE PT4 in log-phase growth (10μL : 250μL). The mixtures were



incubated at 37°C for 20 minutes and then mixed with 7mL of melted NB-agarose (45°C), thoroughly homogenized and overlaid onto 10cm-diameter Nutrient Agar plates (NB added with 15g/L bacteriologic agar) (Kudva *et al.*, 1999). Plaques of lysis were counted after incubation for 24 hours at 37°C. Bacteriophage titers were obtained by multiplying the number of plaques in each dilution by the dilution factor. Cuts were immersed in the bacteriophage suspension, allowed to dry for a few minutes and replaced into the plastic bag for storage at 5°C for 15 days.

### Experimental design

The number of washable viable *Salmonella* cells and bacteriophage particles were assessed in five leg samples taken from each group at three-day intervals. The cuts were randomly chosen and submitted to *Salmonella* and bacteriophage assessment as described below.

### Salmonella isolation

Samples from Groups 2 and 3 were individually rinsed in sterile plastic bags containing 100mL of buffered peptone water. The number of *Salmonella* colony forming units (CFU) in the rinse solution (rinse peptone water) was determined by multiplying the number of viable cells in 1mL of the solution multiplied by 1,000. Rinse peptone water was diluted tenfold in buffered saline and 100µL of each dilution were spread onto 10cm-diameter brilliant green agar plates (BGA) supplemented with novobiocin (40µg/mL) and incubated at 37°C for 48h. *Salmonella* CFU per cut was determined by the number of colonies in plates with 30 to 300 colonies multiplied by the dilution factor. The theoretical limit of detection of this method is  $30 \times 10^2$  CFU per cut, which means that negative results in Groups 2 and 3 would actually represent <3,000 CFU per cut. Slight pink-white opaque colonies surrounded by red medium grown on BGA were confirmed as *Salmonella* by slide agglutination test with polyvalent anti-somatic serum.

Cuts from all groups were also submitted to qualitative analysis of *Salmonella*. Rinse peptone water was incubated at 37°C for 24h for pre-enrichment. Afterwards, 0.3 mL of each culture was enriched in 2.7 mL of Rappaport-Vassiliadis Soya peptone broth (RVS) and incubated at 42°C for 24h. A loopful was then streaked onto BGA plates and incubated at 37°C for 24h. *Salmonella* colonies were identified as described above.

### Bacteriophage titering

Cuts from Group 3 were individually rinsed in 100mL of SM buffer and total bacteriophage PFU was titered as follows. One milliliter of rinse SM buffer was treated with 5% chloroform to lyse all cells and 10µL were serially diluted tenfold in sterile SM buffer. From each dilution, 10µL were then mixed with 250µL of SE PT4 in log-phase growth and incubated for 20 minutes at 37°C to allow infection of *Salmonella* cells by bacteriophages. Afterwards, mixtures were added to 7mL of NB containing 0.7% agarose, thoroughly homogenized and overlaid onto 10cm-diameter nutrient agar plates, which were then incubated for 24h at 37°C. Plaques of lysis were counted on the cultures with 30 to 300 plaques. The number of bacteriophages was determined by multiplying the number of plaques by 100 and expressed as total PFU per mL. Total PFU per chicken cut was obtained multiplying total PFU by 100.

Cuts from Groups 1 and 2 were submitted to qualitative analysis to confirm their bacteriophage-free status. One milliliter of rinse SM buffer from each chicken cut was treated with 5% chloroform, vortexed, centrifuged (12,000 x g for 4 minutes) and 100µL of the supernatant was added to 900µL of a SE PT4 culture, which was incubated at 37°C for 24h. After incubation, this preparation was also treated with 5% chloroform to lyse all bacteria, centrifuged and 10µL of the supernatant was used to prepare overlaying SE PT4 cultures as described before. Positive results were identified by the presence of lysis plaques with about 1cm of diameter on the *Salmonella* lawn.

### Statistical analysis

Analysis of variance followed by t-test was used to compare means of CFU of SE PT4 per piece according to the experimental design. Statistical analysis was performed using a commercial package (SAS, 2001).

## RESULTS AND DISCUSSION

### Salmonella isolation

SE PT4 was recovered from all cuts in all five assessments using the samples of rinse peptone water from Group 2 and Group 3 submitted to selective enrichment. Sampling was performed at each three days after treatment until day 15 (except for day 3). Therefore, the results indicate a relative long period of SE PT4 survival considering the shelf life of refrigerated products. Survival for long periods has been recognized



as an important feature of salmonellae; even in low temperatures, salmonellae were able to survive in non-sanitized poultry carcasses at least for 26 days (Thomson *et al.*, 1979).

The presence of SE PT4 in the bacteriophage-treated cuts indicates that the conditions used in this study did not allow bacteriophages to eliminate *Salmonella* from poultry skin. However, it is unlikely that any single technique applied to poultry carcasses would remove *Salmonella* completely from the skin, nor would *Salmonella* be naturally removed from contaminated skin only by refrigeration. We have previously reported that bacteriophages orally administered to SE PT4-infected broilers reduced the numbers of *Salmonella* significantly, but could not remove completely the bacteria from chicken cecae (Fiorentin *et al.*, 2005). The same is probably also true in other situations, such as when bacteriophages are applied to poultry skin. In regard to food safety, it is extremely important to develop techniques that completely prevent food poisoning in humans. However, it seems hard to achieve this objective using only a single technique and bacteriophages should probably be used together with other methods to completely remove *Salmonella* from chicken skin. The present study is our first evidence of *Salmonella* reduction in chicken skin by the use of bacteriophages. The findings encourage us to perform further research in order to achieve more efficiency regarding bacteriophage treatment applied on SE PT4-contaminated cuts.

### Bacteriophage isolation

Rinse peptone water from cuts from Groups 1 and 2 were inoculated with SE PT4 to permit bacteriophage amplification, in case they were present. This qualitative analysis confirmed the bacteriophage-free status of both groups. On the other hand, bacteriophages were isolated from all pieces from Group 3 in all five samplings according to the qualitative analysis. These results indicate that bacteriophages were not removed from broiler skin contaminated with *Salmonella* within a 15 day-period, which might have resulted from bacteriophage multiplication on the bacteria or because bacteriophages were not inactivated by natural components of the chicken skin.

A panel of three bacteriophages was used to avoid selection of strains of SE PT4 resistant to a particular virus. Although this was a necessary approach, it prevented us from knowing whether one specific bacteriophage lasts longer in the cuts or if the three of them have been recovered from all samples at the same

magnitude. However, we have previously reported that the three bacteriophages share similar lytic properties and multiplication rates when cultured in SE PT4 (Fiorentin *et al.*, 2004) and this might also occur *in vivo*.

### *Salmonella* counting

Enumeration of *Salmonella* was performed only on samples from Group 2 and Group 3, because Group 1 was not contaminated and yielded negative results in the qualitative analysis throughout the experiment. Total CFU ranged from  $0.66 \pm 0.05 \times 10^8$  CFU to  $4,840 \pm 461.09 \times 10^8$  CFU per cut in the first and last sampling of Group 2 (SE PT4 contaminated group), respectively. In the SE PT4-contaminated and bacteriophage-treated cuts (Group 3), total CFU was  $0.29 \pm 0.08 \times 10^8$  CFU and  $3,920 \pm 738.51 \times 10^8$  CFU per cut in the first and the last sampling days, respectively. There was an increasing trend in total CFU in groups of samples taken at each three days. These concentrations of CFU were high, and may be explained because of the high CFU dose used to contaminate the pieces and also because the experimental conditions prevented competition between *Salmonella* and other organisms (Coleman *et al.*, 2003). However, lower magnitudes of total CFU were seen in the bacteriophage-treated group (Group 3) at all sampling days, except for 12 and 15 days post-treatment (Table 2 and Figure 1). The means of treated and non-treated parts were compared by t-test (SAS, 2001), and P values were highly significant on days 3, 6 and 9, indicating higher efficiency of bacteriophages in reducing *Salmonella* CFU within a 9-day period (Table 2).

*Salmonella* numbers in samples from Group 2 were of lower magnitude at 12 days post-inoculation compared to the samples of Group 3 or even to the samples of Group 2 in the previous samplings. This result was probably caused by a technical variation that occurred during dilution or plating. Therefore, we suppose that bacteriophages might have actually had an effect for longer than 9 days. It is also possible, however, that *Salmonella* contamination was lower or bacteriophage effects were greater in the chicken cuts randomly sampled as one group on day 12. The curve presented in Figure 1 suggests that the data of Group 2 on day 12 should logically fall between  $4,000$  and  $8,000 \times 10^8$  CFU.


Reduction of CFU on treated cuts was about 4.49 times on day 9 post-treatment (Table 2). Such a reduction is enough to demonstrate a cause-and-effect relationship, but additional research is required in order



**Table 2** - Mean  $\pm$  standard deviation of total colony forming units (CFU  $\times 10^8$ ) recovered from the rinse peptone water of five cuts contaminated with *Salmonella* Enteritidis phage type 4 (Group 2) or contaminated and treated with bacteriophages (Group 3).

Days after treatment	Group 2*	Group 3	P value**	Reduction***
3	0.66 $\pm$ 0.05	0.29 $\pm$ 0.08	0.0334	2.27
6	340.2 $\pm$ 124.16	74.6 $\pm$ 20.37	0.0026	4.56
9	7,734 $\pm$ 2,875.02	1,720 $\pm$ 360.85	0.0195	4.49
12	772 $\pm$ 51.52	1,714 $\pm$ 813.74	0.269	-0.45
15	4,840 $\pm$ 461.09	3,920 $\pm$ 738.51	0.552	0.97

\*Non-contaminated and non-treated pieces (Group 1) were negative throughout the experiment. \*\*t-test. \*\*\*Difference in mean values between Group 3 (contaminated, treated) and Group 2 (contaminated, non-treated).



**Figure 1** - Mean  $\pm$  standard deviation of total colony forming units (CFU  $\times 10^8$ ) of SE PT4 recovered from cuts contaminated with *Salmonella* (Group 2) or contaminated and treated with bacteriophage (Group 3). Differences between means were statistically significant on days 3, 6 and 9.

to achieve *Salmonella* reduction by several orders of magnitude. Goode *et al.* (2003) carried out a research with the same SE PT4 strain used in the present study and reported that a bacteriophage used for phage typing have also not reduced CFU by several orders of magnitude. Since those authors used a different technique and bacteriophage efficiency assessment was performed only over a 48h-experimental period, any direct comparison between the two experiments would not be adequate. Nevertheless, based on the results from the present study and those reported by Goode *et al.* (2003), it seems reasonable to assume that additional research is necessary to better understand how bacteriophages can be used more efficiently to reduce *Salmonella* in chicken skin.

The fact that total CFU increased even in treated cuts leads us to suppose that higher concentrations of bacteriophages might prevent such multiplication of SE PT4 on the skin. However, this is our first report on the use of bacteriophages to reduce *Salmonella* presence in poultry skin. The convincing results presented herein encourage further research in order

to achieve greater reductions in contamination of poultry carcasses or cuts by *Salmonella*. The use of higher concentrations of bacteriophages, either alone or coupled with other methods, might determine more efficient reductions of SE PT4 contamination of poultry skin.

### Enumeration of bacteriophages

Concentration of bacteriophages recovered from the rinse water in Group 3 ranged from 5.8 to 25.4  $\times 10^5$  PFU per cut (Table 3). The observed variation is probably inherent to the technique used for titrating the viruses, once the difference between titers were still smaller than one order of magnitude. This is a low recovery rate compared to the concentration of bacteriophage ( $10^9$ /mL) used to treat the cuts contaminated with SE PT4. A possible explanation is that only a small portion of SM buffer might have actually remained in the cuts after dipping or perhaps bacteriophages adhered poorly to components on the chicken skin; both conditions would have resulted in lower MOI than the expected MOI of 1,000. However, it is noteworthy that bacteriophages were viable and showed similar titers for 15 days at 5°C even in contact with chicken skin.

**Table 3** - Means of total plaque forming units (PFU) of bacteriophages recovered from rinse peptone water of contaminated and treated cuts (Group 3).

Days after treatment	Total PFU recovered per cut*
3	15 $\times 10^5$
6	5.8 $\times 10^5$
9	25.4 $\times 10^5$
12	8.4 $\times 10^5$
15	13 $\times 10^5$

\*Obtained by titrating 10 $\mu$ l of rinse solution multiplied by the dilution factor.

The fact that bacteriophage titers did not increase in Group 3 might have been caused by the reduction observed on CFU of SE PT4. It is also possible that these



bacteriophages did not target any other bacteria present on the chicken skin, otherwise they would have shown higher PFU per cut with longer shelf storage time.

## CONCLUSIONS

A panel of bacteriophages reduced SE PT4 countings in experimentally contaminated chicken parts stored at 5°C. We demonstrated that *Salmonella* CFU was reduced in the bacteriophage-treated cuts on days 3, 6 and 9 post-treatment when compared to their non-treated counterparts.

## REFERENCES

- Allen VM, Corry JE, Burton CH, Whyte RT, Mead GC. Hygiene aspects of modern poultry chilling. *International Journal of Food Microbiology* 2000; 58:39-48.
- Barrow PA, Lovell MA. Experimental infection of egg-laying hens with *Salmonella enteritidis* phage type 4. *Avian Pathology* 1991; 20:335-348.
- Coleman ME, Sandberg S, Anderson SA. Impact of Microbial Ecology of Meat and Poultry Products on Predictions from Exposure Assessment Scenarios for Refrigerated Storage. *Risk Analysis* 2003; 23:215-228.
- Fiorentin L, Vieira ND, Barioni Júnior W. Oral treatment with bacteriophages reduces the concentration of *Salmonella* Enteritidis PT4 in cecal contents of broilers. *Avian Pathology* 2005; 34(3):1-6.
- Fiorentin L, Vieira ND, Barioni Júnior W, Barros S. *In vitro* characterization and *in vivo* properties of *Salmonellae* lytic bacteriophages isolated from free-range chickens. *Brazilian Journal of Poultry Science* 2004; 6:105-112.
- Goode D, Allen VM, Barrow PA. Reduction of experimental *Salmonella* and *Campylobacter* contamination of chicken skin by application of lytic bacteriophages. *Applied Environmental Microbiology* 2003; 69:5032-5036.
- Kudva IT, Jelacic S, Tarr PP, Youderian P, Hovne CJ. Biocontrol of *Escherichia coli* O157 with O157-Specific bacteriophages. *Applied and Environmental Bacteriology* 1999; 65:3767-3773.
- Mead PS, Slutsker L, Dietz V, McCaig LF, Bresee JS, Shapiro C, Griffin PM, Tauxe RV. Food-related illness and death in the United States. *Emerging Infectious Diseases* 1999; 5(5):607-625.
- Nassar TJ, al-Mashhadi AS, Fawal AK, Shalhat AF. Decontamination of chicken carcasses artificially contaminated with *Salmonella*. *Revue Scientifique et Technique* 1997; 16(3):891-7.
- Nayak R, Kenney PB, Keswani J, Ritz C. Isolation and characterization of *Salmonella* in a turkey production facility. *British Poultry Sciences* 2003;44(2):192-202.
- Persson U, Jendteg S. The economic impact of poultry-borne salmonellosis: how much should be spent on prophylaxis? *International Journal of Food Microbiology* 1992; 15(3-4):207-213.
- SAS Institute. System for Microsoft Windows, Release 8.2. Cary, North Carolina, USA; 1999-2001. CD-ROM.
- Seo KH, Holt PS, Gast RK, Hofacre CL. Elimination of early *Salmonella enteritidis* infection after treatment with competitive-exclusion culture and enrofloxacin in experimentally infected chicks. *Poultry Sciences* 2000; 79:1408-1413.
- Sklar IB, Joerger RD. Attempts to utilize bacteriophage to combat *Salmonella enterica* serovar *Enteritidis* infection in chickens. *Journal of Food Safety* 2001; 21:15-29.
- Thomson JE, Bailey JS, Cox NA. Phosphate and heat treatments to control *Salmonella* and reduce spoilage and rancidity on broiler carcasses. *Poultry Sciences* 1979; 58:139-143.
- Toro H, Price SB, McKee AS, Hoerr FJ, Krehling J, Perdue M, Bauermeister L. Use of bacteriophages in combination with competitive exclusion to reduce *Salmonella* from infected chickens. *Avian Diseases* 2005; 49(1):118-24.
- Van Immerseel F, Cauwerts K, Devriese LA, Haesenbrouck F, Ducatelle R. Feed additives to control *Salmonella* in poultry. *World's Poultry Science Journal* 2002; 58:501-513.
- Waltman WD, Gast RK, Mallinson ET. Salmonellosis. In: Swayne de, editor. A laboratory manual for the isolation and the identification of avian pathogens. 4<sup>th</sup> ed. Kennett Square: The American Association of Avian Pathologists; 1998. p.4-13.
- Whichard JM, Sriranganathan N, Pierson FW. Suppression of *Salmonella* growth by wild-type and large-plaque variants of bacteriophage Felix O1 in liquid culture and on chicken frankfurters. *Journal of Food Protection* 2003; 66(2):220-225.
- Whyte P, Collins JD, McGill K, Monahan C, O'Mahony H. Quantitative investigation of the effects of chemical decontamination procedures on the microbiological status of broiler carcasses during processing. *Journal of Food Protection* 2001; 64(2):179-183.
- Xiong H, Li Y, Slavik MF, Walker JT. Spraying chicken skin with selected chemicals to reduce attached *Salmonella typhimurium*. *Journal of Food Protection* 1998; 61:272-275.
- Yamane Y, Leonard JD, Kobatake R, Awamura N, Toyota Y, Ohta H, Otsuki K, Inoue T. A case study on *Salmonella enteritidis* (SE) origin at three egg-laying farms and its control with an *S. enteritidis* bacterin. *Avian Diseases* 2000; 44(3):519-526.
- Zhang-Barber L, Turner AK, Barrow PA. Vaccination for control of *Salmonella* in poultry. *Vaccine* 1999; 17:2538-2545.