



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e Tecnologia
Avícolas
Brasil

Cardoso, WM; Aguiar Filho, JLC; Romão, JM; Salles, RPR; Câmara, SR; Siqueira, AA; Oliveira, WF;
Sobral, MHNR; Texeira, RSC

Interference of infectious bursal disease virus on antibody production against Newcastle disease and
infectious bronchitis virus

Revista Brasileira de Ciência Avícola, vol. 8, núm. 3, julio-septiembre, 2006, pp. 177-182
Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, SP, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179713991007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Interference of Infectious Bursal Disease Virus on Antibody Production against Newcastle Disease and Infectious Bronchitis Virus

■ Author(s)

Cardoso WM^{3*}
Aguar Filho JLC¹
Romão JM²
Salles RPR¹
Câmara SR¹
Siqueira AA²
Oliveira WF¹
Sobral MHNR¹
Texeira RSC¹

¹ Postgraduate students of Veterinary Sciences of Ceara State University.

² Undergraduate students of Veterinary Faculty of Ceara State University

³ Professor of Veterinary Faculty of Ceara State University

■ Mail Address

William Maciel Cardoso
Av. Rogaciano Leite, 200
Aptº 1303 Bl. Tulipe, Bairro Salinas
60.810-000. Fortaleza, CE, Brazil
Phone/Fax: 55 (085) 3241 1307 ou 3299 2748
ou 9989 4742

E-mail: william.maciell@uol.com.br
ornito_uece@yahoo.com

■ Keywords

Antigenic variation, polyvalent vaccinations, viral interference.

■ Acknowledges

The authors thank Dr. Ronald Carneiro Câmara of the poultry company CIALNE (Northeast Company of Aliments), the BIOLAB Laboratory of Animal Pathology S/C Ltda and FUNCAP - Cearense Foundation for Research Support.

ABSTRACT

This work has the objective of verifying the interference of infectious bursal disease virus in the antibody production against Newcastle disease virus and infectious bronchitis virus. The experiment was carried out with 640 day-old-chicks from a 42 weeks old hen flock. The birds were separated into eight experimental groups (n=80/group) and were submitted to different combinations of vaccinations, with live vaccines, to Newcastle disease, avian infectious bronchitis, and infectious bursal disease with diverse combinations of days of vaccination. We verified that the utilization of polyvalent vaccinal programs have a different efficacy comparing to monovalent vaccinations when Newcastle disease, infectious bronchitis, and infectious bursal disease vaccinations are applied. This way, the use of vaccinations to infectious bursal disease in polyvalent vaccinal programs is desirable due to improvement of NDV response with the presence of IBV by the probable reduction of interference of IBV under NDV.

INTRODUCTION

The infectious bursal disease (IBD) is a highly contagious viral and acute infection with tropism for lymphoid tissue, principally for bursa of Fabricius, in which the virus promotes the cell destruction. This disease is caused by IBDV that belongs to the genus Avibirnavirus of the family Birnaviridae (Murphy *et al.*, 1995) and was identified by the first time in Delmarva region of United States of America. It is a high importance problem for poultry industry. Current thinking is that protection against IBDV may be mediated primarily by anti-IBDV antibodies (Fussell, 1998; Luttkick, 1997; Vakharia *et al.*, 1994). IBDV vaccines used in commercial flocks are selected by the ability of the vaccines to induce vigorous antibody responses (Lasher and Shane, 1994), this way it has been used live and inactivated vaccines from serotype 1 (Jackwood and Saif, 1987). Six variants of this serotype were identified by the virus neutralization test. This antigenic variation can induce failures on the vaccination processes due to the difference of antigenic structures between vaccinal and wild viruses (Cao *et al.*, 1998; Jackwood and Saif, 1987; Kibenge *et al.*, 1988; van den Berg, 2000). Besides the antigenic variation, other factors can interfere on efficacy of a vaccinal program, among them, the viral interference. This phenomenon can occur among different serotypes of the same virus, for example infectious bursal disease virus (IBDV) with intermediate and pathogenic strains (Ashraf *et al.*, 2005), Reovirus (Whitaker-Dowling *et al.*, 1987), Poliovirus (Sabin, 1959) and Avian Influenza virus (Whitaker-Dowling, 1992). It can also occur between different viruses, as between infectious bronchitis virus (IBV) and Newcastle disease virus (NDV) (Cardoso *et al.*, 2005) or avian pneumovirus (Cook *et al.*, 2001). The occurrence of interference among



avian polyvalent vaccines can be promoted by the competition among the vaccinal viruses to the same receptors (Sabin, 1959).

This work has the objective of verifying the occurrence of interference of infectious bursal disease virus in the antibody production against Newcastle disease virus and infectious bronchitis virus.

MATERIAL AND METHODS

Birds

The experiment was carried out with 640 day-old-chicks from a 42 week-old hen flock. The chicks received no previous vaccination and were reared until 50 days of age in experimental broiler houses. The birds were bred simulating industrial conditions, with adequate management, water and feed ad libitum in a 10birds/m² density.

Treatments and vaccination

The birds were separated into eight experimental groups (n=80/group): Control Group to Maternal Antibodies (CG-Mab), Control Group to Avian Infectious Bronchitis (CG-AIB), Control Group to Newcastle disease (CG-ND), Control Group to Infectious Bursal Disease (CG-IBD), Group without IBD vaccination (T0), and the groups with one, two and three IBD vaccinal doses (T1), (T2), and (T3), respectively. All groups, except CG-Mab, were submitted to treatments with vaccinations. The treatments followed the methodology described in the Table 1.

The vaccination was performed by instillation of a 0.03 mL drop by ocular route. All vaccines belonged to the same laboratory and the same group of vaccinators did the vaccinations. The following vaccinal strains were used: HB1 (10^{6.5}) to Newcastle disease, H120 (10^{3.5}) to infectious bronchitis, and Lukert -intermediate classic (10^{3.0}) to infectious bursal disease.

Blood collection and serological tests

The control group to maternal antibodies was

submitted to blood collections at 1st, 25th, 35th and 45th days of age to verification of maternal antibody levels to IBV, NDV, and IBDV. The other groups were submitted to blood collections at 35th and 45th days of age. All blood samples were obtained from the brachial vein and the sera were collected, maintained adequately refrigerated to serological tests. Each serum was identified according to the number of the birds.

The serum samples were analyzed by HI (Haemagglutination inhibition) test to detection of antibodies against NDV and indirect ELISA (enzyme-linked immunosorbent assay) (Kirkegaard & Perry Laboratories - KPL) to detection of antibodies against IBV and IBDV.

Statistical Analyses

The titers obtained by ELISA and HI tests were submitted to the statistic program SAS/STAT 95 from SAS Institute Inc. (User's guide: statistic and graphics, 1995). The results of antibody titers were submitted to Variance Analyses. The antibody titers were transformed by the logarithmic function Log₁₀x+1 to submit the data to variance analyses. The means were compared through the test 't' student with significance level of 5%.

RESULTS


The Graphic 1 presents the maternal antibody titers of control group against Newcastle disease virus, infectious bursae disease virus, and infectious bronchitis virus.

The maternal antibodies to NDV were null since the beginning up to the end of the experiment. The titers against IBDV and IBV were significant. The first one has GMT 3200 by the first day decreasing to zero at 25 days of age and maintained this level until the end of experiment. The maternal antibody curve to IBV was similar to IBDV, but the titers in the first day were around GMT 4500.

Table 1 - Experimental treatments.


Groups	Vaccinations		
	1 st day	8 th day	16 th day
CG-Mab	N/V	N/V	N/V
CG-IB	N/V	H ₁₂₀	N/V
CG-ND	N/V	HB1	N/V
CG-IBD	Lukert-intermediate classic	Lukert-intermediate classic	N/V
T0	N/V	H ₁₂₀ + HB1	N/V
T1	Lukert-intermediate classic	H ₁₂₀ + HB1	N/V
T2	Lukert-intermediate classic	H ₁₂₀ + HB1	Lukert-intermediate classic
T3	Lukert-intermediate classic	H ₁₂₀ + HB1 + Lukert-intermediate classic	Lukert-intermediate classic

N/V= no vaccination.



Graphic 1 - Maternal antibody titers of control group against Newcastle disease virus, infectious bronchitis virus, and infectious bursal disease virus.


The Graphic 2 presents antibody titers ($\text{Log}_{10} + 1$) to NDV at 35 and 45 days of age to the groups that were submitted to vaccinations to Newcastle disease, infectious bronchitis, and one, two or three vaccinations to infectious bursal disease.



Graphic 2 - Antibody titers to Newcastle disease virus of experimental groups.

The birds of control group to Newcastle disease showed the antibody titer (2,318) at 45 days, being statistically superior to T0 antibody titer (0,753) at 45 days. We can observe the antibody titer increasing with the addition of infectious bursal disease vaccinations, one, two or three vaccinations. At 45 days, the anti-NDV antibodies of T2 and T3 were statistically superior to T1 that received only one IBDV vaccination.


The Graphic 3 presents antibody titers ($\text{Log}_{10} + 1$) to IBV at 35 and 45 days of age to the groups that were submitted to vaccinations to Newcastle disease, infectious bronchitis, and one, two or three vaccinations to infectious bursal disease.



Graphic 3 - Antibody titers to Infectious Bronchitis virus of experimental groups.

The Graphic 3 shows that the antibody titers to IBV were similar for most of the groups at 35 and 45 days of age, however T3 at 45 days and T2 at 35 days presented statistic difference ($p < 0.05$) when compared to the other groups in the same age, being the lowest level at 45 days and the highest at 35 days, respectively.

The Graphic 4 presents antibody titers ($\text{Log}_{10} + 1$) to IBDV at 35 and 45 days of age to the groups that were submitted to vaccinations to Newcastle disease, avian infectious bronchitis, and one, two or three vaccinations to infectious bursal disease.



Graphic 4 - Antibody titers against infectious bursal disease virus of experimental groups.

We can observe that the graphic 4 shows that IBDV antibody titers presented no significant difference ($p < 0.05$) when the birds received one, two or three IBD vaccinations.

DISCUSSION

The maternal antibodies to infectious bursal disease in day-old-chicks of CG-Mab group belonging from IBD



immunized breeders were elevated (GMT 3200) according to Sharma *et al.* (1989), and decreased to zero by 25 days of age, agreeing with Al-Natour *et al.* (2004) that found similar results, with maternal antibody titers around GMT 3400 at the first day with a decrease reaching zero around 28 days of age.

This expressive antibody titer drop has been explained by many authors (Knezevic *et al.*, 1999; Knoblich *et al.*, 2000; Alam *et al.*, 2002) that concluded that vaccination does not accelerate decrease in maternal antibodies if chicks are vaccinated at one day of age. Similarly to results reported by Knezevic *et al.* (1999), chicks with passive immunity and vaccinated with an intermediate IBDV strain in the first day of age showed no increase in antibody titers.

In the same way, IBV antibodies decreased to zero around 25 days, like the results observed by Gelb *et al.* (1998). The NDV antibodies were null since the first day up to the end of experiment. The maternal antibody curves to IBDV, NDV, and IBV indicated that, to these three analyzed diseases, there was no interference of wild or vaccinal virus challenges. The relation between the expressive IBDV maternal antibody decrease found in our experiment with absence of interference of wild or vaccinal virus challenges can be explained by inexistence of wild or vaccinal virus challenges or neutralization of the virus by the high IBDV maternal antibodies as demonstrated by Moraes *et al.* (2005) that verified that vaccinated and unvaccinated chicks with high antibody titers (3.4Log10) in the first day of age were protected against the disease after challenge with a very virulent strain of IBD.

We observed that the group of birds that was submitted for just a vaccination to AIB and ND at eighth day presented NDV antibodies inferior to the group that received only a ND vaccination in the same day. Many researchers reported this situation (Beard, 1967; Raggi and Lee, 1963; Raggi and Pignattelli, 1975; Yachida *et al.*, 1986). According to Gelb *et al.*, (2004), the interference between IBV and NDV occurs due to the initial infection tropism that is epithelial cells of respiratory tract in which the viruses replicate in the cytoplasm. Studies carried out by Montgomery *et al.* (1997) in which many vaccinations to AIB and ND were administered, combined or not, verified that IBV induced a decrease in the capacity of immune response of gland of Harder (GH). The reduced capacity of response can decrease the level of antibody response to NDV when the birds are vaccinated to IBV and NDV in the same time. Dohms *et al.* (1988) verified a similar situation to IBDV, showing that there is a decrease of plasma cell

quantity in the gland of Harder during infectious bursal disease virus infection of 3-week-old broiler chickens that might induce deficiency of local immunity in the paraocular region and upper respiratory tract associated with IBD.

However the groups that did not receive only the vaccination to ND and AIB (T1, T2, T3 groups) but also additional IBD vaccinations did not present reduced NDV antibodies as the T0 group. There was a progressive increase of NDV antibody titers according to the addition of IBD vaccinations, that were one, two or three, respectively T1, T2 and T3, with the NDV antibody response to T2 and T3 being statistically superior to T1. These results show a possible positive interference that the use of live vaccine of IBD promotes on NDV antibody response. The progressive increase of antibody titers of T1, T2, and T3 groups in comparison to T0 was not enough to be superior to the antibody response of the birds that received only ND vaccination (CG-ND). These findings suggest that polyvalent vaccination with IBDV does not improve the NDV response, but decrease, modify or interfere with interference of IBV under NDV. This way the live IBDV vaccine induces an NDV antibody response superior to the birds that receive only ND and AIB vaccinations, without IBDV vaccine.

Regarding the IBV and IBDV antibody titers, it wasn't observed significant ($p < 0.05$) variation among the groups. That indicates the low interference of NDV under IBV, as observed by Raggi; Lee, (1964) e Zygraich *et al.*, (1973). In the same way, it wasn't observed interference of IBV and NDV under IBDV, because IBDV antibodies were similar independent of the group or experimental vaccination program administered.

CONCLUSION

We can conclude that the utilization of polyvalent vaccinal programs have a different efficacy comparing to monovalent vaccinations when Newcastle disease, avian infectious bronchitis, and infectious bursal disease vaccinations are applied. This way, the use of vaccinations to infectious bursal disease in polyvalent vaccinal programs is desirable due to improvement of NDV response, in spite of the presence of IBV by the probable reduction of interference of IBV under NDV.

REFERENCES

Alam J, Rahman MM, Sil BK, Khan MSR, Sarker MSK. Effect of maternally derived antibody on vaccination against infectious bursal



disease (Gumboro) with live vaccine in broiler. *International Journal of Poultry Science* 2002; 1(4):98-101.

Al-Natour MQ, Ward LA, Saif YM, Stewart-Brown B, Keck LD. Effect of different levels of maternally derived antibodies on protection against infectious bursal disease virus. *Avian Diseases* 2004; 48:177-182.

Ashraf S, Abdel-Alim G, Al-Natour MQ, Saif YM. Interference between mild and pathogenic strains of infectious bursal disease virus in chickens. *Avian Diseases* 2005; 49:99-103.

Beard CW. Infectious bronchitis virus interference with Newcastle disease virus in monolayers of chicken kidney cells. *Avian Diseases* 1967; 11:399-406.

Bracewell CD, Dawson PS, Allan WH. Antibody responses to a live Newcastle disease vaccine when combined with a live infectious bronchitis vaccine. *Veterinary Record* 1972; 90:248-249.

Cao YC, Yeng WS, Law M, Bi YZ, Leung FC, Lim BL. Molecular characterization of seven Chinese isolates of infectious bursal disease virus: classical, very virulent, and variant strains. *Avian Diseases* 1998; 42:340-351.

Cardoso WM, Aguiar Filho JLC, Romão JM, Oliveira WF, Salles RPR, Teixeira RSC, Sobral MHNR. Effect of associated vaccines on the interference between Newcastle Disease virus and Infectious Bronchitis virus in broilers. *Brazilian Journal of Poultry Science* 2005; 7:181-184.

Cook JKA, Huggins MB, Orbell SJ, Mawditt K, Cavanagh D. Infectious bronchitis virus vaccine interferes with the replication of avian pneumovirus vaccine in domestic fowl. *Avian Pathology* 2001; 30:233-242.

Cosgrove AS. An apparently new disease of chickens-avian nephrosis. *Avian Diseases* 1962; 6:385-389.

Dohms JE, Lee KP, Rosenberger JK, Metz AL. Plasma cell quantitation in the gland of Harder during infectious bursal disease virus infection of 3-week-old broiler chickens. *Avian Diseases* 1988; 32(4):624-31.

Fussell, LM. Poultry industry strategies for control immunosuppressive diseases. *Poultry Science* 1998; 77:1193-1196.

Gelb JJr, Jackwood MW. Infectious bronchitis. In: Swayne DE, Glisson JR, Jackwood MW, Pearson JE, Reed WM, editors. *A laboratory manual for the isolation and identification of avian pathogens*. 4th ed. Kendall: Hunt Publ.; 1988.

Gelb JJr, Campion L, Ladman B. Interferência na replicação entre os vírus da bronquite infecciosa e da doença de Newcastle. In: *Anais da Conferência Apinco 2004 de Ciência e Tecnologias Avícolas*; 2004; Santos. São Paulo-Brasil. p. 63-70.

Jackwood DH, Saif YM. Antigenic diversity of infectious bursal disease viruses. *Avian Diseases* 1987; 31:766-770.

Lasher HN, Shane SM. Infectious bursal disease. *World's Poultry Science Journal* 1994; 50:133-166.

Lutticken D. Viral diseases of the immune system and strategies to control infectious bursal disease by vaccination. *Acta Veterinaria Hungarica* 1997; 45:239-249.

Kibenge FSB, Dhillon AS, Russel RJ. Biochemistry and immunology of infectious bursal disease virus. *Journal of General Virology* 1988; 69:1757-1775.

Knezevic N, Sekler M, Veljovic IJ, Kozlina B, Rodic J. First experiences with poulvac R bursine-2 vaccine against Gumboro disease. In: *Proceeding of the 8^o Yugoslav Symposium on Poultry Production*; 1999; Sokobanja. Yugoslavia.

Knoblich HV, Sommer SE, Jackwood DJ. Antibody titers to infectious bursal disease virus in broiler chicks after vaccination at one day of age with infectious bursal disease virus and Marek's disease virus. *Avian Diseases* 2000; 44:874-884.

Montgomery YRD, Maslin WR, Boyle CR. Effects of Newcastle disease vaccines and Newcastle disease/infectious bronchitis combination vaccines on the head-associated lymphoid tissues of the chicken. *Avian Diseases* 1997; 41(2):399-406.

Moraes HLS, Salle CTP, Nascimento VP, Rocha ACGP, Souza GF, Furian TQ, Artencio JO. Infectious bursal disease: evaluation of maternal immunity and protection by vaccination of one-day old chicks against challenge with a very virulent virus isolate. *Revista Brasileira de Ciência Avícola* 2005; 7(1):51-57.

Murphy FA, Fauquet CM, Bishop DHL, Ghabrial SA, Jarvis AW, Martelli GP, Mayo MA, Summers MD, editors. *Virus taxonomy*. Vienna & New York: Springer-Verlag; 1995. (Report, 6)

Müller H, Scholtissek C, Becht H. The genome of infectious bursal disease virus consists of two segments of double-stranded RNA. *Journal of Virology* 1979; 31:584-589.

Raggi LG, Lee GG. Infectious bronchitis virus interference with growth of Newcastle disease virus. I. Study of interference in chicken embryos. *Avian Diseases* 1963; 7:106-122.

Raggi LG, Lee GG. Infectious bronchitis virus interference with growth of Newcastle disease virus. II. Interference in chickens. *Avian Diseases* 1964; 8:471-480.

Raggi LG, Pignattelli P. Identification of infectious bronchitis virus by interference with the B-1 isolant of Newcastle disease virus. Waxing and waning of interference. *Avian Disease* 1975; 19:334-342.

Sabin AB. Present position of immunization against poliomyelitis with live virus vaccine. *British Medical Journal* 1959; 1:633-680.

Sharma JM, Kim IJ, Rautenschlein S, Yeh HY. Infectious bursal disease virus of chicken: pathogenesis and immunosuppression. *Developmental & Comparative Immunology* 2000; 24:223-235.

Thornton DH, Muskett JC. Effect of infectious bronchitis vaccination on the performance of live Newcastle disease vaccine. *Veterinary Record* 1975; 96:467-468.

USER'S guide: statistic and graphics guide. Ver. 3.1. Cary, NC: SAS Institute; 1995.



Yachida S, Kuwahara E, Iritani Y, Hayashi Y. In egg interference of embryo non-lethal avian infectious bronchitis viruses (IBV) with velogenic Newcastle disease virus and embryo adapted IBV. Research Veterinary Science 1986; 40:1-3.

Van den Berg TP. Acute infectious bursal disease in poultry: a review. Avian Pathology 2000; 29:175-194.

Vakharia VN, Synder DB, Lutticken D, Mengel-Whereat SA, Savage PK, Edwards GH, Goodwin MA. Active and passive protection against variant and classic infectious bursal disease virus strains induced by baculovirus-expressed structural proteins. Vaccine 1994; 12:452-456.

Whitaker-Dowling P, Maassab HF, Younger JS. Dominant negative mutants as antiviral agents: simultaneous infection with the cold adapted live-virus vaccine for influenza A protects ferrets from disease produced by wild-type influenza. American Journal of Infectious Diseases 1992; 164:200-202.

Whitaker-Dowling P, Younger JS. Viral interference dominance of mutant viruses over wild type virus in mixed infections. Microbiology Review 1987; 51:179-191.

Zygraich N, Vasoboinic E, Berge E. The use of a combined vaccine against Newcastle disease and infectious bronchitis. Veterinary Record 1973; 93:516-518.