



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e Tecnologia
Avícolas
Brasil

Cordeiro, AFS; Nääs, IA; Salgado, DD

Field evaluation of broiler gait score using different sampling methods

Revista Brasileira de Ciência Avícola, vol. 11, núm. 3, julio-septiembre, 2009, pp. 149-154

Fundação APINCO de Ciência e Tecnologia Avícolas

Campinas, SP, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179714003002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Field Evaluation of Broiler Gait Score Using Different Sampling Methods

■ Author(s)

Cordeiro AFS¹
Nääs IA²
Salgado DD³

¹ M. Sc. student. Agricultural Engineering College. State University of Campinas.

² Professor. Agricultural Engineering College. State University of Campinas.

³ Ph.D. student. Agricultural Engineering College. State University of Campinas.

■ Mail Address

Irenilza Alencar Nääs

E-mail: irenilza@feagri.unicamp.br

■ Keywords

Animal welfare, environmental rearing conditions, locomotion deficiency.

ABSTRACT

Brazil is today the world's largest broiler meat exporter; however, in order to keep this position, it must comply with welfare regulations while maintaining low production costs. Locomotion problems restrain bird movements, limiting their access to drinking and feeding equipment, and therefore their survival and productivity. The objective of this study was to evaluate locomotion deficiency in broiler chickens reared under stressful temperature conditions using three different sampling methods of birds from three different ages. The experiment consisted in determining the gait score of 28, 35, 42 and 49-day-old broilers using three different known gait scoring methods: M1, birds were randomly selected, enclosed in a circle, and then stimulated to walk out of the circle; M2, ten birds were randomly selected and gait scored; and M3, birds were randomly selected, enclosed in a circle, and then observed while walking away from the circle without stimulus to walking. Environmental temperature, relative humidity, and light intensity inside the poultry houses were recorded. No evidence of interaction between scoring method and age was found however, both method and age influenced gait score. Gait score was found to be lower at 28 days of age. The evaluation using the ten randomly selected birds within the house was the method that presented the less reliable results. Gait score results when birds were stimulated to walk were lower than when they were not simulated, independently of age. The gait scores obtained with the three tested methods and ages were higher than those considered acceptable. The highest frequency of normal gait score (0) represented 50% of the flock. These results may be related to heat stress during rearing. Average gait score increased with average ambient temperature, relative humidity, and light intensity. The evaluation of gait score to detect locomotion problems of broilers under rearing conditions seems subjective and difficult to be properly performed.

INTRODUCTION

Lame broilers cannot walk freely and are not able to reach either the feeder or the drinker when hungry or thirsty, limiting their survival and productivity. The gait score system is an estimation of locomotion deficiency and it is based on the visual judgment of the ability a broiler chicken has to walk on a known surface. The rearing environment is considered to be one of the main aspects related either to the success or failure of broiler production. According to Knowles *et al.* (2008) modern management techniques associated to the genetic characteristics of rapidly-growing broilers have compromised their welfare, as well as their walking ability. Bessei (2006) points out that the light intensity



the incidence of locomotion abnormalities. Literature studies report that broiler chicken lameness problems are related to genetic strains (Bokkers & Koene, 2004; Bizeray *et al.*, 2000), to bird age, and to flock density (Sorensen *et al.*, 2000), in addition to high weight gain rate (Bizeray *et al.*, 2000).

The gait score system developed by Kestin *et al.* (1992) has been used to assess lameness in broilers, and the methodology consists of empirical repetitive visual observations of how birds walk on a surface. The system is divided into six levels of observed leg weakness, as follows: 0 (sound bird); 1 (the bird moves fast, but a slight walking deficiency is observed); 2 (the bird moves fast, but there is significant walking deficiency); 3 (the bird moves fast, but it presents an important deficiency); 4 (the bird moves with serious difficulty); and 5 (the bird barely moves and often uses the wings for crawling).

Garner *et al.* (2002) modified the previous system, simplifying the rules by removing the intermediate scores as they are difficult to estimate. The new system uses four levels of walking ability: 0 (sound bird that walks ten steps normally); 1 (average bird that walks ten steps with some degree of difficulty and presents some unbalance during walking); and 2 (lame bird that walks with great difficulty and sits after 1 to 4 steps) (Dawkins *et al.*, 2004). This is also an individual and subjective evaluation, and therefore it is difficult to compare these two systems as well as the scores given by two different observers.

Grandin (2007) recommends that, in commercial farms, gait score is assessed by randomly selecting around 100 broilers in two different areas of the house, and trying to observe the birds while they move away from that specific place towards other areas. High welfare standard is when 95% - 99% of the broilers present gait score below 1, and welfare is considered acceptable when 70% of the birds walk soundly. Dawkins *et al.* (2004) suggest that one single bird, randomly chosen in the house, needs to be observed and analyzed in ten distinct places.

This study aimed at evaluating lameness in broiler chickens reared in poultry houses, with open sides and natural ventilation and limited environmental control resources, using a simplified system of gait scoring. An association among welfare, observed lameness, and three sampling of gait score evaluation system were studied.

MATERIALS AND METHODS

The experiment was carried out from September to November 2008, in three commercial integrated broiler chicken farms located in the counties of Itatiba and Louveira, in the state of São Paulo, Brazil. The climate of the region is type Cwa, according to Köppen classification. Average regional temperature varied between 18 and 31°C.

Visual observation of the gait score was carried out with 28, 35, and 42-day-old broilers randomly selected from the flocks of the selected houses. Gait score was estimated using a three-scale scoring system (Dawkins *et al.*, 2004), by a skilled observer.

Bird sampling method

Three different sampling methods were used to assess gait score:

- Method 1 (M1) - With walking stimulus: A circle was built using 2-mm thick plywood boards (the same used for building the brooding circles), enclosing 100 randomly-selected birds. Broilers were stimulated to walk away from the circle by a person walking slowly behind them. Half of the total numbers of birds (50%) were gait scored.
- Method 2 (M2) - Ten birds were randomly selected in different areas inside the house, and were gait scored using the system proposed by Dawkins *et al.* (2004).
- Method 3 (M3) - No walking stimulus: One hundred randomly-selected birds were enclosed inside the house, and they were gait the previously described circle, and were allowed to walk freely outside the circle. Half of the birds (50%) were gait scored (Grandin, 2007).

Environmental parameters

The following broiler house environmental parameters were measured during the tests: dry bulb temperature (T, °C), air relative humidity (RH, %) and air velocity (AV, m s⁻¹) using the thermal hydro-anemometer HTA®; light intensity (L, lx), using a digital luximeter ISBB®260; and ammonia concentration (A, ppm) using the manual pump Accuro Dräger® and respective reagents. The parameters were recorded at three different points: geometric center, and both ends of the houses. At each location three measurements were made in the morning, starting at 10:00 AM, and three in the afternoon, starting at 02:00 PM, with intervals of 15



Data analysis

The collected data were submitted to analysis of variance at 95% significance level. Correlations between environmental parameters and gait score were tested using Pearson's linear correlation.

The hypothetical broiler welfare distribution was determined based on Grandin's (2007) concept, which establishes that welfare standard is high when 95 to 99% of all observed broilers present gait scores between 0 and 1 in the six-level scale. Assuming that gait scores 0 and 1 in the six-level scale correspond to the gait score 0 in the three-scale system, it was estimated that the welfare standard was high when 95% of the birds present gait score 0. The remaining 5% would be distributed between gait scores 1 and 2. The Chi Square test was used to estimate welfare, and the observed gait score distribution was compared to the hypothetical distribution.

RESULTS AND DISCUSSION

Table 1 shows the ANOVA results of the effects of the factors Method and Age on gait score; in this case M1, M2, and M3 were taken into account. It was not possible to test the full interaction as M2 was only evaluated in 35 and 42-day-old birds. However, the available data showed significant influence of sampling method (Method) and Age on gait score results ($p < 0.05$).

Table 1. ANOVA results of the effects of sampling Method and Age on gait score.

| Source of variation | Degree of freedom | Sum of Squares | Mean Square | F | P value |
|---------------------|-------------------|----------------|-------------|------|---------|
| Method | 2 | 4.0842 | 2.0998 | 4.55 | 0.011 |
| Age | 2 | 6.0676 | 3.0338 | 6.58 | 0.002 |
| Error | 315 | 145.2358 | 0.4611 | | |
| Total | 319 | 155.3875 | | | |

M2 was removed from the analysis in order to test the interaction between factors. The factor Method (M1 and M3) significantly influenced gait score results ($p < 0.05$, Table 2).

Table 2. ANOVA results of the effect of sampling Method, Age, and Method and Age interaction on gait score.

| Source of variation | Degree of freedom | Sum of Squares | Mean Square | F | P value |
|---------------------|-------------------|----------------|-------------|------|---------|
| Method | 1 | 4.0833 | 4.0833 | 9.06 | 0.003 |
| Age | 2 | 6.1067 | 3.0533 | 6.78 | 0.001 |
| Method * Age | 2 | 0.1867 | 0.0933 | 0.21 | 0.813 |

No evidence of interaction between the factors of Method and Age was found. This showed that sampling Method influenced gait score independently from Age, and likewise, Age influenced gait score independently of the process used for sampling the birds. This means that both how birds are chosen and timing of gait score evaluation in the farm may affect the results obtained.

Figure 1 shows the confidence interval of mean gait score as a function of the sampling Method. It is observed that M2 has a high standard error around the mean, thereby indicating lower reliability of the results. In Figure 2, it is observed that M2 results behaved differently from the other methods (M1 and M3), also indicating that this sampling methods is not as reliable as the other two. The standard error of the mean value in M1 and M3 were relatively lower than M2, showing that the results obtained using methods 1 and 3 to sample birds are more reliable and accurate.




Figure 1. Interval plot graph showing the influence of sampling Method on gait score results.




Figure 2. Distribution of the three levels of gait score according to the three methods used for bird sampling.

Figure 3 shows that, independently from sampling method, gait score was low at 28 days of age



of age. These results are partially consistent with the findings of Sorensen *et al.* (2000) who observed that limb disorders are a relatively reduced problem in young broilers. When using M1 gait score results, there was less lameness than when M3 was used, probably due to the fact that the birds walk more than when using M3. Bokkers *et al.* (2006) found that broilers can be motivated to walk long distances to reach feed. According to Bizeray *et al.* (2000), locomotion activities in broilers during the last week of life may increase if birds are stimulated to move during the first days of life, contributing to reduce lameness.




Figure 3. Interval plot showing the influence of age within two sampling methods on field gait score.

During the field experiment it was observed that, without the stimulus to induce birds to walk they barely gave a few steps and then rested influencing final gait score results. When stimulated they walked more and gait score results were better. Broilers do not walk often inside the poultry house, making it difficult to evaluate final gait score. It is not clear if not walking is related to pain or simply to the low frequency of this activity. On the other hand, when stimulated to walk, broilers make an extra effort to move, and move either faster or more than they usually do, which may interfere in the interpretation of the final gait score results. Weeks *et al.* (2000) found that sound birds spent 76% of their time lying, whereas lame birds spent 86%. The authors observed that lying behavior also increased with age. Similar results were obtained by Webster *et al.* (2008). In the present experiment, it was observed that some broilers reached the feeder or the drinker after few steps, or stopped walking when meeting another bird in their way. These birds received gait score 2; however, it cannot be stated that this meant they presented a

was already tired of walking before the evaluation started, some degree of error may have been introduced in the gait score evaluation when M2 was used, as when birds are randomly selected, it is not known whether they are already tired or not, which may influence their walking behavior independently from its locomotory soundness. According to Webster *et al.* (2008) broilers may ignore some degree of pain either when facing something new or experiencing fear, which may also interfere in the final results. When M1 method was used 2% of the birds presented gait score 2 at 28 days of age, while this proportion was 20% when using M3 sampling (Table 3). Sorensen *et al.* (2000) found that less than 1% of 28-day-old broilers presented gait score 4 or 5. These results are higher than those obtained in the present experiment considering that gait score values of 4 and 5 correspond to score 2 in the three-points gait score scale.

Using M1 in a field study that applied the six-point gait score scale, Sanotra *et al.* (2001) and Kestin *et al.* (1992) found that 25% and 10% of birds of all ages presented gait score 0, respectively, while Brickett *et al.* (2007) found 62.5%. Although in the present study gait score was evaluated using the three-point gait score scale, gait score 0 results were higher than those found by Sanotra *et al.* (2001) and Kestin *et al.* (1992), and lower than those of Brickett *et al.* (2007), which results were higher than those obtained in the present experiment, independently from age or sampling method. Using the three-point broiler gait score scale in field, Dawkins *et al.* (2004) found that 61.1 to 80.8% of the birds presented gait score 0, which is also higher than those obtained in the present study (Table 3).

Table 3. Distribution of the frequency of gait score by age, using the three Methods (M1, M2, and M3).

| Gait Score | M1 | | | M2 | | | M3 | | |
|------------|------|------|------|------|------|----|------|------|------|
| | 42 | 35 | 28 | 42 | 35 | 28 | 42 | 35 | 28 |
| 0 | 0.42 | 0.26 | 0.46 | 0.50 | 0.30 | - | 0.28 | 0.16 | 0.34 |
| 1 | 0.44 | 0.54 | 0.52 | 0.20 | 0.50 | - | 0.54 | 0.52 | 0.46 |
| 2 | 0.14 | 0.20 | 0.02 | 0.30 | 0.20 | - | 0.18 | 0.32 | 0.20 |

Using Grandin's (2007) hypothetical gait score distribution as related to broiler welfare, a statistical test was performed to compare the frequency distribution of estimated *versus* observed gait score. The results suggest that locomotory problems indicate a lack of welfare in the evaluated flocks (Tables 4 and 5, and Figure 3). The probable cause of these results is the


**Table 4.** Chi-square statistical test for adjusting frequency distribution.

| Gait score | Number of Observed Gait score | Proportion | Number of Estimated Gait score | Proportion | N | DF | ÷2 | P value |
|------------|----------------------------------|------------|-----------------------------------|------------|-----|----|---------|---------|
| 0 | 104 | 0.950 | 304 | 131.58 | 320 | 2 | 3256.58 | 0.000 |
| 1 | 158 | 0.025 | 8 | 2812.50 | | | | |
| 2 | 58 | 0.025 | 8 | 312.50 | | | | |

Table 5. Mean and standard deviation of recorded environmental parameter according to age at gait score evaluation.

| Age (days) | A (ppm) | | T (°C) | | RH (%) | | AV (m/s) | | LI (lx) | |
|------------|---------|------|--------|------|--------|------|----------|------|---------|--------|
| | Mean | SD | Mean | SD | Mean | SD | Mean | SD | Mean | SD |
| 28 | - | - | 28.9 | 0.97 | 49.4 | 2.92 | 0.83 | 0.64 | 188.3 | 73.07 |
| 35 | 11.0 | 1.87 | 31.0 | 1.00 | 60.2 | 4.45 | 1.76 | 0.62 | 331.5 | 159.06 |
| 42 | 11.0 | 2.85 | 26.8 | 2.28 | 67.2 | 6.21 | 0.45 | 0.45 | 205.3 | 205.19 |

experiment, gait score increased as average environmental temperature, relative humidity, air speed, and light intensity increased. The only exceptions were the relative humidity values when the M2 sampling method was applied (Table 6).

**Figure 4.** Comparative distribution of observed and estimated gait score and flock welfare condition.**Table 6.** Correlation between environmental parameters and mean gait score values according to sampling method.

| Ambient variables | M1 | M2 | M3 |
|-------------------|------|-------|------|
| RH (%) | 0.53 | -1.00 | 0.24 |
| AV (m/s) | 0.75 | 1.00 | 0.92 |
| L (lx) | 0.95 | 1.00 | 1.00 |
| T (°C) | 0.53 | 1.00 | 0.81 |

According to Dawkins *et al.* (2003), the percentage of broilers presenting poor performance during growth is positively correlated both with high environmental temperature and relative humidity, particularly between three and five weeks of age. The number of sound broilers (gait score=0) was negatively correlated with the temperature, which had the highest correlation

values. It is reported in literature that locomotory problems are related to harsh rearing environments. Bruno *et al.* (2007) found that broilers reared under heat stress presented reduced length and width of the tibia, femur and humerus. Light exposure is also another variable that is reported to influence broiler gait score results. Lin *et al.* (2006) determined that broiler locomotion may be affected when light management is beyond a 6 - 10 lx range. Bird performance is usually affected by the simultaneous occurrence of extreme values in factors related to the thermal and aerial broiler rearing environment (Martrenchar *et al.*, 1997; Dawkins, 2003; Moura *et al.*, 2006; Bessei, 2006).

CONCLUSIONS

It is difficult to estimate broiler gait score under field conditions as all evaluation methods are subjective. When comparing three distinct sampling methods, the use of ten broilers randomly chosen inside the house was shown to be the least reliable method.

The method adopted to estimate gait score influenced the final results.

High gait score values were related to the harsh environmental conditions recorded inside the broiler house.

REFERENCES

- Bessei W. Welfare of broilers. *World's Poultry Science Journal* 2006; 62(3):455-66.
- Bizeray D, Laterrier C, Constantin P, Picard M, Faure JM. Early locomotor behaviour in genetic stocks of chickens with different growth rates. *Applied Animal Behaviour Science* 2000; 68:231-242.
- Bokkers EAM, Koene P. Motivation and ability to walk for a food



Bokkers EAM, Zimmerman PH, Rodenburg TB, Koene P. Walking behaviour of heavy and light broilers in an operant runway test with varying durations of feed deprivation and feed access. *Applied Animal Behaviour Science* 2006; 108:129-142.

Weeks CA, Danbury TD, Davies HC, Hun TP, Kestin SC. The behaviour of broiler chickens and its modification by lameness. *Applied Animal Behaviour Science* 2000; 67(1/2):111-125.

Brickett KE, Dahiya JP, Classen HL, Gomis S. Influence of dietary nutrient density, feed form, and lighting on growth and meat yield of broiler chickens. *Poultry Science* 2007; 86(10):2172-2181.

Bruno LDG, Luquetti BC, Furlan RL, Macari M. Influence of early qualitative feed restriction and environmental temperature on long bone development of broiler chickens. *Journal of Thermal Biology* 2007; 32:349-354.

Dawkins MS. Behavior as a tool in the assessment of animal welfare. *Zoology* 2003; 106(4):383-7.

Dawkins MS, Donnelly CA, Jones TA. Chicken welfare is influenced more by housing conditions than by stocking density. *Nature* 2004; 427:342-344.

Garner JP, Falcone C, Wakenell P, Martin M, Mench JA. Reliability and validity of a modified gait scoring system and its use in assessing tibial dyschondroplasia in broilers. *British Poultry Science* 2002; 43(3):355-363.

Grandin, T. Poultry slaughter plant and farm audit: critical control points for bird welfare (2007) [cited 2008 jan]. Available from: <http://www.grandin.com/poultry.audit.html>.

Kestin SC, Knowles TG, Tinch AE, Gregory NG. Prevalence of leg weakness in broiler chickens and its relationship with genotype. *Veterinary Record* 1992; 131:190-194.

Knowles TG, Kestin SC, Haslam SM, Brown SM, Green LE. Leg disorders in broiler chickens: Prevalence, risk factors and prevention [cited 2008 jan]. Available from: <http://www.plosone.org/article/info:doi/10.1371/journal.pone.0001545>.

Lin H, Jiao HC, Buyse J, Decuypere E. Strategies for preventing heat stress in poultry. *World's Poultry Journal* 2006; 62(1):71-85.

Martrenchar A, Morisse JP, Huonnic D, Cotte JP. Influence of stocking density on some behavioural, physiological and productivity traits broiler. *Veterinary Research* 1997; 28(5):473-80.

Moura DJ, Nääs IA, Pereira DF, Silva RBTR, Camargo GA. Animal welfare concepts and strategy for poultry production: a review. *Revista Brasileira de Ciência Avícola* 2006; 8(1):137-48.

Sanotra GS, Lund JD, Ersboll JS, Vestergaard KS. Monitoring leg problems in broilers: A survey of commercial broiler production in Denmark. *World's Poultry Science* 2001; 57:55-59.

Sorensen P, Su G, Kestin SC. Effects of age and stocking density on leg weakness in broiler chickens. *Poultry Science* 2000; 79(6): 864-870.

Webster AB, Fairchild BD, Cummings TS, Stayer PA. Validation of a three point gait scoring system for field assessment of walking ability of broiler chickens. *Applied Animal Behaviour Science* 2006; 108:143-150.