



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e Tecnologia
Avícolas
Brasil

Raghiante, F; Rocha, TS; Rossi, DA; Silva, PL
Penetration time of Salmonella Heidelberg through shells of white and brown commercial eggs
Revista Brasileira de Ciência Avícola, vol. 12, núm. 4, outubro-diciembre, 2010, pp. 273-277
Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, SP, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179715864009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Penetration Time of Salmonella Heidelberg Through Shells of White and Brown Commercial Eggs

■ Author(s)

Raghianti F¹
Rocha TS^{2*}
Rossi DA³
Silva PL³

- ¹ DVM, Universidade Federal de Uberlândia, MG.
- ² Pos Graduate Student in Avian Pathology, Universidade Estadual Paulista, Campus Botucatu, SP.
- ³ Professors at the School of Veterinary Medicine, Universidade Federal de Uberlândia, MG.

■ Mail Address

TS Rocha
Universidade Estadual Paulista Júlio de Mesquita Filho, Campus Botucatu.
Hospital Veterinário s/ nº
Prédio da Patologia Veterinária
Laboratório de Ornitopatologia
18.618-970. Distrito de Rubião Júnior. Botucatu, SP, Brazil
Phone: 55 14 38116293
Fax: +55 14 38116293

E-mail: ticirocha@yahoo.com.br

■ Keywords

Eggs, penetration time, Salmonella Heidelberg.

ABSTRACT

This study aimed at determining the minimum time required for the penetration of Salmonella Heidelberg inside the eggs after contact with contaminated material. Recently-collected brown and white eggs from laying hens between 45-50 weeks of age, reared in a commercial poultry house, were artificially contaminated by contact with wood shavings moistened with liquid inoculum of Salmonella Heidelberg in stationary-growth phase (10^3 - 10^4 CFU g⁻¹). According to type (white or brown), eggs were distributed into three different groups, with four replicates each: negative control group (no artificial contamination), positive control group (analyzed externally immediately after contamination and internally after the maximum storage period of the test group) and test group. Eggs were stored at controlled environmental temperature varying from 25°C to 30°C. In the test group, eggs contents (yolk and albumen) were pooled and analyzed after 1:00, 1:30, 2:00, 2:30, 3:00, 3:30, and 4:00 hours after contamination for the presence of Salmonella Heidelberg in 25g of this pool. The experimental unit consisted of five eggs in each test. The analysis protocol included pre-enrichment, selective enrichment, plating on selective agar, and biochemical and serological tests. The results obtained were submitted to logistic regression, which indicated that the presence of Salmonella Heidelberg was verified after 2:16 h and 2:44 h of contact with white and brown eggs, respectively.

INTRODUCTION

The egg is probably the food item most frequently involved in outbreaks of foodborne infections with Salmonella spp as etiological agent (Gantois et al., 2009; Gast et al., 2005). Eggs are commonly contaminated by Salmonella through the shell. Carrier birds shed the pathogen in the feces, where it may persist for long periods in empty poultry houses, highlighting the importance of the environment in horizontal transmission (Silva & Duarte, 2002).

Salmonella Enteritidis is the serovar most commonly involved in human disease; however, other serovars, such as S. enterica serovar Heidelberg (S. Heidelberg), are becoming increasingly frequent. (Mammìna et al., 2003). Salmonella Heidelberg has been detected in Brazilian poultry stocks since 1982, characterizing the importance of the emergence of this serotype (Hofer et al., 1997). The Centers for Disease Control and Prevention have implicated Salmonella Heidelberg as a potentially significant source of egg-associated human disease (Gast et al., 2005).

Many studies report that Salmonella is capable of penetrating the eggshell and replicating inside the egg (Gast & Holt, 2000; Gast et al., 2005). Several factors are involved in the time required for its penetration in the egg, including eggshell, albumen, and yolk quality (Messens et al., 2004); bird age (Lapão et al., 2005); feed type and physical form



(Pappas et al., 2005); egg storage time (Chen et al., 2005); genetics (Dunn et al., 2005); and photoperiod (Backhouse et al., 2005).

The presence of *Salmonella* in the egg yolk may result from the migration of bacteria from the contaminated shell. It is known that the eggshell pores allow the penetration of *Salmonella*, and its penetration index is related to storage time and temperature rather than to the number of pores (Kanashiro et al., 2002).

The presence of *Salmonella* spp in poultry flocks is constant and requires the poultry industry adopt continuous control measures during all stages of poultry production (Rodrigues, 2005). Maintaining flocks in good health status and cleaning and disinfecting eggs with chemical products after lay are the most frequent practices applied to prevent or to reduce contamination and bacterial replication in eggs (Hammack et al., 1993).

In addition to the use of effective antimicrobial products, efficient egg disinfection requires knowing precisely when microorganisms cross the eggshell in order to apply any treatments before this happens (Hammack et al., 1993; Board & Tranter, 1995).

Considering the health and economic importance of *Salmonella* and aiming at providing the poultry industry useful data on the time limit treatments must be performed in order to prevent internal contamination of commercial eggs, the objective of the present study was to determine the penetration time of *Salmonella* Heidelberg in artificially-contaminated commercial white and brown eggs.

MATERIAL AND METHODS

The study used 280 commercial white eggs and 280 commercial brown eggs of 45 to 40-week-old layers housed in a commercial poultry house. Layers were free from *Salmonella* spp., as tested before the beginning of this experiment. Eggs were collected immediately after lay (first morning collection) and were not washed or submitted to any cleaning process.

The serovar *Salmonella* Heidelberg used was isolated from birds. The isolated sample was cultivated in Brain Heart Infusion Broth, and the inoculum with a known concentration (10^3 - 10^4 CFU g⁻¹) was prepared when the culture presented stationary growth phase.

The experimental contamination was performed by placing sterile wood shavings in bags, moistening with distilled water, sprayin with liquid inoculum in quantity sufficient to achieve 10^3 to 10^4 CFU g⁻¹ of *Salmonella* Heidelberg per gram, and placing the eggs inside the bag. Eggs remained in contact with contaminated

shavings for 10 minutes at 25°C (\pm 5°C).

The study was carried out in two different phases (pilot test and determination phase) for each egg type (brown and white). During both phases, eggs were distributed in three experimental groups with four replicates each: Group 1: negative control - no inoculation, analyzed at the end of the storage period; Group 2, positive control - inoculated, analyzed immediately after experimental contamination, at hour zero; and Group 3: test group - eggs previously contaminated by contact with wood shavings containing *Salmonella* Heidelberg and stored at 25°C (\pm 5°C) for predetermined periods.

The pilot tests determined the shortest period in which *Salmonella* could be detected inside the egg. Eggs were tested 2, 4, 8, 10, and 24h after contact with wood shavings containing the inoculum. After the time of penetration of bacteria through the eggshell was determined, the verification phase started, analyzing internal egg contamination 1:00, 1:30, 2:00, 2:30, 3:00, 3:30, and 4:00h after contamination.

Each sample consisted of five eggs. For microbiological analysis, the eggshells were cleaned with alcohol 70°GL, aseptically broken in a laminar flow hood, their content poured into a sterile beaker, and homogenized with a sterile glass stick to form a yolk and albumen pool. Samples were analyzed using conventional culture methods according to the protocol recommended by the USDA/FSIS in 2004.

Data were analyzed by simple logistic regression analysis. The dependent variable was the presence or absence of *Salmonella* Heidelberg in egg contents, while the independent variable was evaluation time after contamination. SISVAR software program (Ferreira, 2000) was used.

RESULTS AND DISCUSSION

The contamination of wood shavings was standardized in 10^3 to 10^4 CFU/g. Average *Salmonella* counts in the eggshell determined immediately after experimental contamination in the first and second experimental phases were 8×10^4 CFU/g¹ and 6×10^4 CFU/g, respectively.

In the first phase, the presence or absence of *Salmonella* Heidelberg in the egg content after inoculation and storage at 22.5°C (\pm 2°C) was evaluated 2, 4, 8, 10, and 24h in four replicates. The negative control and positive control groups presented negative and positive counts, respectively, in all replicates.

The results demonstrate that *Salmonella* Heidelberg was detected in the egg contents four hours after



contact. These results are similar to those obtained by Schoeni et al. (1995) and Oliveira & Silva (2000), who observed that, after external contamination of eggs, different *Salmonella* serovars were detected inside the egg before 24 hours of storage. On the other hand, Messens et al. (2005) observed in their study on the penetration time of *Salmonella* Enteritidis through the eggshell that, in average, 38.7% of the eggs tested had their contents contaminated by *Salmonella* and that penetration occurred on the third day after contamination.

In the second phase of this study, the experimental eggs were analyzed at 1:00, 1:30, 2:00, 2:30, 3:00, 3:30 and 4:00 hours after contamination. The results showed that internal contamination occurred after 2:00 hours, but before 2:30 hours in white eggs, whereas in brown eggs, between 2:30 and 3:00 hours after their external contamination with 10^3 to 10^4 CFU/g of *Salmonella* Heidelberg. These results are consistent with the findings of Miyamoto et al. (1998), who externally contaminated eggs for 15 minutes to 3 hours after lay and stored them at 25°C and obtained significant recovery of *Salmonella* after two hours of incubation.

The time in minutes after contamination and *Salmonella* Heidelberg counts (log CFU/g) in the eggshells immediately after experimental contamination were analyzed by logistic regression to determine the precise time of egg content contamination. Figures 1 and 2 present the logistic regression results of brown and white eggs, respectively.

Data analysis showed that white and brown eggs externally exposed to 10^3 - 10^4 CFU/g of *Salmonella* Heidelberg were internally contaminated 2:16 and 2:44 hours after contact, respectively.

Egg quality includes several aspects related to eggshell (external quality) and to the albumen and yolk (internal quality). It has a genetic basis and parameter values are different among bird strains (Silversides et al., 2006). Several investigators compared eggs of white and brown egg layer strains (Scott & Silversides, 2000; Jones et al., 2010), but there are no studies investigating if these two egg types present different contamination patterns.

Singh et al. (2009), in a study comparing eggs of different layers strains reared in two different housing systems (cages or floor pens), found higher *E. coli* contamination in brown than in white eggs, but this may be explained by the fact that brown layers were reared in the floor-pen system and did not use the nests properly, laying half of their eggs on the floor.




Figure 1 - Logistic regression of *Salmonella* Heidelberg penetration time in artificially inoculated white commercial eggs. Values in the Y axis indicate the absence (-0.1) to the presence (1.1) of the bacterium in the egg content.




Figure 2 - Logistic regression of *Salmonella* Heidelberg penetration time in artificially inoculated brown commercial eggs. Values in the Y axis indicate the absence (-0.1) to the presence (1.1) of the bacterium in the egg content.

The egg has an impressive arsenal of antimicrobial protective mechanisms, including both nonspecific physical barriers and highly efficient microbiocidal molecules. However, *Salmonella* can survive and replicate on the eggshell despite the absence of fecal contamination, even at low environmental temperatures and low relative humidity (Messens et al., 2006). *Salmonella* probably survives longer at low temperatures due to the slower metabolism induced by the unfavorable conditions on the dry eggshell surface (Radkowski, 2002).

The penetration of bacteria through the eggshell is directly influenced by shell quality, as well as egg storage time and temperature (de Reu, 2006; Schoeni et al., 1995). According to Miyamoto et al. (1998), the longer the time and the higher the temperature of egg storage, the faster the penetration of bacteria through the eggshell.

Salmonella spp may also contaminate egg through the ovary. In this case, the bacteria are located in the yolk, and consequently, conventional processes of egg disinfection are useless. Therefore, adequate hygiene



measures should be a priority in layer farms. It must be stressed that no disinfection procedure per se can prevent egg contamination.

The results obtained in the present study indicate that cleaning and disinfection procedures aiming at controlling internal egg contamination should be performed before 2:16 hours in white eggs, and before 2:44 hours in brown eggs.

Ensuring layer flock health and proper disinfection of the rearing environment, facilities and eggs are important to prevent or to reduce contamination by pathogens and to ensure consumers' health.

CONCLUSION

After contaminating eggshells, *Salmonella* Heidelberg reaches the egg contents in 2:16 hours in commercial white eggs and in 2:44 hours in commercial brown eggs.

REFERENCES

- Backhouse D, Lewis PD, Gous RM. Constant photoperiods and eggshell quality in broiler breeder pullets. *Poultry Science* 2005; 46(2):211-213.
- Board RG, Tranter HS. The microbiology of eggs. In: Stadelman WJ, Cotterill OJ, editors. *Egg science and technology*. London 1995; 81-104.
- Brasil, Ministério da Agricultura e Abastecimento. Instrução Normativa nº 62, de 26 de Agosto; 2003. Seção 1. p. 14.
- Chen J, Thesmar HS, Kerr WL. Outgrowth of *Salmonella* and the physical property of albumen and vitelline membrane as influenced by egg storage conditions. *International Journal of Food Protection* 2005; 68(12):2553-2558.
- De Reu K, Grijspeerdt K, Messens W, Heyndrickx M, Uyttendaele M, Debevere J, Herman L. Eggshell factors influencing eggshell penetration and whole egg contamination by different bacteria, including *Salmonella* Enteritidis. *International Journal of Food Microbiology* 2006; 112:253-260.
- Dunn IC, Bain M, Edmond A, Wilson PW, Joseph N, Solomon S. et al. Heritability and genetic correlation of measurements derived from acoustic resonance frequency analysis; a novel method of determining eggshell quality in domestic hens. *Poultry Science* 2005; 46(3):280-286.
- Ferreira DF. Sistema SISVAR de análise estatística. Versão 4.6. Lavras:UFLA; 2000. Available from: <http://www.dex.ufla.br/danielff/dff02.htm>.
- Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Gast R, Humphrey TJ, Van Immerseel F. Mechanisms of egg contamination by *Salmonella* Enteritidis. *FEMS Microbiology Reviews* 2009; 33:718-738.
- Gast RK, Holt PS, Murase T. Penetration of *Salmonella* enteritidis and *Salmonella* Heidelberg into egg yolks in an in vitro contamination model. *Poultry Science* 2005; 84:621-625.
- Gast, RK, Holt PS. Deposition of phage type 4 and 13a *Salmonella* enteritidis strains in the yolk and albumen of eggs laid by experimentally infected hens. *Avian Disease*. 2000; 44:706-710.
- Hammack TS, Sherrod PS, Bruce VR. Growth of *Salmonella* Enteritidis in grade A eggs during prolonged storage. *Poultry Science* 1993; 72:373-377.
- Hofer E, Silva Filho SJ, Reis EMF. Prevalência de sorovares de *Salmonella* isolados de aves no Brasil. *Pesquisa Veterinária Brasileira* 1997; 17(2):55-62.
- Jones DR, Musgrove MT, Anderson KE, Thesmar HS. Physical quality and composition of retail shell eggs. *Poultry Science* 2010; 89: 582-587.
- Kanashiro AMI, Castro AGM, Cardoso ALSP, Tessari ENC, Jesus CAM, Ferreira E, Souza ECA. Isolamento de *Salmonella* Enteritidis em ovos comerciais, durante rastreamento de possível fonte de infecção em humanos. *Higiene Alimentar* 2002; 16(101):76-79.
- Lapão C, Gama LT, Soares MC. Effects of broiler breeder age and length of egg storage on albumen characteristics and hatchability. *Poultry Science* 2005; 78(5): 640-645.
- Mammìna C, Talini M, Pontello M, Di Noto AM, Natasi A. Clonal circulation of *Salmonella* enterica serotype Heidelberg in Italy. *Euro Surveillance* 2003; 11(8):222-225.
- Messens W, Dubocage L, Grijspeerdt K, Heyndrickx M, Herman L. Growth of *Salmonella* serovars in hens' egg albumen as affected by storage prior to inoculation. *Food Microbiology* 2004; 21:25-32.
- Messens W, Grijspeerdt K, Herman L. Eggshell characteristics and penetration by *Salmonella* enterica serovar Enteritidis through the production period of a layer flock. *British Poultry Science* 2005; 46(6):694-700.
- Messens W, Grijspeerdt K, Herman L. Eggshell penetration of hen's eggs by *Salmonella* enterica serovar Enteritidis upon various storage conditions. *British Poultry Science* 2006; 47(5):554-560.
- Miyamoto T, Horie T, Baba E, Sasai K, Fukata T, Arakawa A. *Salmonella* penetration through eggshell associated with freshness of laid and refrigeration. *Journal of Food Protection* 1998; 61(3):350-353.
- Oliveira DD, Silva EM. *Salmonella* em ovos comerciais: ocorrência, condições de armazenamento e desinfecção da casca. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 2000; 52(6):655-661.
- Pappas AC, Acamovic T, Sparks NH, Surai PF, Mcdevitt RM. Effects of supplementing broiler breeder diets with organic selenium and polyunsaturated fatty acids on egg quality during storage. *Poultry Science* 2005; 84(6):865-874.
- Radkowski M. Effect of moisture and temperature on survival of *Salmonella* Enteritidis on shell eggs. *Archiv für Geflügelkunde* 2002; 66:119-123.



Rodrigues DP. Ecologia e prevalência de *Salmonella* spp em aves e material avícola no Brasil. Anais da Conferência Apinco de Ciência e Tecnologia Avícolas; 2005; Santos, São Paulo. Brasil. p.223-237.

Sauter EA, Petersen CF. The effect of egg shell quality on penetration by various *Salmonellae*. Poultry Science 1974; 53:2159-2162.

Schoeni JL, Glass KA, Mcdermott JL, Wong ACL. Growth and penetration of *Salmonella* Enteritidis, *Salmonella* Heidelberg and *Salmonella* Typhimurium in eggs. International Journal of Food Microbiology 1995; 24:385-396.

Scott TA, Silversides FG. The effect of storage and strain of hen on egg quality. Poultry Science 2000; 79:1725-1729.

Silva EN, Duarte A. *Salmonella* Enteritidis em aves: retrospectiva no Brasil. Revista Brasileira de Ciências Avícolas 2002; 4(2):85-100.

Silversides FG, Villeneuve P. Is the Haugh unit correction for egg weight valid for eggs stored at room temperature? Poultry Science 1994; 73:50-55.

Singh R, Cheng KM, Silversides FG. Production performance and egg quality of four strains of laying hens kept in conventional cages and floor pens. Poultry Science 2009; 88:256-264.

United States Department of Agriculture/ Food Safety and Inspection Service. Isolation and identification of *Salmonella* from meat, poultry and egg products, 2004 [cited 2010 may 15]. Available from: http://www.fsis.usda.gov/PDF/MLG_4_03.pdf.