



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e Tecnologia
Avícolas
Brasil

Sadeghi, GH; Karimi, A; Padidar Jahromi, SH; Azizi, T; Daneshmand, A
Effects of cinnamon, thyme and turmeric infusions on the performance and immune response in of 1-
to 21-day-old male broilers

Revista Brasileira de Ciência Avícola, vol. 14, núm. 1, marzo, 2012, pp. 15-20

Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, SP, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179722273003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Effects of Cinnamon, Thyme and Turmeric Infusions on the Performance and Immune Response in of 1- to 21-Day-Old Male Broilers

■Author(s)

Sadeghi GH
Karimi A
Padidar Jahromi SH
Azizi T
Daneshmand A

Department of Animal Science
Faculty of Agriculture
University of Kurdistan
Kurdistan, Iran.

■Mail Address

Ghorbanali.Sadeghi
Department of Animal Science
College of Agriculture, University of Kurdistan
Kurdistan, Sanandaj, P.O. Box: 416, Iran.
Tel: +98-9183717052
Fax: +98-8716660070

E-mail: ghorbanalis@yahoo.com
gsadeghi@uok.ac.ir

■Keywords

Broiler, cinnamon, infusion, thyme, turmeric.

■Acknowledgement

The authors are grateful to Dr. Kenneth J. Gruber at North Carolina A&T State University, Greensboro, North Carolina for his critical and constructive reading of the manuscript.

ABSTRACT

This study was conducted to investigate the effects of the replacement of drinking water by herb infusions on the performance, relative weight of internal organs, hematocrit and immune response to Newcastle disease virus of broiler chickens. A total of 540 male broiler chicks (Ross 308) were divided into five groups, corresponding to four different treatments and one control group. Treatments were replicated eight times, and the control group four times. Experimental treatments included infusions (5 grams per liter) of cinnamon, thyme and turmeric in equal ratios in replacement of drinking water. Experimental period lasted 21 days and all chicks were fed with a corn-soybean based diet. Results showed that all herbs infusions caused significant ($p < 0.05$) decrease in live body weight compared with the control group at 21 days of age. Mix treatment significantly decreased relative carcass weight relative to the control group ($p < 0.05$). Herbs infusions increased the relative weight of some organs. None of herb additives affected hematocrit in comparison to control group. Cinnamon and herb mix infusion significantly improved bird immune response to the NDV vaccine in comparison to the control group and those that received only turmeric infusion. When all in-water additives were compare to each other, the birds supplemented with turmeric infusion showed the worst performance and immunity. The results of this experiment suggest that these herbs infusions did not favor the performance of broiler chickens.

INTRODUCTION

The use of additives like antibiotics as growth promoters in livestock has been banned due to concerns about their residues in animal tissues and subsequent induction of bacterial resistance. Due to this concern, researchers have evaluated alternative growth promoters such as probiotics and prebiotics, which benefits to the health of the digestive tract and its habitants have been suggested (Patterson & Burkholder, 2003). However, manufacturing these alternatives is currently costly and time-consuming and their effects have not been clearly proven. Some researchers have evaluated herbs and their essential oils. It has been shown that the dietary incorporation of herbs and their associated essential oils may provide beneficial effects on poultry performance and health due to the antimicrobial activity of their phytochemical components (Lee *et al.*, 2004). However, other studies have not found positive effects of herbs and their related essential oils. These latter findings may be related to experimental conditions, such as hygiene and dietary agents (Lee *et al.*, 2003).

Thyme (*Thymus vulgaris*), cinnamon (*Cinnamomum verum*) and turmeric (*Curcuma longa*) are herbs commonly used as spices in human food and have received considerable attention as additives in poultry



nutrition. Several studies have reported the beneficial effects of thyme in poultry nutrition (Al-Kassie, 2009; El-Ghousein & Al-Beitawi, 2009; Najafi & Torki, 2010). Canan Bölükbai & Kuddusi Erhan (2007) also showed that supplementation of layer diets with 0.1 and 0.5% thyme improved feed conversion and egg production and also reduced *E. colifecal* content. Cross *et al.* (2007) studied the effects of five herbs (10 g/kg) or their essential oils (1 g/kg) on broiler growth, digestibility, and intestinal microflora and showed that dietary thyme oil or yarrow herb had the most positive effects on broiler performance. Ocak *et al.* (2008) reported that the inclusion of 0.2% dry thyme leaves in broiler diets from 7 to 42 days of age had no significant effects on feed efficiency, growth performance, relative weight of internal organs and the relative length of the whole gut. However, the relative weight of abdominal fat was reduced.

Turmeric is a member of Zingiberaceae family and contains curcumin, demethoxycurcumin, bisdemethoxycurcumin and tetrahydrocurcuminoids as active constituents (Kiuchi *et al.*, 1993). Antioxidant, anti-inflammatory and nematocidal activities of turmeric and their relative constituents have been demonstrated (Kiuchi *et al.*, 1993; Ammon *et al.*, 1993; Osawa *et al.*, 1995). Studies have shown that supplementation of broilers diets with turmeric enhance their performance (Al-Sultan, 2003; Durrani *et al.*, 2006). However, Mehala & Moorthy (2008) demonstrated that 0.1 and 0.2% turmeric powder used as feed additive had no significant effects on the performance and carcass yield of broiler chickens.

Cinnamaldehyde is the predominant compound of cinnamon, and presents antimicrobial and antioxidant activity (Lee *et al.*, 2004; Faix *et al.*, 2009). Hernandez *et al.* (2004) showed that 200 ppm essential oil extract from oregano, cinnamon and pepper improved nutrient digestibility in broilers. Moreover, the antibacterial activity of cinnamaldehyde and the essential oil obtained from cinnamon leaves was demonstrated by Chang *et al.* (2001).

The inclusion of those herbs in several forms (i.e., powder, fluid extract and essential oil) in poultry diets has been the subject of several studies (Al-Sultan, 2003; Hernandez *et al.*, 2004; Durrani *et al.*, 2006; Mehala & Moorthy, 2008). However, one form that has not received much attention is herbal infusion. This process is commonly used for preparing the more delicate and aromatic parts of a plant, and the obtained components are different from those obtained from the same plant using other methods. The objective of

the present study was to assess the effects of infusions of cinnamon, thyme, turmeric and their combinations as replacement for drinking water on the performance of male broilers.

MATERIAL AND METHODS

Birds, diets and management

A total of 540 newly-hatched male broiler chicks (Ross 308) were purchased from a local hatchery and randomly allocated to 36 litter-floor pens (15 birds per pen). All chicks were fed a corn-soybean based diet daily for 21 days of experimental period (Table 1). Diets were formulated to meet broiler nutrient requirements according to the Ross 308 Management Guideline (2008). Chicks were randomly distributed to one of the four treatment groups or to the control group in a completely randomized design. The experimental treatment involved the replacement of regular drinking water by herb-infused water. The four treatments were infusions (5 grams per liter) of cinnamon, thyme, turmeric and their combination, which contained equal ratios of thyme, turmeric and cinnamon). Each treatment was replicated eight times and the control group four times. Feed, water and herb infusions were provided *ad libitum* and chicks had access to light according to a 23L/1D program. All chicks were vaccinated against Newcastle disease at seven and 17 days, by eye-drop and drinking water, respectively. Vaccination was carried out according to the regional vaccination program routine.

Preparation of herb infusions

Thyme (*Thymus vulgaris*), cinnamon (*Cinnamomum verum*) and turmeric (*Curcuma longa*) were purchased commercially as dry powder. The infusions were prepared daily, according to Durrani *et al.* (2007). The herbal powders were placed in a recipient and freshly boiled water was poured over the herbs (5 grams per liter). The container was sealed and placed at environmental temperature for 8 h. After cooling, the prepared infusion was freshly added to drinkers.

Measured Parameters

The initial and final body weight and the feed intake were weekly recorded for each group. Average daily gain (ADG), average daily feed intake (ADFI) and feed conversion ratio (FCR) were also calculated for each treatment. Live body weight (LBW) was individually measured on day 21. In order to assess blood and carcass



Table 1 - Ingredient and calculated nutritional composition of the diet (g per 100g as fed).

Ingredients	
Corn	56.36
Soybean meal	33.79
Canola meal	2.00
Fish meal	3.00
Soybean oil	1.47
Dicalcium phosphate	1.24
CaCO ₃	1.10
Common salt	0.31
Vitamin premix ¹	0.25
Mineral premix ²	0.25
L-lysine HCL	0.10
DL-methionine	0.14
Calculated composition	
Crude protein(g/kg)	225
Metabolizable energy(MJ/kg)	12.13
Fiber (g/kg)	386
Methionine (g/kg)	5.49
Total sulfur amino acids(g/kg)	9.2
Lysine (g/kg)	13.8
Calcium(g/kg)	9.8
Available phosphorus(g/kg)	4.7

1 - Each kilogram of vitamin supplement contains: Vitamin A, 3600000 IU; vitamin D3, 800000 IU; vitamin E, 7200 IU; vitamin K₃, 800 mg; vitamin B₁, 720 mg; vitamin B₂, 2640 mg; vitamin B₃, 4000 mg; vitamin B₅, 12000 mg; vitamin B₆, 1200 mg; vitamin B₉, 400 mg; vitamin B₁₂, 6 mg; biotin, 40 mg; choline chloride, 100000 mg; antioxidant, 40000 mg.

2 - Each kilogram of mineral supplement contains: Mn, 40000 mg; Zn, 33880 mg; Fe, 20000 mg; Cu, 4000 mg; I, 400 mg; Se, 80 mg; choline chloride, 100000 mg.

characteristics, one bird per replicate were sacrificed at 21 days of age. Blood samples were collected in two tubes, one containing EDTA for hematocrit analysis and the other to evaluate the immune response to NDV by enzyme-linked immunosorbent assay (ELISA). The clotted blood was centrifuged and the serum was separated and stored at -24°C until subsequent analysis. The analysis was performed using the Newcastle Disease Antibody Test Kit from IDEXX laboratories Inc. (Westbrook, ME04092). Carcass, liver, pancreas, heart, spleen, bursa, abdominal fat, gizzard (empty), proventriculus, cecum, duodenum, jejunum and ileum as a percentage of live body weight (relative weight) relative weights and duodenum, jejunum and ileum relative lengths were calculated.

Data analysis

The results of this study were analyzed as

a completely randomized design with unequal replications by analysis of variance (ANOVA), using the general linear model (GLM) procedure of SAS (2001). Means were compared using Duncan's Multiple Range Test with accepted level of significance of 0.05.

RESULTS

Performance parameters

The effects of herbal infusions on weekly performance parameters are presented in Table 2. Individual cinnamon, thyme and turmeric infusions or their combination did not influence LBW, ADG and ADFI in comparison to the control group between 1-7 days of age ($p > 0.05$). However, during this period, turmeric-fed birds presented significantly ($p < 0.05$) higher FCR than those in the control and cinnamon groups. All herbal infusions significantly ($p < 0.05$) increased both ADFI and FCR compared with the control group between 7-14 days of age; however, LBW and ADG were not affected by these additives. In comparison with the control and cinnamon groups, the turmeric-treated group presented significantly lower ($p < 0.05$) ADG between 14-21 days of age. The other evaluated performance parameters were not affected by infusion additives for this age period. None of the herbal infusions had a significant effect on ADG, ADFI and FCR in comparison to control group ($p > 0.05$) during the entire experimental period (1-21 days of age), while all infusion additives significantly decreased LBW ($p < 0.05$).

Blood parameters and carcasses characteristics

The relative weights of the internal organs at 21 days of age are shown in Table 3. The use of a mixed infusion of additives, which contained thyme, turmeric and cinnamon, significantly decreased the weight of ready-to-cook poultry carcasses in comparison to the control group ($p < 0.05$). Thyme-treated broilers showed the highest gizzard and proventriculus relative weights relative to the control broilers ($p < 0.05$). None of the treatments significantly affected the relative weights of the liver, pancreas, heart, abdominal fat, spleen, cecum and bursa.

The relative weights and lengths of the different parts of intestine and entire small intestine are shown in Table 3. These data confirm that thyme and turmeric significantly increased ($p < 0.05$) entire small intestine weight in comparison to the control birds. Turmeric, as compared to the control group, significantly ($p < 0.05$) increased ileum weight. None of the other treatments



Table 2 - Live body weight (LBW), average daily gain (ADG), average daily feed intake (ADFI) and feed conversion ratio (FCR) of experimental male broilers at 1-21 days of age.

Parameters	Age	Treatments ¹					SEM
		Control	Thyme	Turmeric	Cinnamon	Mix	
Live body weight (g)	7	144	141	140	145	139	1.1
	14	336	324	320	334	326	2.4
	21	725a	681b	658b	690b	675b	5.4
Average daily gain (g)	1-7	14.7	14.2	14.1	14.8	13.9	0.15
	7-14	27.4	25.6	25.7	26.5	26.3	0.27
	14-21	51.5a	49.1ab	47.2b	50.6a	48.6ab	0.46
	1-21	30.0	29.4	29.0	30.4	29.3	0.25
Average daily feed intake (g)	1-7	19.2	19.1	19.7	19.3	18.8	0.16
	7-14	34.3b	40.1a	39.3a	39.7a	38.6a	0.53
	14-21	78.8	77.2	74.0	76.7	75.3	0.69
	1-21	48.3	49.0	48.7	48.6	48.0	0.26
Feed conversion ratio (g/g)	1-7	1.30b	1.34ab	1.40a	1.30b	1.35ab	0.012
	7-14	1.24b	1.57a	1.53a	1.50a	1.47a	0.022
	14-21	1.52	1.58	1.57	1.52	1.55	0.016
	1-21	1.61	1.67	1.68	1.60	1.64	0.013

1- Infusion (5 grams per liter) of thyme, cinnamon, turmeric or equal ratio of their combination (mix) as replacement of the drinking water. SEM: Standard error of the mean. a,b - Means within each row followed by the same letter are not significantly different at 0.05.

Table 3 - Relative weight and length of organs (% of live weight) of the experimental birds at 21 days of age.

	Treatments ¹					SEM
	Control	Thyme	Turmeric	Cinnamon	Mix	
Relative weight						
Ready-to-cook carcass	55.29a	54.90a	52.97ab	54.84a	50.51b	0.529
Liver	2.72	2.70	2.87	3.01	2.77	0.061
Pancreas	0.416	0.452	0.425	0.477	0.433	0.0120
Heart	0.577	0.624	0.580	0.600	0.638	0.0135
Gizzard	2.77b	3.44a	3.20ab	3.01ab	2.90ab	0.084
Proventriculus	0.585b	0.692a	0.613ab	0.644ab	0.699a	0.0133
Abdominal Fat	0.754	0.822	0.760	0.796	0.981	0.0499
Bursa	0.278	0.281	0.246	0.272	0.300	0.0107
Spleen	0.071	0.081	0.070	0.073	0.071	0.0024
Cecum	0.525	0.545	0.621	0.598	0.554	0.0244
Entiresmall intestine	4.30b	4.84a	4.89a	4.71ab	4.70ab	0.075
Duodenum	1.17	1.27	1.21	1.28	1.18	0.025
Jejunum	1.84	2.02	2.00	1.85	2.01	0.045
Ilium	1.29b	1.55ab	1.68a	1.57ab	1.51ab	0.042
Relative length						
Entiresmall intestine	25.59	28.24	27.67	26.04	27.52	0.458
Duodenum	0.047	0.050	0.047	0.046	0.046	0.0009
Jejunum	0.100	0.116	0.112	0.102	0.115	0.0024
Ilium	0.109	0.116	0.118	0.113	0.114	0.0023

1 - Infusion (5 grams per liter) of thyme, cinnamon, turmeric or equal rates of their combination (mix) as replacement of drinking water. SEM: Standard error of the means. a,b - Means within each row followed by the same letter are not significantly different at 0.05.



significantly influenced the relative weights and lengths of the duodenum, jejunum or entire small intestine.

The effects of the additive infusions on the immune response to the NDV vaccine and on the hematocrit are presented in Table 4. None of the herbal infusions significantly affected the hematocrit relative to the control group ($p>0.05$). The cinnamon and herbal mix infusions significantly improved the immune response to the NDV vaccine in comparison to the control group and those that received only the turmeric infusion ($p<0.05$).

DISCUSSION

Performance parameters

The results of this study indicate that the use of herbal infusions as dietary supplementation to young male broiler chickens did not significantly improve performance parameters in comparison to control group.

The study showed that thyme supplied as an herbal infusion did not significantly improve the performance of male broilers. This is consistent with Ocak *et al.* (2008), who reported that the inclusion of 0.2% thyme leaves in broiler diets had no effects on body weight gain, feed intake or feed conversion ratio. Cross *et al.* (2007) also reported that the performance parameters of male broilers were not improved when their diets were supplemented with 1% thyme, whereas the results of the present study showed that thyme increased LBW and weight gain of the thyme-treated broilers. The results also agree with the findings of Mehala & Moorthy (2008), who reported that diets supplemented with 0.1 and 0.2% turmeric had no detrimental effect on broiler performance. In contrast to this study, Durrani *et al.* (2006) showed that adding 0.5% turmeric to broiler diets resulted in worse ADFI and FCR both during the starter and finisher periods, but weight gain decreased only during the finisher period. Al-Sultan (2003) reported that adding 0.5% turmeric to broilers diet improved all their performance characteristics. Our results showed that turmeric infusion reduced live body weight more than the other evaluated infusions.

The performance of cinnamon-treated broilers was not different from those receiving the other treatments. These findings are in agreement with those of Hernandez *et al.* (2004), who reported that the incorporation of 200 ppm of oregano, cinnamon and pepper essential oils in broilers diets had no effect on their performance at 21 and 42 days of age (Hernandez *et al.*, 2004). Lee *et al.* (2003) also reported that weight gain, feed intake and feed conversion ratio were not affected by 100 ppm dietary thymol or cinnamaldehyde.

In most of the previous studies, herb powder or essential oil were added to the broiler diets, but in the present study, the aqueous extracts of these herbs was used in a different way and via drinking water. Therefore, some of the inconsistencies observed in this study in comparison to others may be due to the method of preparation of the additives.

Blood parameters and carcasses characteristics

Relative weight of ready-to-cook carcasses was not affected by the herbal infusions tested in present study. However, the mixture of these herbal medicinal plants significantly reduced carcass weight ($p<0.05$). Ocak *et al.* (2008) reported that broiler carcass relative weight was not affected by the inclusion of 0.2% thyme leaves in the diet. Consistent with present results, some previous studies showed that gizzard relative weight was not affected by turmeric herb or cinnamon essential oil (Durrani *et al.*, 2006; Hernandez *et al.*, 2004). The present data showed that thyme and turmeric increased small intestine relative weight. In addition, turmeric infusion increased ileum relative weight; however, the length of the intestinal sections were not affected by the treatments. Therefore, the higher gut weights in the turmeric-treated birds may have been due to higher diameters of the gut wall that affected digestive processes, resulting in lower live body weight. In contrast with findings of the present study, Ocak *et al.* (2008) reported that no changes were detected in the gut relative weight of broilers fed with 0.2% thyme leaves.

Cinnamon and herbal mix infusions improved antibody titers against NDV in comparison to control group. There is no published work about the effects of

Table 4 - Immune response to live Newcastle disease virus (NDV) and hematocrit percent of the experimental birds at 21 days of age.

	Treatments ¹					SEM
	Control	Thyme	Turmeric	Cinnamon	Mix	
Hematocrit (%)	32.5	35.6	34.0	32.6	32.6	0.62
Antibody titer	310.3b	485.1ab	177.2b	529.0a	767.3a	0.193

1 - Infusion (5 g/L) of thyme, cinnamon, turmeric or equal ratio of their combination (mix) as replacement of the drinking water. SEM: Standard error of means. a, b - Means within each row followed by the same letter are not significantly different at 0.05.



herb infusions on antibody titers. However, Al-Ankari *et al.* (2004) reported that mint powder had no effects on antibody titers to NDV vaccine.

CONCLUSION

The three herb infusions (cinnamon, thyme and turmeric) used as drinking water did not improve the performance of broilers. However, cinnamon and herbal mix infusions significantly improved the immune response of birds to the NDV vaccine. When the additives were compared, the broilers supplied with the turmeric infusion presented the worst performance and immunity.

REFERENCES

- Al-Ankari AS, Zaki MM, Al-Sultan SI. Use of habek mint (*menthalongifolia*) in broiler chicken diets. *International Journal of Poultry Science* 2004; 3:629-634.
- Al-Kassie GAM. Influence of two plant extracts derived from thyme and cinnamon on broiler performance. *Pakistan Veterinary Journal* 2009; 29:169-173.
- Al-Sultan SI. The effect of curcuma longa (turmeric) on overall performance of broiler chickens. *International Journal of Poultry Science* 2003; 2: 351-353.
- Ammon HPT, Safayhi H, Mack T, Sabieraj J. Mechanism of antiinflammatory actions of curcumin and boswellic acids. *Journal of Ethnopharmacology* 1993; 38:113-119.
- Canan Bölükbaşı, Kuddusi Erhan M. Effect of dietary thyme (*Thymus vulgaris*) on laying hens performance and escherichia coli (*E. coli*) concentration in feces. *International Journal of Natural and Engineering Sciences* 2007; 1:55-58.
- Chang ST, Chen PF, Chang SC. Antibacterial activity of leaf essential oils and their constituents from *Cinnamomum mosophloeum*. *Journal of Ethnopharmacology* 2001; 77:123-127.
- Cross DE, Mcdevitt RM, Hillman K, Acamovic T. The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in chickens from 7 to 28 days of age. *British Poultry Science* 2007; 48:496-506.
- Durrani FR, Ismail M, Sultan A, Suhail SM, Chand N, Durrani Z. Effect of different levels of feed added turmeric (*Curcuma longa*) on the performance of broiler chicks. *Journal of Agricultural and Biological Science* 2006; 1:9-11.
- Durrani FR, Sultan A, Marri ML, Chand N, Durrani Z. Effect of wild mint (*menthalongifolia*) infusion on the overall performance of broiler chicks. *Pakistan Journal of Biological Sciences* 2007; 10:1130-1133.
- El-Ghoussein SS, Al-Beitawi NA. The effect of feeding of crushed thyme (*Thymus vulgaris* L.) on growth, blood constituents, gastrointestinal tract and carcass characteristics of broiler chickens. *The Journal of Poultry Science* 2009; 46:100-104.
- Faix Š, Faixová Z, Plachá I, Koppel J. Effect of cinnamomumzeylanicum essential oil on antioxidative status in broiler chickens. *Acta Veterinaria Brno* 2009; 78:411-417.
- Hernandez F, Madrid J, García V, Orengo J, Megías MD. Influence of two plant extracts on broilers performance, digestibility, and digestive organ size. *Poultry Science* 2004; 83:169-174.
- Kiuchi F, Goto Y, Sugimoto N, Akao N, Kondo K, Tsuda Y. Nematocidal activity of turmeric: synergistic action of curcuminoids. *Chemical and Pharmaceutical Bulletin* 1993; 41:1640-1643.
- Lee KW, Everts H, Beynen AC. Essential oils in broiler nutrition, *International Journal of Poultry Science* 2004; 3:738-752.
- Lee KW, Everts H, Kappert HJ, Frehner M, Losa R, Beynen AC. Effects of dietary essential oil components on growth performance, digestive enzymes and lipid metabolism in female broiler chickens. *British Poultry Science* 2003; 44:450-457.
- Mehala C, Moorthy M. Production performance of broilers fed with aloe vera *Curcuma longa* (turmeric). *International Journal of Poultry Science* 2008; 7:852-856.
- Najafi P, Torki M. Performance, blood metabolites and immunocompetence of broiler chicks fed diets included essential oils of medicinal herbs. *Journal of Animal and Veterinary Advances* 2010; 9:1164-1168.
- Ocak N, Erener G, Burak Ak F, Sungu M, Altop A, Özmen A. Performance of broilers fed diets supplemented with dry peppermint (*Mentha piperita* L.) or thyme (*Thymus vulgaris* L.) leaves as growth promoter source Czech. *Journal of Animal Science* 2008; 53:169-175.
- Osawa T, Sugiyama Y, Inayoshi M, Kawakishi S. Antioxidative activity of tetrahydrocurcuminoids. *Bioscience, Biotechnology and Biochemistry* 1995; 59:1609-1612.
- Patterson JA, Burkholder KM. Application of prebiotics and probiotics in poultry production. *Poultry Science* 2003; 82:627-631.
- SAS Institute. SAS user's guide: statistic. Version 8.2. Cary; 2001.