



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e
Tecnologia Avícolas
Brasil

Elerolu, E; Yldrm, A; Duman, M; ekerolu, A
Edible GIBLETS and Bone Mineral Characteristics of Two Slow-Growing Chicken
Genotypes Reared in an Organic System
Revista Brasileira de Ciência Avícola, vol. 19, núm. 1, enero-marzo, 2017, pp. 1-6
Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179750697001>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



■ Author(s)

Eleroğlu E^I
Yıldırım A^{II}
Duman M^{III}
Şekeroğlu A^{IV}

^I Şarkışla Aşık Veysel Vocational School,
Cumhuriyet University, Sivas/Turkey
e-mail: eleroglu@cumhuriyet.edu.tr

^{II} Department of Animal Science, Faculty of
Agriculture, University of Gaziosmanpaşa,
Tokat/Turkey
e-mail: arda.yildirim@gop.edu.tr

^{III} Bor Vocational High School, Ömer Halisde-
mir University, Niğde/Turkey
e-mail: mustafa.duman@ohu.edu.tr

^{IV} Department of Animal Production and Te-
chnologies, Faculty of Agricultural Sciences
and Technologies, Ömer Halisdemir Univer-
sity, Niğde/Turkey
e-mail: ahmet.sekeroglu@ohu.edu.tr

■ Mail Address

Corresponding author e-mail address
Hasan Eleroğlu
Informatics Department, Cumhuriyet
University, 58146 Sivas/Turkey
Tel: +90 532 2041066
Email: eleroglu@cumhuriyet.edu.tr

■ Keywords

Organic production, broiler strain, bone
mineral density, bone mineral content.

Edible Giblets and Bone Mineral Characteristics of Two Slow-Growing Chicken Genotypes Reared in an Organic System

ABSTRACT

This study was conducted to compare edible giblets weight, tibial bone mineral density (BMD), and bone mineral content (BMC) of two slow-growing broiler genotypes (Hubbard S757; S757 and Hubbard Grey Barred JA; GB-JA) reared with outdoor access, and to determine the relationship between these variables. Day-old chicks (straight-run) of the genotypes S757 (n=120) and GB-JA (n=120) were housed for 98 days. Each genotype was assigned to six pens of 20 birds each. Birds were reared in indoor floor pens and moving shelters with outdoor access (during daylight hours). Absolute body (BW), heart (HW), spleen (SW), liver (LW), gizzard (GW), and abdominal fat pad (AFW) weights of the genotype S757 and male birds were statistically higher than that of the genotype GB-JA and female birds. Genotype statistically affected relative HW, whereas sex affected relative GW. Although BMD values were not influenced by genotype or sex, S757 birds and males presented statistically higher tibial BMC, lean, lean+BMC, total mass values (g) and area (cm²) compared with GB-JA birds and females. BW, HW, SW, LW, GW and AFW were positively correlated with BMC obtained by DXA. In conclusion, the measured traits influenced by genetic strain and sex. The use of the Hubbard S757 genotype in organic production systems with outdoor access is recommend.

INTRODUCTION

Although environmental factors such as management, feeding, movement, toxins, and infectious diseases can play very important roles in the development of skeletal structure, genetic background also has a fundamental influence (Rath *et al.*, 2000). Modern broiler genotypes have been subjected to intensive selection for body weight gain. This selection has resulted in major changes in the anatomy and physiology of broilers, such as increased fat deposition, metabolic diseases, and abnormal skeletal development, including leg disorders, lameness, twisted legs, tibial dyschondroplasia, and crooked toes (Bradshaw *et al.*, 2002; Decuyper *et al.*, 2003; Oviedo-Rondon *et al.*, 2006; Scawah, 2000).

Conventional intensive production conditions do not allow sufficient exercise and typically present poor hygiene, resulting in leg weakness and skin infections (Bessei, 2006; Talaty *et al.*, 2009). It has been demonstrated that the bones of fast growing meat-type chickens are often abnormally developed, and present high porosity and much lower density than those of slow-growing broilers (Bennett, 2008; Williams *et al.*, 2004). Chickens living freely in their natural habitats did not appear to have such problems (Newberry, 1995), and present much stronger muscle structures (Balog, 1997; Le Van, 2000; Shields, 2004). The genetic selection for growth rate and the feed conversion ratio has altered the relative growth of organs in modern broiler lines when



compared with heritage lines (Lippens, 2003; Schmidt *et al.*, 2009) and the increasing muscle weight of modern broiler lines is accompanied by a decrease in edible giblets weight (Janiszewska *et al.*, 1998; Plavnik & Hurwitz, 1982). The changes in the weight of edible giblets occur at different rates (Murawska *et al.*, 2011; Tüzün & Aktan, 2012).

Because of bone ash (Onyango *et al.*, 2003) and its high correlation with live scans (Schreiweis *et al.*, 2005), excised bones can also be scanned using DXA to assess bone quality. Bone mineral density (BMD, g/cm²) (Kim *et al.*, 2006; Lian *et al.*, 2004; Onyango *et al.*, 2003; Rath *et al.*, 2000; Shim *et al.*, 2012; Watkins and Southern, 1992) and bone mineral content (BMC, g/cm) (Akpe *et al.*, 1987; Almeida *et al.*, 2008; Kim *et al.*, 2006; Onyango *et al.*, 2003) measurements have been used to evaluate the quality and the status of the porous structure of bone matrix. These two parameters are greatly influenced by mineral intake. BMD is a noninvasive method using dual-energy x-ray absorptiometry (DXA) that has also been employed to predict osteoporosis in humans (Bolotin, 2007; Koo, 2000). DXA could also be used to measure BMC in broilers and laying hens (Schreiweis *et al.*, 2005; Shim *et al.*, 2012; Talaty *et al.*, 2010).

Comparative research among different slow-growing broiler strains may be used to establish physiological and morphological adaptations in response to distinct selective pressures. The purpose of this study was to compare the BMD, BMC, and giblet weights of two slow-growing broiler genotypes selected for organic production.

MATERIAL AND METHODS

Two hundred and forty male and female Hubbard S757 (S757) and Grey Barred JA (GB-JA) were equally divided between strains were used. Although both these strains present slow growth rates to be used in organic production systems, Hubbard S757 (S757) and Grey Barred JA (GB-JA) were selected to achieve market weight at 84 and 70 days of age respectively. In the study, one-d-old chicks were weighed, identified by wing band, and randomly allocated to two treatments (genotypes: Hubbard S757 or Hubbard Grey Barred JA) with six replicates of 20 birds each. The experiment was approved by the Ethics Committee of the University of Cumhuriyet in Sivas, Turkey (20.06.2011/50).

Twelve portable shelters (1.5 x 1.5 m), each housing 20 birds per replicate at a stocking density of 10 birds/m² placed in each of the 100-m² grazing areas. The research was carried out according to the principles and

implementation of regulation on organic agriculture (OFL, 2010) published by the Republic of Turkey, Ministry of Food, Agriculture and Livestock. During the first 14 days, chicks were housed in the portable shelters, and offered feed and water *ad libitum*. After this period chicks were allowed to access outdoors and to graze freely. All feed and water were provided between 07:00-19:00 for all chicks during the entire experimental period.

Birds were fed with a three-phase organic diet: a starter diet (197 g crude protein (CP)/kg and 13.00 MJ of metabolizable energy (ME) per kg of diet) from 0 to 28 days, a grower diet (201 g/kg CP and 12.72 MJ of ME per kg of diet) from 29 to 81 days, and a finisher diet (180 g/kg CP and 12.91 MJ of ME per kg of diet) from 82 to 98 days (Table 1).

Only natural day length lighting was provided for chickens from first days to slaughter age. Ceramic heaters, which are sources of far infrared rays and do not emit light, were used for heating.

All birds used in the experiment were managed following the recommendations of Sirri *et al.* (2011). Birds were fed certified organic feedstuffs, and provided with a cultivated poultry pasture, consisting of equal amounts of *Lotus corniculatus* and *Bromus inermis*.

At the end of the 98-day trial period, bone properties were assessed in 24 males and 24 females whose body weights (BW) were close to the pen average (Eleroğlu *et al.*, 2014a). These 48 birds were fasted for 10 h (with free access to water), weighed, and slaughtered by severing the throat and major blood vessels of the neck in the local processing plant producing organic chicken products (Eleroğlu *et al.*, 2014b). After slaughter, hot carcass weight was determined in a semi-analytic digital scale with 0.001-g precision, and then the left tibial bone, heart, spleen, liver, gizzard, and abdominal fat pad were removed and weighed to determine absolute weight and relative weight as a percentage of live body weight. Bone mineral content (BMC), bone mineral density (BMD), lean, lean+BMC, total mass and mass area of the left tibial bone were determined using a total-body dual energy X-ray absorptiometry DXA scanner (QDR 4500W Acclaim, Hologic, USA) at the Nuclear Medicine Department of the Medicine Faculty at Cumhuriyet University.

Data were analyzed using the General Linear Model (GLM) procedure. The effects of genotype, sex, and their interactions on the absolute and relative weights of the body (BW), heart (HW), spleen (SW), liver (LW), gizzard (GW), and abdominal fat (AFW), as well as



Table 1 – Ingredients and calculated nutritional composition of the experimental diets

Feedstuffs, %	0–28 days	29–81 days	82–98 days	Nutritional composition (g/kg)			
Barley	3.45	4.50	4.50	ME (MJ/kg)	13.00	12.72	12.91
Vegetable oil	4.36	5.00	5.00	Dry matter	899.00	903.00	901.00
Wheat bran	5.00	5.00	5.00	Crude protein	197.00	201.00	180.00
White wheat	12.40	4.00	4.00	Crude ash	4.70	5.90	4.80
Rye	3.00	4.00	4.00	Lysine	10.80	10.60	8.50
Corn	40.00	20.00	20.00	Met + Cys.	6.60	6.70	5.90
Triticale	-	22.00	32.00	Threonine	7.30	7.20	6.20
Oats	2.10	5.00	-	Calcium	10.00	11.60	9.00
Fish meal	7.30	5.00	-	Phosphorus	7.70	6.00	5.90
Soybean meal	20.00	22.00	22.00	Sodium	1.90	1.80	1.50
Dicalcium phosphate	1.10	2.10	2.10	Tryptophan	2.40	2.60	2.50
Limestone	0.74	0.80	0.80	Linoleic acid	31.9	32.1	31.3
Salt	0.30	0.30	0.30				
Vitamin-mineral premix*	0.25	0.30	0.30				

Explanation: * - Each kg of vitamin-mineral premix contained: vitamin A, 4,400,000 IU; vitamin D₃, 1,600,000 IU; vitamin E, 20,000 mg; vitamin K₃, 1,600 mg; vitamin B₁, 1,200 mg; vitamin B₂, 3,200 mg; vitamin B₃, 20,000 mg; vitamin B₅, 6,000 mg; vitamin B₆, 1,600 mg; vitamin B₉, 800 mg; vitamin B₁₂, 8 mg; biotin, 80 mg; antioxidant, 50,000 mg; Cu, 6,000 mg; Fe, 20,000 mg; Mn, 48,000 mg; Se, 80 mg; Zn, 40,000 mg; Co, 80 mg; I, 500 mg.

on BMD and BMC were evaluated using multifactor analysis of variance. The percentage data were converted to arcsines prior to analysis. The relationship between bone mineral characteristics and edible giblets were determined using one-tailed Pearson correlation procedure (SPSS Inc. 2010, Release 16.0). Treatment effects were considered significant at $p < 0.05$. Data were expressed as mean values with pooled standard errors (standard errors of the mean, SEM).

RESULTS

The results of the effects of genotype and sex on edible giblets, abdominal fat, and bone characteristics are given in Table 2 and 3, respectively.

Body weight at slaughter (BW) of the two genotypes investigated under the conditions of the present experiment was accurately measured, and was different from each other. Before slaughter, genotype and sex significantly influenced BW ($p < 0.01$). As shown in Table 2, S757 males were heavier (2847 g) than GB-JA males (2128 g) and S757 females (2194 g) were heavier than GB-JA females (1634 g). Males of both evaluated strains were heavier than females ($p < 0.01$). There were significant differences between males and females of the same genotypes ($p < 0.01$). S757 males showed higher BW (2847 g) than females (2194 g), whereas the calculated BW of males and females of the GB-JA genotype were 2128 g and 1634 g, respectively.

Table 2 – Absolute and relative weights of edible giblets and abdominal fat of male and female broilers of two different slow-growing genotypes.

	GB-JA ¹		S757 ¹		SEM ²	p value	
	Female	Male	Female	Male		G ³	S ⁴
Absolute weight (g)							
Body	1634	2128	2194	2847	65.31	**	**
Heart	7.13	9.57	8.59	11.91	0.31	**	**
Spleen	2.07	2.40	3.17	3.74	0.13	**	*
Liver	24.55	31.35	31.73	38.71	0.9	**	**
Gizzard	32.14	37.16	41.37	46.42	1.04	**	**
Abdominal fat pad	27.75	38.44	39.14	50.34	2.25	**	**
Relative weight (g/100 g body weight)							
Heart	0.44	0.45	0.39	0.42	0.008	**	NS
Spleen	0.13	0.11	0.15	0.13	0.005	NS	NS
Liver	1.51	1.47	1.45	1.36	0.024	NS	NS
Gizzard	1.97	1.75	1.90	1.63	0.042	NS	**
Abdominal fat pad	1.70	1.79	1.78	1.78	0.087	NS	NS

Explanations: ¹ - GB-JA = Hubbard Grey Barred JA; S757 = Hubbard S757; ² - Standard error of the mean, G³ - Genotype; S⁴ - Sex; * - $p < 0.05$; ** - $p < 0.01$; NS - Not significant ($p > 0.05$)



Table 3 – Bone characteristics of male and female broilers of two different slow-growing genotypes

Genotype (G) ¹	Sex (S)	BMC (g)	Lean (g)	Lean+BMC (g)	Total mass (g)	Mass Area (cm ²)	BMD (g/cm ²)
GB-JA	F	2.37	55.78	58.15	57.10	9.96	0.237
	M	4.55	86.75	91.30	89.38	16.52	0.269
S757	F	3.57	81.65	85.22	84.31	14.45	0.256
	M	5.88	109.45	115.35	113.78	21.18	0.269
SEM ²		0.29	3.54	3.78	3.76	0.76	0.009
Main effect of							
G ³		**	**	**	**	**	NS
S ⁴		**	**	**	**	**	NS

Explanations: ¹ - GB-JA = Hubbard Grey Barred JA; S757 = Hubbard S757; ² - Standard error of the mean; G³ – Genotype; S⁴ – Sex; * - $p < 0.05$; ** - $p < 0.01$, NS - Not significant ($p > 0.05$)

Absolute HW, SW, LW, GW and AFW (g) were influenced by genotype, with S757 birds presenting higher values than GB-JA birds ($p < 0.01$). On the other hand, no differences in HW, SW, LW, GW, and AFW relative to body weight (g/100 g body weight) were detected between genotypes, except for HW, which was higher ($p < 0.01$) in GB-JA (0.45 g) than in S757 (0.41 g) birds.

The absolute HW (11.91 g), SW (3.74 g), LW (38.71 g), GW (46.42 g) and AFW (50.34 g) of S757 males were higher ($p < 0.01$) than absolute HW (9.57 g), SW (2.40 g), LW (31.75 g), GW (37.16 g) and AFW (38.44 g) of GB-JA males. Similarly, absolute HW (8.59 g), SW (3.17 g), LW (31.73 g), GW (41.37 g) and AFW (39.14 g) values of S757 females were higher compared with GB-JA females, which presented HW (7.13 g), SW (2.07 g), LW (24.55 g), GW (32.14 g) and AFW (27.75 g) ($p < 0.01$, $p < 0.05$). Neither genotype nor sex influenced ($p > 0.05$) relative SW, LW and AFW (g/100 g BW). However, relative HW ($p < 0.01$) was significantly higher in S757 birds compared with GB-JA birds, independently of sex, and males presented higher relative GW than females ($p < 0.01$), independently of genotype.

A strong effect of genotype and sex was observed on BMC, Lean, Lean+BMC, total mass and mass area ($p < 0.01$). In contrast, genotype and gender effects on BMD were not significant ($p > 0.05$).

As shown in Table 3, the average BMC (4.72 g), Lean (95.55 g), Lean+BMC (100.28 g), total mass (99.05 g) and mass area (17.82 cm²) values of S757 birds were higher ($p < 0.01$) than the average BMC (3.46 g), Lean (71.27 g), Lean+BMC (74.73 g), Total mass (73.24 g) and Mass area (13.24 cm²) values of GB-JA genotype. On the other hand similar differences were observed between sex, the average BMC (5.51 g), Lean (98.1 g), Lean+BMC (103.33 g), Total mass (101.58 g) and Mass area (18.85 cm²) values of male were higher than the average BMC (2.97 g), Lean (68.72 g), Lean+BMC (71.69 g), Total mass (70.71 g) and Mass area (12.21 cm²) values of female ($p < 0.01$).

Table 4 shows the correlations between BW, giblets, abdominal fat, BMC, and BMD values. HW, SW, LW, GW and AFW were significantly correlated with BW ($r = 0.839, 0.711, 0.826, 0.668$ and 0.526 , respectively, $p < 0.01$, Table 4). A high correlation of 0.839 ($p < 0.01$) was especially noted between HW and BW, whereas the correlation of AFW between HW was low ($r = 0.307$; $p < 0.05$). In addition, BMD, BW, HW, SW, LW and AFW were highly significantly correlated with BMC ($r = 0.687, 0.543, 0.415, 0.396, 0.462$ and 0.434 , respectively, $p < 0.01$) and the correlation GW with BMC was low ($r = 0.319$; $p < 0.05$).

DISCUSSION

The obtained absolute and relative HW, SW, LW, GW and AFW results account indicate that bird growth was isometric, and their genotype, management, and feeding program allowed their normal growth and development under the organic rearing conditions (Plavnik & Hurwitz, 1982; Lippens, 2003).

The differences in the relative HW detected between the S757 and GB-JA genotypes may be explained by the selection for different growth rates. The differences in BW led to different relative GW values. Edible internal organ weights relative to hot carcass weight were expected to be higher as the BW at slaughter of broilers reared in organic systems is higher (Eleroglu et al., 2014b). The findings of the current study confirmed this fact.

BMD was not influenced by genotype or sex, on account of this can be said to be similar applied to environmental conditions for both genotype and gender.

Simsek et al. (2009) observed BMC and BMD values of 1.94; 0.12 and 2.01; 0.14 in 42-d-old broilers reared in an enriched and a conventional housing system, respectively. In the present study, BMC values ranged between 2.37 and 5.88, and BMD values between 0.237 and 0.269 in broilers reared in an organic system. The higher BMC and BMD values obtained in



Table 4 – Correlation coefficients among BW, edible giblets, abdominal fat, BMC and BMD

Items	BMD	BMC	BW	HW	SW	LW	GW
BMD	1						
BMC	0.687**	1					
BW	0.125	0.543**	1				
HW	0.065	0.415**	0.839**	1			
SW	0.136	0.396**	0.711**	0.537**	1		
LW	0.058	0.462**	0.826**	0.767**	0.668**	1	
GW	0.059	0.319*	0.668**	0.620**	0.509**	0.679**	1
AFW	0.145	0.434**	0.526**	0.307*	0.399**	0.484**	0.136

Explanations: ** - Correlation is significant at * $p < 0.05$; ** - $p < 0.01$

the present study may be attributed to age at slaughter (98 days), whereas in Simsek *et al.* (2009), broilers were slaughtered with 42 days of age. Rath *et al.* (2000) indicated that BMC and BMD vary according to the age of the skeletal structure, Williams *et al.* (2004) and Bennett (2008) slow-growing genotypes present better skeletal structure compared fast-growing broilers.

In the present study, a positive correlation between BW and edible giblets and abdominal fat was determined. Genetics and environmental factors, such as feeding, housing, and management, influence BW and the relative growth of internal organs (Ramadan *et al.*, 2014; Tavárez *et al.*, 2016), resulting in changes in the coefficients of correlation between BW and organ relative weights. The correlation coefficients were unexpected and possibly related with allometric internal organ growth.

The positive correlation between BW and BMC observed in the present study is in agreement with previous studies (Schreiweis *et al.*, 2004; Salas *et al.*, 2012). In addition, this correlation indirectly affected the correlation of BMC with the internal organs. Bone breaking strength, bone ash concentration, and bone weight are positively correlated with BMD and BMC obtained by DXA in chickens (Fleming *et al.*, 1994; Mazzuco and Hester, 2005; Kim *et al.*, 2006, 2008). There is a strong correlation between BMD and BMC ($r=0.81$; $p<0.01$) in White Leghorns (Hester *et al.*, 2004) as well in male broilers of commercial lines (Talaty *et al.*, 2010). In addition, relatively larger bones present higher BMC correlation with total bone mineral content and are stronger and BMD and BMC values increase with broiler BW (Schreiweis *et al.*, 2004).

The correlation between BMC and tibia ash content is reported as 86% in commercial broilers (Onyango *et al.*, 2003), and BMC has been described as a better indicator of Ca and P nutrition than BMD (Yan *et al.*, 2005). It is concluded that DXA can be used to assess bone mineralization status and bird welfare as a function of their correlation with BMC and BMD.

CONCLUSIONS

The results of this study showed the importance of genetic and environmental influences on the bone development of broilers. The positive correlation between BMD values and the weight of internal organs should be further studied.

ACKNOWLEDGEMENTS

This study was supported by the Research Fund of Cumhuriyet University (Project No: ENF-003), which was approved by the local Ethical Committee of Cumhuriyet University for Experimental Animals.

REFERENCES

- Akpe MP, Waibel PE, Larntz K, Metz AL, Noll SL, Walser MM. Phosphorous availability bioassay using bone ash and bone densitometry as response criteria. *Poultry Science* 1987; 66: 713–720
- Almeida Paz ICL, Mendes AA, Balog A, Vulcano LC, Ballarin AW, Almeida ICL, Takahashi SE, Komiyama CM, Silva MC, Cardoso KFG. Study on the bone mineral density of broiler suffering femoral joint degenerative lesions. *Brazilian Journal of Poultry Science* 2008; 10: 103–108
- Balog JM, Bayyari GR, Rath NC, Huff WE, Anthony NB. Effect of intermittent activity on broiler production parameters. *Poultry Science* 1997; 76: 6–12
- Bennett MB. Post-hatch growth and development of the pectoral and pelvic limbs in the black nobby, Anous minutus. *Comparative Biochemistry and Physiology* 2008; A150: 159–168
- Bessei W. Welfare of broilers: a review. *World Poultry Science* 2006; J 62: 455–466
- Bolotin HH. DXA in vivo BMD methodology: An erroneous and misleading research and clinical gauge of bone mineral status, bone fragility, and bone remodelling. *Bone* 2007; 41: 138–154
- Bradshaw RH, Kirkden RD and Broom DM. A review of the aetiology and pathology of leg weakness in broilers in relation to welfare. *Avian And Poultry Biology Reviews*. 2002; 13: 45–103
- Decuypere E, Bruggeman V, Barbato GF and Buyse J. Growth and reproduction problems associated with selection for increased broiler meat production. *Poultry Genetics, Breeding and Technology*, CABI-publishing, Wallingford 2003.
- Eleroğlu H, Yıldırım A, Işıklı ND, Şekeroğlu A, Duman M. Comparison of meat quality and fatty acid profile in slow-growing chicken genotypes fed diets supplemented with *Origanum vulgare* or *Melissa officinalis*



- leaves under the organic system under the organic system, Italian Journal of Animal Science 2014a; 12: 395-403
- Eleroğlu H, Yıldırım A, Şekeroğlu A, Duman M. Comparison of the growth performance and carcass characteristics of two slow-growing broiler genotypes fed diets supplemented with dry oregano (*Origanum vulgare* L.) or lemon balm (*Melissa officinalis* L.) leaves under the organic system, Kafkas Üniversitesi Veteriner Fakültesi Dergisi. 2014b; 20: 49-58
- Fleming RH, Whitehead CC, Alvey D, Gregory NG, Wilkins LJ. Bone structure and breaking strength in laying hens housed in different husbandry systems. British Poultry Science. 1994; 35: 651-662
- Hester PY, Schreiweis MA, Orban JI, Mazzucco H, Kopka MN, Ledur MC, Moody DE. Assessing bone mineral density in vivo: Dual energy X-ray absorptiometry. Poultry Science 2004; 83: 215-221
- Janiszewska M, Bochno R, Lewczuk A, Brzozowski W. Changes in the weight of body, parts of carcass and tissue components in broiler and laying chickens during the growing period. Acta Academiae Agriculturae ac Technicae Olstenensis Veterinaria 1998; 48: 103-114
- Kim WK, Donalson LM, Mitchell AD, Kubena LF, Nisbet DJ, Ricke SC. Effects of alfalfa and fructooligosaccharide on molting parameters and bone qualities using dual energy X-ray absorptiometry and conventional bone assays. Poultry Science 2006; 85: 15-20
- Kim WK, Herfel TM, Dunkley CS, Hester PY, Crenshaw TD, Ricke SC. The effects of alfalfa-based molt diets on skeletal integrity of white leghorns. Poultry Science. 2008; 87: 2178-2185
- Koo WWK. Body composition measurements during infancy. Annals of the New York Academy of Sciences 2000; 904: 383-392
- Le Van NF, Estevez I, Stricklin WR. Use of horizontal and angled perches by broiler chickens. Applied Animal Behaviour Science 2000; 65: 349-365
- Lian JB, Javed A, Zaidi SK, Lengner C, Montecino M, van Wijnen AJ, Stein JL, Stein GS. Regulatory controls for osteoblast growth and differentiation: role of Runx/Cbfa/AML factors. Crit. Rev. Eukaryot. Gene Expr 2004; 14: 1-41
- Lippens M. The influence of feed control on the growth pattern and production parameters of broiler chickens. PhD Thesis, Universiteit Gent, Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen 2003; 203 pp.
- Mazzucco H, Hester PY. The effect of an induced molt and a second cycle of lay on skeletal integrity of White Leghorns. Poultry Science 2005; 84: 771-781
- Murawska D, Kleczek K, Wawro K, Michalik D. Age-related changes in the percentage content of edible and non-edible components in broiler chickens. Asian-Australasian Journal of Animal Sciences 2011; 24: 532-539
- Newberry RC. Environmental enrichment. Increasing the biological relevance of captive environments. Applied Animal Behaviour Science. 1995; 44: 229-243
- OFL. Organic Farming Legislation, published by the Republic Of Turkey Ministry Of Food, Agriculture and Livestock, Turkey Official Gazette 2010; 27676
- Onyango EM, Hester PY, Strohshen RL, Adeola O. Bone densitometry as an indicator of percent tibia ash in broiler chicks fed varying dietary calcium and phosphorus levels. Poultry Science 2003; 82: 1787-1791
- Oviedo-Rondon EO, Ferket PR, Havenstein GB. Understanding long bone development in broilers and turkeys. Avian and Poultry Biology Reviews. 2006; 17: 77-88
- Plavnik I, Hurwitz S. Organ weights and body composition in chickens as related to the energy and amino acid requirements: Effects of strain, sex, and age. Poultry Science 1982; 62: 152-163
- Ramadan GS, Moghaieb RE, EL-Ghamry AA, El-Komy EM, Nassar FS, Abdou AM, Ghaly MM, Stino, FKR. Effect of selection for high live body weight on slaughter performance of broiler breeders. Egyptian Poultry Science Journal 2014; 34: 289-304
- Rath, NC, Huff GR, Huff WE, Balog JM. Factors regulating bone maturity and strength in poultry. Poultry Science 2000; 79: 1024-1032
- Salas C, Ekmay RD, England J, Cerrate S, Coon CN. Determination of chicken body composition measured by dual energy X-Ray absorptiometry. International Journal of Poultry Science 2012; 11: 462-468
- SCAHAW. Scientific Committee on Animal Health and Welfare, The Welfare of Chickens Kept for Meat Production (Broilers). European Commission, Health and Consumer Protection Directorate-General, Brussels, Belgium 2000; SANCO.B.3/AH/R15/2000
- Schmidt CJ, Persia ME, Feierstein E, Kingham B, Saylor WW. Comparison of a modern broiler line and a heritage line unselected since the 1950s. Poultry Science 2009; 88: 2610-2619
- Schreiweis MA, Orban JI, Ledur MC, Moddy DE, Hester PY. Effects of ovulatory and egg laying cycle on bone mineral density and content of live White Leghorns as assessed by Dual-Energy X-Ray Absorptiometry. Poultry Science 2004; 83: 1011-1019
- Schreiweis MA, Orban JI, Ledur MC, Moody DE, Hester PY. Validation of dual-energy X-ray absorptiometry in live White Leghorns. Poultry Science 2005; 84: 91-99
- Shields SJ, Garner JP, Mench JA. Dustbathing by broiler chickens: a comparison of preference for four different substrates. Applied Animal Behaviour Science 2004; 87: 69-82
- Shim MY, Karnuah AB, Mitchell AD, Anthony NB, Aggrey SE. The effects of growth rate on leg morphology and tibia breaking strength, mineral density, mineral content, and bone ash in broilers. Poultry Science 2012; 91: 1790-1795
- Simsek UG, Dalkilic B, Ciftci M, Cerci IH, Bahsi M. Effects of enriched housing design on broiler performance, welfare, chicken meat composition and serum cholesterol. Acta Veterinaria Brunensis 2009; 78: 67-74
- Sirri F, Castellini C, Bianchi M, Petracci M, Meluzzi A, Franchini A. Effect of fast-, medium- and slow-growing strains on meat quality of chickens reared under the organic farming method. Animal 2011; 5: 312-319
- SPSS. Statistical Package in Social Sciences for Windows. Statistical Innovations Inc., Chicago, USA 2010
- Talaty PN, Katanbaf MN and Hester PY. Life cycle changes in bone mineralization and bone size traits of commercial broilers. Poultry Science 2009; 88: 1070-1077
- Talaty PN, Katanbaf MN, Hester PY. Bone mineralization in male commercial broilers and its relationship to gait score. Poultry Science 2010; 89: 342-348
- Tavárez MA, de los Santos FS. Impact of genetics and breeding on broiler production performance: a look into the past, present, and future of the industry. Animal Frontiers 2016; 6: 37-41
- Tüzün CG, Aktan S. Residual feed consumption in poultry. Süleyman Demirel University Journal of the Faculty of Agriculture 2012; 7: 115-123
- Watkins KL, Southern LL. Effect of dietary sodium zeolite A and graded levels of calcium and phosphorous on growth, plasma, and tibia characteristics of chicks. Poultry Science 1992; 71: 1048-1058
- Williams B, Waddington DD, Murray H and Farquharson C. Bone strength during growth: Influence of growth rate on cortical porosity and mineralization. Calcified Tissue International 2004; 74: 236-245
- Yan F, Angel R, Ashwell C, Mitchell A, Christman M. Evaluation of the broiler's ability to adapt to an early moderate deficiency of phosphorus and calcium. Poultry Science 2005; 84: 1232-1241