



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e
Tecnologia Avícolas
Brasil

Sereno, MJ; Ziech, RE; Druziani, JT; Pereira, JG; Bersot, LS
Antimicrobial Susceptibility and Biofilm Production by Salmonella sp. Strains Isolated from
Frozen Poultry Carcasses
Revista Brasileira de Ciência Avícola, vol. 19, núm. 1, enero-marzo, 2017, pp. 103-107
Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179750697014>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



■ Author(s)

Sereno MJ^I
Ziech RE^{II}
Druziani JT^I
Pereira JG^{III}
Bersot LS^I

^I Universidade Federal do Paraná, Setor Palotina

^{II} Universidade Federal de Santa Maria

^{III} Universidade Federal do Pampa (UNIPAM-PA) - Campus Uruguaiana

■ Mail Address

Corresponding author e-mail address
Mallu Jagnow Sereno
Rua Pioneiro, 2153, Jardim Dallas, Palotina –
PR, Brasil. Zip code: 85950-000.
Tel: +55 (44) 3211-8513
Email: mallusereno@gmail.com

■ Keywords

Biofilm formation; food pathogens;
multidrug-resistance; poultry meat.

Antimicrobial Susceptibility and Biofilm Production by *Salmonella* sp. Strains Isolated from Frozen Poultry Carcasses

ABSTRACT

The objectives of this study were to evaluate the antimicrobial resistance and the biofilm-producing ability of *Salmonella* sp. strains isolated from frozen poultry carcasses. Antimicrobial susceptibility was tested by the disk-diffusion method. Biofilm-producing ability was determined in 96-well polystyrene microplates stained with crystal violet at 1%. Out of the 22 strains tested, all were multiresistant, that is, resistant to more than three antimicrobial classes, and 72.7% were able to form biofilms. The highest resistance rates obtained were against sulfonamides, tetracycline, and quinolones. On the other hand, 100% of the strains were sensitive to chloramphenicol. According to the rate of biofilm formation, 3 (13.6%) and 13 (59.1%) strains were classified as moderate and weak biofilm-producers, respectively, and 27.3% did not form biofilms. Biofilms increase the tolerance of microorganisms to stress, reducing their sensitivity to disinfectants and antimicrobials; favor equipment corrosion; and act as substrates for the adhesion of bacteria with lower biofilm-producing capacity. The results of the present study stress the importance of cleaning procedures in food processing plants and highlight the public health risks related to the emergence of multiresistant strains.

INTRODUCTION

Animal products, particularly poultry products, have been reported as the main sources of *Salmonella* sp. foodborne diseases (FD) (Dallal *et al.*, 2010). In 2012, about 91,034 cases of salmonellosis were confirmed in humans in Europe (European Food Safety Authority & European Centre for Disease Prevention and Control, 2014), and it is estimated that, in the US, one million people are annually infected by *Salmonella* sp., with an average of 19,000 hospital admittances and 380 deaths (Scallan *et al.*, 2011). In Brazil, according to the Health Surveillance Agency of the Ministry of Health, between 2000 and 2014, there were 9,719 FD outbreaks, and *Salmonella* sp. was the most frequent microorganism associated with these outbreaks. This microorganism was identified as the etiological agent of 1,564 outbreaks in the period (Brasil, 2014).

Salmonellosis is generally clinically manifested by to self-limiting gastroenteritis, and does not require treatment in healthy patients. Treatment is only required in cases of immunosuppressed patients, neonates, elderly patients, or those affected by septicemia, as well as in patients with typhoid fever or enteric fever caused by *S. Typhi* or *S. Paratyphi*, respectively. However, the routine and indiscriminate use of antibiotics have favored the emergence of resistant bacterial populations (Brasil, 2012), increasing the possibility of occurrence of treatment failures, and limiting therapeutic options both in human and veterinary medicine, when treatment is necessary (Souza *et al.*, 2010a).



Microorganisms may be naturally resistant to a given antimicrobial agent or a specific class of antimicrobials due to the molecular structure of the compound, but resistance may also be acquired. Acquired resistance is a specific property of a bacterial strain and may result from the acquisition of resistance genes, which are frequently located in mobile genetic elements, such as plasmids, transposons, gene cassettes inserted in integrons, genomic islands, or mutations in chromosomal genes (Kadlec *et al.*, 2012).

Public health impacts may be more severe when microorganisms, in addition of being resistant to antimicrobials, also have the ability to form biofilms. Biofilms are defined as complex microbial communities embedded in a self-produced extracellular polymeric matrix that attach to surfaces and are the predominant mode of microbial growth in nature (Steenackers *et al.*, 2012). Biofilm formation begins with the adhesion of organic or inorganic molecules to surfaces forming a conditioning film. The reversible phase begins with Van der Waals force and electrostatic attraction of planktonic cells to the preformed substrate and the irreversible phase is the result of the extracellular polymeric matrix production, strengthening the bonds between the bacteria and the surface (Oliveira *et al.*, 2010).

Biofilms on equipment and tools used in food processing are reservoirs of pathogenic and spoilage microorganisms, increasing the risk of microbial contamination in food processing plants (Shi & Zhu, 2009; Xu *et al.*, 2010; Hasegawa *et al.*, 2011; Wang *et al.*, 2013). Bacteria in biofilms are protected from damaging environmental agents, such as disinfectants. Consequently, they are extremely difficult to eliminate, contributing for the resistance and persistence of these bacteria in these sites (Shi & Zhu, 2009).

The objective of the present study was to evaluate the susceptibility pattern of *Salmonella* sp. strains isolated from frozen poultry carcasses to antimicrobials, and to test their ability to produce biofilms.

MATERIAL AND METHODS

The *Salmonella* sp. strains tested in the present study were obtained from two previous studies on the prevalence of this microorganism in 80 poultry carcasses sold in the western region of the state of Paraná, Brazil (Sereno *et al.*, 2012; Druziani *et al.*, 2013). In those studies, only frozen carcasses with their packages intact, bearing the Federal Inspection Service stamp, and before the expiration date were acquired.

Samples were plated on selective and differential media, and typical *Salmonella* colonies were submitted to phenotypical and serological confirmation. Out of each of the 22 positive carcasses obtained by Sereno *et al.* (2012) and Druziani *et al.* (2013), a typical *Salmonella* colony was randomly selected and subjected to the antimicrobial susceptibility test and the evaluation of biofilm-producing ability.

Antimicrobial susceptibility was tested by the disk diffusion method, according to the recommendations of the National Committee for Clinical Laboratory Standards (Clinical and Laboratory Standards Institute, 2011), against 11 agents of nine different classes. (1) aminoglycosides: streptomycin (STR - 10 µg); (2) quinolones: nalidixic acid (NAL - 30 µg) and enrofloxacin (ENR - 5 µg); (3) cephalosporins: Cephalotin (CF - 30 µg) and ceftiofur (CFT - 30 µg); (4) phenicols: chloramphenicol (C - 30 µg); (5) sulfonamides: sulfonamide (SSS - 300 µg); (6) penicillins: ampicillin (AMP - 30 µg); (7) monobactams: aztreonam (AZT - 30 µg); (8) tetracyclines: tetracycline (TET - 30 µg); (9) sulfonamides and pyrimidines: sulfamethoxazole-trimethoprim (SXT - 23.75 µg/1.25). For the interpretation of results, strains that show halos in the resistance and intermediate zones were considered resistant. Strains that were resistant to at least three antimicrobial classes were considered multiresistant (Ngoi & Thong, 2013).

The biofilm-producing test applied the methodology described by Stepanović *et al.* (2000), with some adaptations according to Steenackers *et al.* (2012). Each strain was diluted to 10⁸ CFU/mL (0.5 in the MacFarland scale) using Luria-Bertani LB broth (Difco™), and 200 µL were cultured in four wells of the 96-well flat-bottom polystyrene microplate (Nest®). A total of 88 wells were used to test 22 strains; the other 4 wells received the *Salmonella* Typhimurim ATCC 14028 as positive control (Oliveira *et al.*, 2014), and 4 wells received the negative control (non-inoculated culture medium). The plates containing *Salmonella* sp. strains and controls were incubated at 35°C for 96h. The plate was washed three times with phosphate buffered saline (PBS pH 7.2) and stained with crystal violet at 1% for 15 minutes. After washing three times with distilled water and drying at room temperature, absorbance was read in a Polaris (Celer®) microplate reader at 492 nm wave length.

In order to determine absorbance results, optical density of each sample (OD) was compared with the three standard deviations above the mean OD of the negative control (OD₋); to determine biofilm-producing



rate, the following classification was used: non-producer ($OD \leq OD_c$), weak biofilm-producer ($OD_c < OD \leq 2 \times OD_c$), moderate biofilm-producer ($2 \times OD_c < OD \leq 4 \times OD_c$), and strong biofilm-producer ($4 \times OD_c < OD$). All tests were carried out two times and the results were averaged.

RESULTS AND DISCUSSION

All 22 strains tested were considered multiresistant in the antimicrobial susceptibility test, with 100% resistant to quinolones, sulfonamide, and tetracycline; 81.8% resistant to cephalosporins; and 77.3% resistant to penicillins. Only 36.4% strains were resistant to aminoglycosides, and 13.6% to monobactams and sulfonamides associated with pyrimidines (Table 1). Among these agents, quinolones, third-generation cephalosporins, penicillins, and monobactams are considered critically important antimicrobials in human medicine, according to the World Health Organization (World Health Organization, 2012).

Table 1 – Sensitivity and resistance of each of the 22 strains of *Salmonella* sp. to each antimicrobial tested.

Antimicrobial Class	Antimicrobial Tested	Strains Tested	
		Resistant (n/%)	Sensitive (n/%)
Aminoglycosides	STR	8/36.4	14/63.6
Quinolones	NAL	22/100	0/0
	ENR	2/9.1	20/90.9
Cephalosporins	CF	18/81.8	4/18.2
	CFT	15/68.2	7/31.8
Phenicol	C	0/0	22/100
Sulfonamides	SSS	22/100	0/0
Penicillins	AMP	17/77.3	5/22.7
Monobactams	AZT	3/13.6	19/86.4
Tetracyclines	TET	22/100	0/0
Sulfonamides/ Pyrimidines	SXT	3/13.6	19/86.4

STR – Streptomycin; NAL – Nalidixic acid; ENR – Enrofloxacin; CF – Cephalotin; CFT – Ceftiofur; C – Chloramphenicol; SSS – Sulfonamide; AMP – Ampicillin; AZT – Aztreonam; TET – Tetracycline; SXT – Sulfamethoxazole-trimetoprim.

The obtained results are in agreement with those of Cortez *et al.* (2006), Dallal *et al.* (2010), and Wang *et al.* (2013), who reported tetracycline resistance rates of 72.4%, 69%, and 70%, respectively. Tetracycline was widely used in Brazil as a prophylactic agent in poultry feed until it was banned (Brasil, 2009). The high rates of tetracycline resistance observed may be related to the transference of several *tet* genes that confer resistance to *Salmonella* sp. (Pezzella *et al.*, 2004). Other results presented by Cortez *et al.* (2006), in Brazil, and Wang *et al.* (2013) in China, showed that

86.2% and 78% of the isolated strains, respectively, were also resistant to ampicillin, as also detected in the present study. Bacci *et al.* (2012) in Italy, observed that only 33.3% of the *Salmonella* strains isolated from chicken carcasses were resistant to ampicillin.

Among the tested quinolones, 100% of the strains showed resistance to nalidixic acid, and among cephalosporins, 68.2% showed resistance to ceftiofur (Table 1). Ceftiofur is a third-generation cephalosporin, and these high levels of resistance represent a public health concern, as after the emergence of strains resistant to chloramphenicol around the 1990s, quinolones and third-generation cephalosporins started to be the antibiotics of choice for the treatment of salmonellosis in humans (Pokharel *et al.*, 2006; Souza *et al.*, 2010b; Lopes 2014).

On the other hand, none of the strains was resistant to chloramphenicol (Table 1), as previously reported by Cardoso *et al.* (2006), Lima *et al.* (2009), Begum *et al.* (2010), and Galdino *et al.* (2013). These authors also did not find any resistance to norfloxacin in *Salmonella* strains isolated from chicken products. According to Lima *et al.* (2009), the sensitivity to chloramphenicol is partially due to the fact that the use of this antimicrobial, both as growth promoter and therapeutic agent, has been banned from animal production since the 1970s, thereby reducing the exposure of bacteria to this active principle.

As for the assessment of biofilm production, 72.7% of the strains tested were able to form biofilms on polystyrene microplates. Ziech *et al.* (2016a) reported that 100% of *Salmonella* sp. strains isolated in poultry slaughterhouses and processing plants in Brazil were able to produce biofilm. Oliveira *et al.* (2014), in Brazil, and Solano *et al.* (2002), in Spain, found rates of 98.3% and 97%, respectively. According to the classification by biofilm production rate, in the present study, 13.6% isolates were moderate producers, 59.1% were weak biofilm-producers, and 27.3% did not form biofilm (Table 2).


Table 2 – Classification of the 22 strains of *Salmonella* sp. according to the rate of biofilm production.

Rate of biofilm production	Strains tested	
	Nº.	%
No formation	6	27,3
Weak	13	59.1
Moderate	3	13.6
Strong	0	0
Total of strains	22	100



According to Gilbert *et al.* (2002), the resistance of microbial biofilms to a wide variety of antimicrobial agents is related with the organization of bacterial cells inside the polymer matrix, which hinders the penetration of antimicrobial agents into the biofilm. In addition, cells under stress, showing slow growth rate and undergoing restricted access to nutrients, may express phenotypes related to mechanisms of resistance to antimicrobials. This is due to the exchange of genetic material, especially of genes that encode resistance, among bacteria within the biofilm, where bacterial multiplication is slow, but conjugation rates are fast (Ghigo, 2001).

In the present study, the evaluated strains showed multiresistance to antibiotics, and most were able to form biofilms to some degree. Wang *et al.* (2013), analyzing *Salmonella* sp. isolated from chicken carcasses and from contact surfaces of processing plants, did not find any positive correlation between biofilm formation and antimicrobial resistance. In the present study, the most frequent profile of the isolates was weak biofilm formation and resistance to quinolones, sulfonamides, tetracyclines, cephalosporin, and penicillins (Figure 1). On the other hand, Ziech *et al.* (2016b), evaluating *Salmonella* sp. strains isolated in broiler processing plants, detected 86% multiresistant strains, and the most frequent profile was weak biofilm formation, production of wide-spectrum beta-lactamase, and resistance to beta-lactams, quinolones, and tetracyclines.



QUI - Quinolones; SSS - Sulfonamides; TET - Tetracyclines; CEF - Cephalosporins; PEN - Penicillins; AMINO - Aminoglycosides; MON - Monobactams; SSS/PIR - Sulfonamides/Pyrimidines

Figure 1 – Distribution of resistance pattern and ability to produce biofilm in the 22 strains of *Salmonella* sp. that were tested.

CONCLUSIONS

Data obtained in the present study suggest that poultry carcasses, particularly those found in retail stores, show a high prevalence of *Salmonella* sp. resistant to the antimicrobials commonly used in human medicine to control severe infections. These results point out to an alarming public health situation.

In addition to the fact that the strains tested showed high resistance to antimicrobials, a high percentage presented the ability to form biofilms. Biofilms increase the tolerance of microorganisms to stress, reducing their sensitivity to disinfectants and antimicrobials; favor equipment corrosion; and aid the adhesion of bacteria that are less able to form biofilms, as these act as substrates. Consequently, biofilms contribute to the persistence of resistant microorganisms both in the food industry environment and in processed foods, mainly due to cross contamination.

ACKNOWLEDGMENT

The authors Bersot thank CNPq and Sereno and Druziani CAPES for the financial support.

REFERENCES

- Bacci C, Boni E, Alpighiani I, Lanzoni E, Bonardi S, Brindani F. Phenotypic and genotypic features of antibiotic resistance in *Salmonella enterica* isolated from chicken meat and chicken and quail carcasses. *International Journal of Food Microbiology* 2012;160(1):16-23.
- Begum K, Reza TA, Haque M, Hossain A, Hassan FMK, Hasan SN, et al. Isolation, identification and antibiotic resistance pattern of *Salmonella* sp. from chicken eggs, intestines and environmental samples. *Bangladesh Pharmaceutical Journal* 2010;13(1):23-27.
- Brasil. Agência Nacional de Vigilância Sanitária. Programa Nacional de Monitoramento de Prevalência e da Resistência Bacteriana em Frangos - PREBAF. Monitoramento de prevalência e do perfil de susceptibilidade aos antimicrobianos em enterococos e salmonelas isolados de carcaças de frango congeladas comercializadas no Brasil. Brasília, DF: ANVISA; 2012. p.1-171.
- Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa nº 26, de 9 de julho de 2009. Regulamento técnico para a fabricação, o controle de qualidade, a comercialização e o emprego de produtos antimicrobianos de uso veterinário. *Diário Oficial da República Federativa do Brasil*, Brasília, DF; 2009.
- Brasil. Vigilância Epidemiológica das Doenças Transmitidas por Alimentos (VE-DTA). Dados de surtos de DTA no Brasil, 2000-2014 [cited 2014 Nov 19]. Available from: http://www.anrbrasil.org.br/new/pdfs/2014/3_PAINEL_1_ApresentacaoRejaneAlvesVigilanciaEpidemiologica-VE-DTA-Agosto_2014_PDF.pdf.
- Cardoso MO, Ribeiro AR, Santos LR, Pilotto F, Moraes HLS, Salle CTP, et al. Antibiotic resistance in *Salmonella* Enteritidis isolated from broiler carcasses. *Brazilian Journal of Microbiology* 2006;37(1):368-371.
- Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-First Informational



- Supplement. CLSI document M100-S21. Wayne: Clinical and Laboratory Standards Institute; 2011. p.1-172.
- Cortez ALL, Carvalho ACFB, Ikuno AA, Burger KP, Vidal-Martins AMC. Resistência antimicrobiana de cepas de *Salmonella* sp. isoladas de abatedouros de aves. Arquivos do Instituto Biológico 2006;73(2):157-163.
- Dallal MMS, Doyle MP, Rezadehbashi M, Dabiri H, Sanaei M, Modarresi S, et al. Prevalence and antimicrobial resistance profiles of *Salmonella* serotypes, *Campylobacter* and *Yersinia* spp. isolated from retail chicken and beef, Tehran, Iran. Food Control 2010;21(1):388-392.
- Druziani JT, Sereno MJ, Lampugnani C, Pereira JG, Bersot, LS. Prevalência de *Salmonella* em carcaças de frango na região Oeste do Estado do Paraná e perfil de resistência a antimicrobianos. Resumos do 21º Evento da Iniciação científica; 6º Evento de Iniciação em Desenvolvimento Tecnológico e Inovação; 2013; Curitiba, Paraná. Brasil: Fundação Araucária; 2013. p.513.
- European Food Safety Authority & European Centre for Disease Prevention and Control. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2012. EFSA Journal 2014;12(2):1-312.
- Galdino VMCA, Melo RT, Oliveira RP, Mendonça EP, Nalevaiko PT, Rossi DA. Virulência de *Salmonella* sp. de origem avícola e resistência a antimicrobianos. Bioscience Journal 2013;29(4):932-939.
- Ghigo JM. Natural conjugative plasmids induce bacterial biofilm development. Nature 2001;412(6845):442-445.
- Gilbert P, Allison DG, McBain AJ. Biofilms in vitro and in vivo: do singular mechanisms imply. Journal of Applied Microbiology Symposium Supplement 2002;92(1):98-110.
- Hasegawa A, Hara-KudoY, Kumagai S. Survival of *Salmonella* strains differing in their biofilm-formation capability upon exposure to hydrochloric and acetic acid and to high salt. Journal of Veterinary Medical Science 2011;73(1):1163-1168.
- Kadlec K, Michael GB, Fessler AT, Schwarz S. Molecular mechanisms of antibiotic resistances. Antibiotics in Food Chain. Proceedings of the Max Rubner Conference; 2012; Karlsruhe. Netherlands. p.16.
- Lima ET, Andreatti Filho RL, Pinto JPAN. Perfil de susceptibilidade antimicrobiana de sorotipo de *Salmonella* isolados de produtos avícolas. Veterinária e Zootecnia 2009;16(2):394-400.
- Lopes GV. Caracterização de determinantes de resistência a antimicrobianos em isolados de *Salmonella enterica* subsp. *enterica* provenientes da cadeia produtiva de suínos no sul do Brasil [tese]. Porto Alegre (RS): Universidade Federal do Rio Grande do Sul; 2014.
- Ngoi ST, Thong KL. Molecular characterization showed limited genetic diversity among *Salmonella* Enteritidis isolated from humans and animals in Malaysia. Diagnostic Microbiology and Infectious Disease 2013;77(1):304 - 311.
- Oliveira DCV, Fernandes Júnior A, Kaneno R, Silva MG, Araújo Júnior JP, Silva NCC, et al. Ability of *Salmonella* sp. to produce biofilm is dependent on temperature and surface material. Foodborne Pathogens and Disease 2014;1(1):1-7.
- Oliveira MM, Brugnera DF, Piccoli RH. Biofilmes microbianos na indústria de alimentos: uma revisão. Revista do Instituto Adolfo Lutz 2010;69(3):277-284.
- Pezzella C, Ricci A, Digiannatale E, Luzzi I, Carattoli A. Tetracycline and streptomycin resistance genes, transposons and plasmids in *Salmonella enterica* isolates from animals in Italy. Antimicrob Agents Chemother 2004;48(3):903 - 908.
- Pokharel BM, Koirala J, Dahal RK, Mishra SK, Khadga PKL, Tuladhar NR. Multidrug-resistant and extended-spectrum beta-lactamase (ESBL)-producing *Salmonella enterica* (serotypes Typhi and Paratyphi A) from blood isolates in Nepal: surveillance of resistance and a search for newer alternatives. International Journal of Infectious Diseases 2006;10(6):434-438.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL. Foodborne illness acquired in the United States e major pathogens. Emerging Infectious Diseases 2011;17(1):7-15.
- Sereno MJ, Druziani JT, Gallego J, Bonatto CB, Bersot LS. Prevalência de *Salmonella* spp. em carcaças de frango na região Oeste do Estado do Paraná e perfil de resistência a antimicrobianos. Resumos do 20º Evento da Iniciação científica; 4º Evento de Iniciação em Desenvolvimento Tecnológico e Inovação; 2013; Curitiba, Paraná. Brasil: Fundação Araucária; 2011. p.505.
- Shi X, Zhu X. Biofilm formation and food safety in food industries. Food Science and Technology 2009;20(1):407-413.
- Solano C, Garcia B, Valle J, Berasain C, Ghigo JM, Gamazo C, Lasa I. Genetic analysis of *Salmonella enteritidis* biofilm formation: Critical role of cellulose. Molecular Microbiology 2002;43(1):793-808.
- Souza RB, Ferrari RG, Magnani M, Kottwitz LBM, Alcocer I, Tognim MCB, et al. Ciprofloxacin susceptibility reduction of *Salmonella* strains isolated from outbreaks. Brazilian Journal of Microbiology 2010b;41(1):497-500.
- Souza RB, Magnani M, Oliveira TCRM. Mecanismos de resistência às quinolonas em *Salmonella* sp. Semina Ciências Agrárias 2010a;31(2):413-428.
- Steenackers H, Hermans K, Vanderleyden J, Keersmaecker SCJ. *Salmonella* biofilms: An overview on occurrence, structure, regulation and eradication. Food Research International 2012;45:502-531.
- Stepanovic S, Vukovic D, Dakic I, Savic B, Svabic-Vlahovic M. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. Journal of Microbiological Methods 2000;40(1):175-179.
- Wang H, Ye K, Wei X, Cao J, Xu X, Zhou G. Occurrence, antimicrobial resistance and biofilm formation of *Salmonella* isolates from a chicken slaughter plant in China. Food Control 2013;33(1):78-84.
- WHO - World Health Organization. Critically important antimicrobials for human medicine [cited 2015 Jan 5]. World Health Organization; 2012. Available from: http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485_eng.pdf.
- Xu H, Lee HY, Ahn J. Growth and virulence properties of biofilm-forming *Salmonella enterica* serovar Typhimurium under different acidic conditions. Applied Environmental Microbiology 2010;76(1):910-917.
- Ziech RE, Lampugnani C, Perin AP, Sereno MJ, Sfacciotte RAP, Viana C, et al. Multidrug resistance and ESBL-producing *Salmonella* sp. isolated from broiler processing plants. Brazilian Journal of Microbiology 2016b;47:191-195.
- Ziech RE, Perin AP, Lampugnani C, Sereno MJ, Viana C, Soares VM, et al. Biofilm-producing ability and tolerance to industrial sanitizers in *Salmonella* sp. isolated from Brazilian poultry processing plants. Food Science and Technology 2016a;68:85-90.