



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e
Tecnologia Avícolas
Brasil

Zendehdel, M; Tirgari, F; Shohre, B; Deldar, H; Hassanpour, S
Involvement of Gaba and Cannabinoid Receptors in Central Food Intake Regulation in
Neonatal Layer Chicks: Role of CB1 and GABAA Receptors
Revista Brasileira de Ciência Avícola, vol. 19, núm. 2, abril-junio, 2017, pp. 221-229
Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179751953007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

■ Author(s)

Zendehtdel M^I
Tirgari F^{II}
Shohre B^{II}
Deldar H^{II}
Hassanpour S^{III}

^I Department of Basic Sciences, Faculty of Veterinary Medicine, University of Tehran, 14155-6453, Tehran, Iran

^{II} Department of Animal Science, Sari Agricultural Sciences & Natural Resources University, Sari, Iran

^{III} Department of Basic Sciences, Faculty of Veterinary Medicine, Science and Research Branch, Islamic Azad University, Tehran, Iran

■ Mail Address

Corresponding author e-mail address
Morteza Zendehtdel
Department of Basic Sciences, Faculty of Veterinary Medicine, University of Tehran, 14155-6453, Tehran, Iran
Tel: +98-21-61117186
Email: zendehtdel@ut.ac.ir

■ Keywords

Cannabinoidergic, GABAergic, Food intake, Layer-type chicken.

Involvement of Gaba and Cannabinoid Receptors in Central Food Intake Regulation in Neonatal Layer Chicks: Role of CB₁ and GABA_A Receptors

ABSTRACT

Feeding behavior is regulated via a complex network which interacts via diverse signals from central and peripheral tissues. Endocannabinoids modulate release of GABA in a variety of regions of the central nervous system. Endocannabinoids and GABAergic system have an important role in the central regulation of appetite. Thus, the present study examines the possible interaction of central cannabinoidergic and GABAergic systems on food intake in 3-h food-deprived (FD₃) neonatal layer-type chicks. The results of this study showed that intracerebroventricular (ICV) injection of 2-AG (2-Arachidonoylglycerol, selective CB₁ receptors agonist, 2μg) significantly increased food intake and this effect of 2-AG was attenuated by Picrotoxin (GABA_A antagonist, 0.5μg) (P<0.001); but 21ng CGP54626 (GABA_B antagonist) had no effect (p>0.05). Also, hyperphagic effect of CB₆₅ (CB₂ receptors agonist, 1.25μg) was not affected by Picrotoxin or CGP54626 (p>0.05). Moreover, the food intake of chicks was significantly increased by ICV injection of GABA_A agonist (Gaboxadol, 0.2 μg) and SR141716A (CB₁ receptors antagonist, 6.25μg) significantly decreased Gaboxadol-induced hyperphagia (P<0.001) but CB₂ receptors antagonist (AM630, 1.25μg) had no effect. In contrast, co-injection of SR141716A or AM630 with GABA_B agonist (baclofen, 0.2μg) had no effect on the hyperphagia induced by baclofen (p>0.05). These data showed there might be an interaction between central cannabinoidergic and GABAergic systems via CB₁ and GABA_A receptors in control of food intake in neonatal layer chicks.

INTRODUCTION

Feeding behavior is a complex physiologic phenomenon which interacts via various signals from central and peripheral tissues. In the central nervous system (CNS), several neurotransmitters interact by a wide distributed neurological network on food intake regulation (Zendehtdel *et al.*, 2014; Honda, 2016). γ-aminobutyric acid (GABA) is one of the most abundant neurotransmitter in mammalian brain. The GABAergic is known as an inhibitory neurotransmitter in the CNS (Bonaventura *et al.*, 2012). GABA exerts its effect via two distinct receptors including the bicuculline-sensitive GABA_A receptor and the bicuculline-insensitive GABA_B receptors (Stratford and Wirtshafter, 2013). The GABA_A and GABA_C receptors are part of a macromolecular complex coupled to a Cl⁻ ionophore while GABA_B is metabotropic receptor belong to G-protein coupled receptors (GPCRs) (Jonaidi *et al.*, 2012).

The appetite-stimulating effects of marijuana (*Cannabis sativa*) are known in science for the past decades. Animal tissues express at least two cannabinoidergic (CBergic) receptors: cannabinoid type 1 (CB₁) and type 2 (CB₂) receptors which belong to the GPCRs (Kangas



et al., 2013). CB₁ receptors mainly expressed in the presynaptic terminals of the brain (Sharkey *et al.*, 2014) and CB₂ receptors formerly exist on cells and organs of the immune system in the peripheral nervous system (PNS) but they also expressed in the CNS (Onaivi *et al.*, 2008). The endocannabinoids (ECBs) mediates many physiological functions like pain relief, motor control, learning and memory formation as well as appetite regulation (D'Addario *et al.*, 2014).

GABA and CBs involved control of food intake (Volkow *et al.*, 2011). It appears GABA has orexigenic effect in domestic fowls (Jonaidi *et al.*, 2012). ICV injection of Muscimol (a GABA_A agonist) and Baclofen (a GABA_B agonist) increased food consumption in rats (Ebenezer, 1990), turkeys (Denbow, 1991), meat-type (broiler) and layer-type chicks (hens) (Jonaidi *et al.*, 2002). Both CB₁ and CB₂ receptors have a role in food intake regulation in the rat (Chen *et al.*, 2006). Scarce information exists about the role of ECBs on feeding behavior in domestic fowl. Reports claim just CB₁ receptors have a regulatory role on appetite in broilers (*et al.*, 2011; Novoseletsky *et al.*, 2011). Recent reports revealed both CB₁ and CB₂ receptors have hyperphagic effect in neonatal layer-type chicken (Alizadeh *et al.*, 2015; Hassanpour *et al.*, 2015).

There is central evidence mechanisms for food intake regulation are different between mammalian and birds (Zendehdel & Hassanpour 2014). Comparative physiological studies suggested there are differences award pathways between the meat-type and layer-type chicks (Alizadeh *et al.*, 2015). For example, food intake mediates via both CB₁ and CB₂ receptors in layer-type chicken (Alizadeh *et al.*, 2015) but only CB₂ receptors interact on feeding in meat-type chicken (Emadi *et al.*, 2011; Novoseletsky *et al.*, 2011). For instance, ICV injection of Baclofen had no effect on food intake in broilers (Jonaidi *et al.*, 2002; Tajalli *et al.*, 2006; Zendehdel *et al.*, 2008, 2009).

During the last several years, GABA and ECBs have been the subject of an increasing number of behavioral studies. Several types of researches are done to find the mediatory effect of neurotransmitters on feeding behavior in mammals but aspects of food intake regulation in avian are still unclear (Zendehdel & Hassanpour, 2014). ECBs modulate the release of GABA in a variety of regions of the CNS, including the hippocampus, basal ganglia, cerebellum, and brainstem. Anatomical investigations have shown high levels of CB₁ receptor immune-reactivity and mRNA associated with GABAergic neurons (Irving *et al.*, 2002). In the brain, CB₁ receptors is predominantly expressed presynaptically and modulating the release

of neurotransmitters, such as GABA, dopamine, noradrenaline, glutamate and serotonin (Cota *et al.*, 2003). To the best of our knowledge, no report exists on the interconnection of the CBergic and GABAergic systems on food intake regulation and energy balance in neonatal layer type chicken. So, the aim of this study was to investigate the possible involvement of the CBergic and GABAergic systems on feeding behavior in 3 h food deprived (FD₃) neonatal layer type chicken.

MATERIALS AND METHODS

Animals

In this study, a total 384 one-day-old layer-type chickens (Hy-Line) purchased from a local hatchery (Morghak Co. Iran). Birds were kept as flocks for two days then randomly transferred into individual cages at a temperature of 30 ± 1°C with 50 ± 2 percent humidity (Olanrewaju *et al.*, 2006). A commercial diet provided during the study containing 21 % crude protein and 2850 kcal/kg of metabolizable energy (Chineh Co. Iran). All birds received *ad libitum* food and fresh water during the study. Just three h prior the ICV injections, chicken, were food deprived (FD₃) but had free access to water. The injections were applied to all birds at five days of age. Animal handling and experimental procedures were performed according to the Guide for the Care and Use of Laboratory animals by the National Institutes of Health (USA) and the current laws of the Iranian government for animal care.

Experimental Drugs

Gaboxadol (GABA_A agonist), Baclofen (GABA_B agonist), Picrotoxin (GABA_A antagonist), CGP54626 (GABA_B antagonist), 2-AG (2-Arachidonoylglycerol, a selective CB₁ receptors agonist), CB65 (a selective CB₂ receptors agonist), SR141716A (a selective CB₁ receptors antagonist), AM630 (a selective CB₂ receptors antagonist) and Evans Blue purchased from Sigma Co. (Sigma, USA). Drugs except 2-AG at first dissolved in absolute dimethyl sulfoxide (DMSO) then diluted with 0.85% saline containing Evans blue at a ratio of 1/250. 2-AG solved in 0.1 % Evans blue solution which was prepared in 0.85 % saline. DMSO with this ratio does not have a cytotoxic effect (Blevins *et al.*, 2002; Qi *et al.*, 2008).

ICV injection procedures

In this study, eight experiments designed to investigate interconnection of CBergic and GABAergic systems on cumulative food intake in neonatal layer-type birds (each includes four groups within 12



replicates in each group). Before each treatment, the chicks were weighed and based on their body weight allocated into groups so the average weight between treatment groups was as uniform as possible. ICV injection was applied using a microsyringe (Hamilton, Switzerland) without anesthesia according to the technique previously described by Davis *et al.*, (1979) and Furuse *et al.*, (1997) in which the head of the birds was held with an acrylic device while the bill holder was 45° and calvarium parallel to the surface of table (Van Tienhoven and Juhasz, 1962). In a plate, a hole was drilled in the skull over the right lateral ventricle and immediately overlaid through this plate. A microsyringe inserted into the right ventricle via the hole and tip of the needle penetrated 4 mm beneath the skin of the skull. It is revealed that there is no injection-induced physiological stress using this method in neonatal chicks (Saito *et al.*, 2005). Each bird received an ICV injection (with vehicle or drug solution) in a volume of 10 µL. Right away after the injection, FD₃ birds returned to their individual cages and offered fresh water and food (pre-weighed). Cumulative food intake (gr) was measured at 30, 60 and 120 min after the injection. Food consumption was calculated as a percentage of body weight to minimize the effect of body weight on the amount of food intake. Each bird was just used once in each experimental group. At the end, accuracy of placement of the injection in the ventricle was verified by the presence of Evans Blue followed by slicing the frozen brain tissue. In each group, 12 birds received the injection, but just data of those individuals where the dye was present in their lateral ventricle (12 chickens per group), was used for the analysis. All experimental procedures were done from 8:00 A.M. until 3:30 P.M. Also, the time course for feeding behavior was selected by previous studies (Jonaidi *et al.*, 2012; Zendehdel & Hassanpour 2014; Alimohammadi *et al.*, 2015; Alizadeh *et al.*, 2015; Hassanpour *et al.*, 2015).

Feeding experiments

In this study, eight experiments were designed, each with four treatment groups: A, B, C and D groups (n=48 in each). In experiment 1, chickens received intracerebroventricular (ICV) injection of (A) saline, (B) 2-AG (2-Arachidonoylglycerol, selective CB₁ receptors agonist, 2µg), (C) Picrotoxin (GABA_A antagonist, 0.5µg) and (D) co-injection of 2-AG+Picrotoxin. In experiment 2, birds injected with (A) saline, (B) 2-AG (2µg), (C) CGP54626 (GABA_B antagonist, 21ng) and (D) co-injection of 2-AG+CGP54626. In experiment 3, animals received: (A) saline, (B) CB65 (selective

CB₂ receptors agonist, 1.25µg), (C) Picrotoxin (0.5µg) and (D) CB65+ Picrotoxin. In experiment four, chicks ICV injected with (A) saline, (B) CB65 (1.25µg), (c)

Table 1 – Treatments procedure in experiments 1-8

Exp. 1	ICV Injection
Treatment groups	
A	CS*
B	2-AG (2µg)
C	Picrotoxin (0.5µg)
D	2-AG (2µg)+ Picrotoxin (0.5µg)
Exp. 2	ICV Injection
Treatment groups	
A	CS*
B	2-AG (2µg)
C	CGP54626 (21ng)
D	2-AG (2µg)+ CGP54626 (21ng)
Exp. 3	ICV Injection
Treatment groups	
A	CS*
B	CB65 (1.25µg)
C	Picrotoxin (0.5µg)
D	CB65 (1.25µg) + Picrotoxin (0.5µg)
Exp. 4	ICV Injection
Treatment groups	
A	CS*
B	CB65 (1.25µg)
C	CGP54626 (21ng)
D	CB65 (1.25µg) +CGP54626 (21ng)
Exp. 5	ICV Injection
Treatment groups	
A	CS *
B	Gaboxadol (0.2µg)
C	SR141716A (6.25µg)
D	Gaboxadol (0.2µg) + SR141716A (6.25µg)
Exp. 6	ICV Injection
Treatment groups	
A	CS *
B	Gaboxadol (0.2µg)
C	AM630 (1.25µg)
D	Gaboxadol (0.2µg)+ AM630 (1.25µg)
Exp. 7	ICV Injection
Treatment groups	
A	CS *
B	baclofen (0.2µg)
C	SR141716A (6.25µg)
D	baclofen (0.2µg) + SR141716A (6.25µg)
Exp. 8	ICV Injection
Treatment groups	
A	CS *
B	baclofen (0.2µg)
C	AM630 (1.25µg)
D	baclofen (0.2µg) + AM630 (1.25µg)

CS: control solution, 2-AG: 2-Arachidonoylglycerol, selective CB₁ receptors agonist, Picrotoxin: GABA_A antagonist, CGP54626: GABA_B antagonist, CB65: selective CB₂ receptors agonist,CGP54626: GABA_B antagonist, Gaboxadol: GABA_A agonist, SR141716A: selective CB₁ receptors antagonist, AM630: selective CB₂ receptors antagonist, Baclofen: GABA_B agonist.



CGP54626 (21ng) and (D) co-administration of CB65+CGP54626. In experiment 5, birds injected with (A) saline, (B) Gaboxadol (GABA_A agonist, 0.2μg), (C) SR141716A (selective CB₁ receptors antagonist, 6.25μg) and (D) Gaboxadol+ SR141716A injected. In experiment 6, ICV injection of (A) saline, (B) Gaboxadol (0.2μg), (C) AM630 (selective CB₂ receptors antagonist, 1.25μg) and (D) co-injection of Gaboxadol+AM630 were applied. In trial 7, avian injected with (A) saline, (B) baclofen (GABA_B agonist, 0.2μg), (C) SR141716A (6.25μg) and (D) baclofen+SR141716A. In experiment 8, (A) saline, (B) baclofen (GABA_B, 0.2μg), (C) AM630 (1.25μg) and (D) co-injection of baclofen+AM630 used. The injection procedure is shown in Table 1. To find the possible interaction between these CBergic and GABAergic systems, effective and sub-effective doses of pharmacologic agents were administered to confront nullifying effects of the agents. In other words, when an effective dose of a system was administered, the sub-effective dose of the other system was considered. These doses of drugs were determined according to the previous studies (Jonaidi *et al.*, 2002; Chen *et al.*, 2006; Emadi *et al.*, 2011; Novoseletsky *et al.*, 2011; Alizadeh *et al.*, 2015; Hassanpour *et al.*, 2015).

Statistical analysis

Data is presented as mean ± SEM (standard error of the mean). Cumulative food intake (as the percentage of body weight) was analyzed by repeated measure two-way analysis of variance (ANOVA) using SPSS 16.0 for Windows (SPSS, Inc., Chicago, IL, USA). For treatment showing a main effect by ANOVA, means were compared by Tukey-Kramer test. $p < 0.05$ was significant differences between treatments.

RESULTS

Effects and interactions of central CBergic and GABAergic systems on cumulative food intake in FD₃ neonatal layer-type bird are shown in Figures 1-8.

In experiment 1, ICV injection of effective dose of CB₁ receptors agonist (2-AG, 2μg) significantly increased food intake (% BW) in comparison with control group [$F(1,42)=67.02$; $p < 0.001$]. Also, the ICV injection of sub effective dose of GABA_A antagonist Picrotoxin (0.5μg) had no effect on food intake compared to control group [$F(1,42)=0.01$; $p > 0.05$]. Co-injection of CB₁ receptors agonist plus GABA_A antagonist (Picrotoxin + 2-AG) significantly decreased CB₁ receptors-induced hyperphagia [Time, $F(2,14)=93.45$, $p < 0.001$; Picrotoxin × 2-AG interaction, $F(1,42)=48.95$, $p < 0.001$] [Fig. 1].




Figure 1 – Effect of ICV injection of 2-AG (2μg), Picrotoxin (0.5μg) and their combination on percent of body weight cumulative food intake in neonatal layer type chickens (n=48). 2-AG: 2-Arachidonoylglycerol, selective CB₁ receptors agonist, Picrotoxin: GABA_A antagonist. Data are expressed as mean ± SEM. F and P value for within and between subject factors are as follows: Time, $F(2,14)=93.45$, $p < 0.001$; 2-AG, $F(1,42)=67.02$, $p < 0.001$; Picrotoxin, $F(1,42)=0.01$, $p > 0.05$; Picrotoxin × 2-AG interaction, $F(1,42)=48.95$, $p < 0.001$. Different letters (a, b and c) indicate significant differences between treatments ($p < 0.001$). The average body weight for each group was as follows: control, 62.14 g; 2-AG, 61.71 g; Picrotoxin, 61.48 g; 2-AG + Picrotoxin, 62.83 g.

In experiment 2, ICV injection of 2-AG, (2μg) had hyperphagic effect compared to control group in neonatal layers [$F(1,38)=39.51$; $p < 0.001$]. Also, ICV injection of sub effective dose of GABA_B antagonist (CGP54626, 21ng) had no effect on food consumption [$F(1,38)=0.62$; $p > 0.05$]. Co-injection of CGP54626 + 2-AG was not able to attenuated 2-AG-induced hyperphagia in neonatal layers [Time, $F(2,69)=112.04$, $p < 0.001$; CGP54626 × 2-AG interaction, $F(1,38)=0.08$, $p > 0.05$] [Fig 2].




Figure 2 – Effect of ICV injection of 2-AG (2μg), CGP54626 (21ng) and their combination on percent of body weight cumulative food intake in neonatal layer type chickens (n=48). 2-AG: 2-Arachidonoylglycerol, selective CB₁ receptors agonist, CGP54626: GABA_B antagonist. Data are expressed as mean ± SEM. F and P value for within and between subject factors are as follows: Time, $F(2,69)=112.04$, $p < 0.001$; 2-AG, $F(1,38)=39.51$, $p < 0.001$; CGP54626, $F(1,38)=0.62$, $p > 0.05$; CGP54626 × 2-AG interaction, $F(1,38)=0.08$, $p > 0.05$. Different letters (a and b) indicate significant differences between treatments ($p < 0.001$). The average body weight for each group was as follows: control, 64.26 g; 2-AG, 63.92 g; CGP54626, 63.19 g; CGP54626 + 2-AG, 62.38 g.

In experiment 3, ICV administration of CB₂ receptors agonist, 1.25μg) significantly increased food



intake compared to control group [$F(1,54)=76.01$; $p<0.001$]. ICV administration of Picrotoxin ($0.5\mu\text{g}$) had no effect on food intake compared to control group [$F(1,54)=0.08$; $p>0.05$]. Additionally, co-injection of Picrotoxin+CB65 was not able to fluctuate hyperphagic effect of CB₂ receptors in neonatal layer-type chicks [Time, $F(2,81)=67.52$, $p<0.001$; Picrotoxin \times CB65, $F(1,54)=1.36$; $p>0.05$] [Fig 3].




Figure 3 – Effect of ICV injection of CB65 ($1.25\mu\text{g}$), Picrotoxin ($0.5\mu\text{g}$) and their combination on percent of body weight cumulative food intake in neonatal layer type chicks ($n=48$). CB65: selective CB₂ receptors agonist, Picrotoxin: GABA_A antagonist. Data are expressed as mean \pm SEM. F and P value for within and between subject factors are as follows: Time, $F(2,81)=67.52$, $p<0.001$; CB65, $F(1,54)=76.01$, $p<0.001$; Picrotoxin, $F(1,54)=0.08$, $p>0.05$; Picrotoxin \times CB65interaction, $F(1,54)=1.36$; $p>0.05$. Different letters (a and b) indicate significant differences between treatments ($p<0.001$). The average body weight for each group was as follows: control, 63.41 g; CB65, 63.75 g; Picrotoxin, 62.74 g; CB65 + Picrotoxin, 62.51 g.

In experiment 4, ICV injection of CB65 ($1.25\mu\text{g}$) significantly amplified feeding behavior in neonatal layer-type chicks in comparison with control group [$F(1,26)=47.15$; $p<0.001$]. ICV injection of sub effective dose of GABA_B antagonist, CGP54626 (21ng), had no role on food consumption compared to control group [$F(1,26)=0.49$; $p>0.05$]. Also, their combination (CGP54626+CB65) had no effect on CB₂ receptors agonist-induced hyperphagia [Time, $F(2,55)=124.13$, $p<0.001$; CGP54626 \times CB65interaction $F(1,26)=1.27$; $p>0.05$] [Fig 4].




Figure 4 – Effect of ICV injection of CB65 ($1.25\mu\text{g}$), CGP54626 (21ng) and their combination on percent of body weight cumulative food intake in neonatal layer type chicks ($n=48$). CB65: selective CB₂ receptors agonist, CGP54626: GABA_B antagonist. Data are expressed as mean \pm SEM. F and P value for within and between subject factors are as follows: Time, $F(2,55)=124.13$, $p<0.001$; CB65, $F(1,26)=47.15$, $p<0.001$; CGP54626, $F(1,26)=0.49$, $p>0.05$; CGP54626 \times CB65interaction, $F(1,26)=1.27$; $p>0.05$. Different letters (a and b) indicate significant differences between treatments ($p<0.001$). The average body weight for each group was as follows: control, 61.43 g; CB65, 62.04 g; CGP54626, 63.27 g; CB65 + CGP54626, 62.17 g.

In experiment 5, ICV injection of effective dose of GABA_A agonist, (Gaboxadol, $0.2\mu\text{g}$) significantly increased cumulative food intake compared to control group [$F(1,53)=87.04$; $p<0.001$]. ICV injection of sub effective dose of selective CB₁ receptors antagonist, (SR141716A, $6.25\mu\text{g}$) had no effect on food intake [$F(1,53)=0.31$; $p>0.05$]. Also, co-administration of SR141716A + Gaboxadol significantly decreased Gaboxadol-induced hyperphagia at 60 and 120 min after the injection [Time, $F(2,79)=56.27$, $p<0.001$; SR141716A \times Gaboxadol interaction, $F(1,53)=65.93$; $p<0.001$] [Fig 5].




Figure 5 – Effect of ICV injection of Gaboxadol ($0.2\mu\text{g}$), SR141716A ($6.25\mu\text{g}$) and their combination on percent of body weight cumulative food intake in neonatal layer type chicks ($n=48$). Gaboxadol: GABA_A agonist, SR141716A: selective CB₁ receptors antagonist. Data are expressed as mean \pm SEM. F and P value for within and between subject factors are as follows: Time, $F(2,79)=56.27$, $p<0.001$; Gaboxadol, $F(1,53)=87.04$, $p<0.001$; SR141716A, $F(1,53)=0.31$, $p>0.05$; SR141716A \times Gaboxadol interaction, $F(1,53)=65.93$; $p<0.001$. Different letters (a, b and c) indicate significant differences between treatments ($p<0.001$). The average body weight for each group was as follows: control, 64.26 g; Gaboxadol, 64.74 g; SR141716A, 64.06 g; Gaboxadol + SR141716A, 63.81 g.

In experiment 6, ICV injection of Gaboxadol at dose of $0.2\mu\text{g}$ amplified food consumption [$F(1,21)=38.12$; $p<0.001$] but ICV injection of sub effective dose of selective CB₂ receptors antagonist (AM630, $1.25\mu\text{g}$) had no effect on feeding behavior in neonatal layer-type chicks [$F(1,21)=1.02$; $p>0.05$]. Co-injection of AM630+ Gaboxadol had no effect on GABA_A receptors-induced hyperphagia [Time, $F(2,17)=48.91$, $p<0.001$; AM630 \times Gaboxadol interaction, $F(1,21)=2.37$; $p>0.05$] [Fig 6].




Figure 6 – Effect of ICV injection of Gaboxadol (0.2µg), AM630 (1.25µg) and their combination on percent of body weight cumulative food intake in neonatal layer type chicks (n=48). Gaboxadol: GABA_A agonist, AM630: selective CB₂ receptors antagonist. Data are expressed as mean ± SEM. F and P value for within and between subject factors are as follows: Time, $F(2,17)=48.91$, $p<0.001$; Gaboxadol, $F(1,21)=38.12$, $p<0.001$; AM630, $F(1,21)=1.02$, $p>0.05$; AM630 × Gaboxadol interaction, $F(1,21)=2.37$, $p>0.05$. Different letters (a and b) indicate significant differences between treatments ($p<0.001$). The average body weight for each group was as follows: control, 62.35 g; Gaboxadol, 61.96 g; AM630, 61.47 g; Gaboxadol + AM630, 62.94 g.

In experiment 7, ICV injection of effective dose of GABA_B agonist (baclofen, 0.2µg) significantly improved food intake in comparison to the control group [$F(1,31)=46.73$; $p<0.001$]. ICV injection of selective CB₁ receptors antagonist (SR141716A, 6.25µg) had no effect on food intake in neonatal layer-type chicks [$F(1,31)=0.09$; $p>0.05$]. Co-administration of SR141716A + baclofen had no effect on baclofen-induced hyperphagia [Time, $F(2,83)=102.17$, $p<0.001$; SR141716A × baclofen interaction, $F(1,31)=0.04$; $p>0.05$] [Fig 7].




Figure 7 – Effect of ICV injection of Baclofen (0.2µg), SR141716A (6.25µg) and their combination on percent of body weight cumulative food intake in neonatal layer type chicks (n=48). Baclofen: GABA_B agonist, SR141716A: selective CB₁ receptors antagonist. Data are expressed as mean ± SEM. F and P value for within and between subject factors are as follows: Time, $F(2,83)=102.17$, $p<0.001$; Baclofen, $F(1,31)=46.73$, $p<0.001$; SR141716A, $F(1,31)=0.09$, $p>0.05$; SR141716A × Baclofen interaction, $F(1,31)=0.04$; $p>0.05$. Different letters (a and b) indicate significant differences between

treatments ($p<0.001$). The average body weight for each group was as follows: control, 64.11 g; baclofen, 64.26 g; SR141716A, 63.21 g; Baclofen + SR141716A, 62.75 g.

In experiment 8, ICV injection of baclofen (0.2µg) showed hyperphagic effect [$F(1,46)=63.01$; $p<0.001$]; but ICV injection of AM630 (selective CB₂ receptors antagonist, 1.25µg) had no effect on feeding behavior [$F(1,46)=0.06$; $p>0.05$]. Also, the combined injection of baclofen + AM630 was not able to fluctuate GABA_B receptors-induced food intake in neonatal layer-type chicks [Time, $F(2,65)=70.31$, $p<0.001$; AM630 × Baclofen interaction, $F(1,46)=0.83$; $p>0.05$] [Fig 8].




Figure 8 – Effect of ICV injection of Baclofen (0.2µg), AM630 (1.25µg) and their combination on percent of body weight cumulative food intake in neonatal layer type chicks (n=48). Baclofen: GABA_B agonist, AM630 (selective CB₂ receptors antagonist). Data are expressed as mean ± SEM. F and P value for within and between subject factors are as follows: Time, $F(2,65)=70.31$, $p<0.001$; Baclofen, $F(1,46)=63.01$, $p<0.001$; AM630, $F(1,46)=0.06$, $p>0.05$; AM630 × Baclofen interaction, $F(1,46)=0.83$; $p>0.05$. Different letters (a and b) indicate significant differences between treatments ($p<0.001$). The average body weight for each group was as follows: control, 61.48 g; baclofen, 61.92 g; AM630, 63.19 g; baclofen + AM630, 62.38 g.

DISCUSSION

So far, numerous researches were done to investigate the role of neurotransmitters on appetite regulation in mammals, but aspects of food intake regulation in avian are still unclear (Zendehdel & Hassanpour 2014). To the best of our knowledge, there are no studies describing the interaction between CB₁ and GABA_B systems on food intake in FD₃ neonatal layer-type chicks. According to the results, ICV injection of CB₁ and CB₂ agonists amplified food consumption in layer-type chicks which was similar to mammals (Chen *et al.*, 2006; Wiley *et al.*, 2012) and recent studies on layer-type chicks (Alizadeh *et al.*, 2015; Hassanpour *et al.*, 2015). However, these results were dissimilar compared to broilers which only CB₂ receptors were responsible for feeding (Emadi *et al.*, 2011; Novoseletsky *et al.*, 2011). The ECBs is an important modulator in the central and peripheral regulation of energy and



metabolism (Bellocchio *et al.*, 2013). Broilers are genetically selected for rapid muscle and body weight gain while layer has indirectly been selected for slow growth and low body weight. So, presumably, genetic selection altered chicken's brain neurological pathways associated with food intake (Denbow 1994; Richards 2003). It is a hypothetical effect of ECBs on appetite regulation which mediates at two levels. First, ECBs reinforces the motivation to find and eating, possibly via interaction with the mesolimbic pathways involved in reward mechanisms. Second, it activated 'on demand' in the hypothalamus after short-term food deprivation, then transiently regulates the action of the other orexigenic and anorectic neurotransmitters. The dual action hypothesis in mesolimbic and hypothalamic regions revealed by the finding that injection of ECBs into these brain areas provoke food intake in rodents (Di Marzo & Matias, 2005).

Based on the results, ICV injection of GABA_A and GABA_B receptors agonists increased food intake in neonatal layer-type chicks. These results were in agreement with observation in rats (Di Marzo *et al.*, 2004), broilers (Jonaidi *et al.*, 2002), turkeys (Denbow, 1991) sheep (Seone *et al.*, 1984) and pigs (Baldwin *et al.*, 1990). Regarding the site of action of GABA, with rats and sheep suggest GABA might alter food intake on some intra hypothalamic or extra-hypothalamic structures (Jonaidi *et al.*, 2002). Several possible mechanisms introduced the mechanism of action of GABA. For example, it may inhibit the satiety centers e.g. ventromedial hypothalamus (VMH). GABA may act via dis-inhibition of the brain feeding centers by inhibitory GABAergic projection from the medial accumbens to the lateral hypothalamus (LH). Finally, GABA can exert orexigenic effect via a direct excitatory effect in some regions of the brain including the hypothalamus (Jonaidi *et al.*, 2002).

As observed in this study, hyperphagic effects of 2-AG (CB₁ receptors agonist) attenuated by co-injection of 2-AG + Picrotoxin (GABA_A receptors antagonist). CB₁ receptors control the activity of many neurotransmitter systems involved in central regulation of food intake (Pagotto *et al.*, 2006). In the CNS, CB₁ is predominantly expressed presynaptically, modulating the release of neurotransmitters, such as GABA, dopamine, noradrenaline, glutamate and serotonin (Cota *et al.*, 2003). These data showed there might be an interaction between central CB₁ and GABAergic systems via CB₁ and GABA_A receptors. The broad distribution of CB₁ receptors in the CNS, along with its modulatory role on other neurotransmitter

systems, complicates the mechanisms underlying its effect on food intake regulation (Rey *et al.*, 2012).

Retrograde signaling by ECBs released from depolarized postsynaptic neurons inhibits presynaptic GABA release in the lateral hypothalamus and arcuate nucleus (ARC) (Di Marzo *et al.*, 2004). The ARC plays a crucial role in the integration of signals regulating appetite. The ARC is accessible to circulating signals of energy balance via the median eminence, this is not protected by the blood-brain barrier. In the ARC two types of neurons identified which integrates signals of nutritional status and influence energy homeostasis: pro-opiomelanocortin (POMC) and cocaine and amphetamine regulated transcript (CART) which are anorexigenic neuropeptides while another neuronal circuit stimulates feed intake via neuropeptide Y (NPY) and agoutirelated protein (AgRP) (Zendeheel & Hassanpour 2014). The AgRP, POMC and NPY contain the GABA_A R alpha (3) subunit in the ventromedial and ventrolateral part of ARC (Jonaidi *et al.*, 2002). ECBs impress their hyperphagic effect by modulating POMC neurons via CB₁ receptors in the ARC (Hentges *et al.*, 2005). Also, ICV injection of the CB₁ receptors agonist increases NPY levels (D'Addario *et al.*, 2014). A strong indication exists for CB₁ receptors on modulating appetite by enhancement of NPY (Cota *et al.*, 2003). One potential target for cannabinoids in the arcuate nucleus is NPY/ AgRP /GABA neurons (Menzies *et al.*, 2010). This is most likely a direct inhibitory effect on spontaneous release mediated via pre-synaptic CB₁ receptors on NPY/ AgRP /GABA neurons or other GABAergic terminals in the explanation. It is reported that CB₁ receptor mRNA are not expressed by arcuate NPY/ AgRP /GABA neurons and target probably underlies the direct effect (Menzies *et al.*, 2010). CB₁ receptors are expressed in presynaptic terminals of excitatory neurons (Sharkey *et al.*, 2014). Functionally, GABA_A receptors seem to be located both pre- and post-synapses (Tellez *et al.*, 2012). GABA_A receptors are ionotropic receptors which couple to a Cl⁻ ionophore (Jonaidi *et al.*, 2012). ECBs synthesized in postsynaptic neurons and binds to CB receptors on the presynaptic membrane. These receptors belong to G protein, inhibits Ca²⁺ influx which decreases the release of other neurotransmitters (Williams and Kirkham 2002). CBs modulate GABA release in a variety of CNS regions, including the hippocampus, basal ganglia, cerebellum, and brainstem. Activation of CB₁ receptors modulates the release of GABA in the brain. However, minute information exists about the mechanism behind the interaction of CB₁ with GABA_A receptors.



Anatomical investigations have shown high levels of CB₁ receptor immune reactivity and mRNA associated with GABAergic neurons. CBs modulate GABA release by a presynaptic mechanism. Presumably, effect of CB₁ receptors on GABA_A release is mediated via inhibition of presynaptic Ca²⁺ channels (Irving *et al.*, 2002).

To the best of our knowledge, CB₁ receptors mRNA expressed by POMC neurons and CBs directly inhibit arcuate POMC neurons. In turn, POMC neurons inhibit GABA release from NPY/AgRP neurons (Menzies *et al.*, 2010). As observed in this study, the hyperphagic effect of CB₁ receptors diminished by GABA_A receptors antagonist (experiment 1). For instance, hyperphagic effects of GABA_A receptors minimized by CB₁ receptors antagonist (experiment 5). Presumably, ECBs, by inhibition of the hypothalamic POMC, leads to increasing GABA release in the CNS of neonatal layer-type chicken. So, it seems both CBergic and GABAergic systems impress their hyperphagic effects via CB₁ receptors in neonatal layer-type chicken. Also, perhaps CBergic and GABAergic systems interact with other systems such as opioidergic system (Menzies *et al.*, 2010).

Most research on central food intake regulation was done on rat models because few investigations were in avian. These results can use as base information on central appetite regulation in layer-type chicken. Finally, the authors recommend merit investigation needed to find direct cellular and molecular signaling pathways of CBergic and GABAergic mechanism with other neurotransmitters in the physiology of eating behavior in poultry.

CONFLICT OF INTEREST

Authors declare that they have no conflict of interest.

INFORMED CONSENT

This manuscript does not contain any studies with human subjects performed by any of the authors.

HUMAN AND ANIMAL RIGHTS

All experiments executed according to the Guide for the Care and Use of Laboratory Animals and approved by the institutional animal ethics committee.

REFERENCES

Alimohammadi S, Zendeheel M, Babapour V. Modulation of opioid-induced feeding behavior by endogenous nitric oxide in neonatal layer-type chicks. *Veterinary Research Communications* 2015;39:105-113.

Alizadeh A, Zendeheel M, Babapour V, Charkhkar S, Hassanpour S. Role of cannabinoidergic system on food intake in neonatal layer-type chicken. *Veterinary Research Communications* 2015;39:151-157.

Baldwin BA, Ebenezer IS, De La Riva C. Effects of intracerebroventricular injection of muscimol or GABA on operant feeding in pigs. *Physiology & Behavior* 1990;48:417-421.

Bellocchio L, Soria-Gómez E, Quarta C, Metna-Laurent M, Cardinal P, Binder E, et al. Activation of the sympathetic nervous system mediates hypophagic and anxiety-like effects of CB₁ receptor blockade. *PANS* 2012;110(12):4786-4791.

Blevins JE, Stanley BG, Reidelberger RD. DMSO as a vehicle for central injections: tests with feeding elicited by norepinephrine injected into the paraventricular nucleus. *Pharmacology, Biochemistry, and Behavior* 2002;71:277-282.

Bonaventura MM, Crivello M, Ferreira ML, Repetto M, Cymeryng C, Libertun C, et al. Effects of GABAB receptor agonists and antagonists on glycemia regulation in mice. *European Journal of Pharmacology* 2012;677:188-196.

Chen RZ, Frassetto A, Fong TM. Effects of the CB₁ cannabinoid receptor inverse agonist AM251 on food intake and body weight in mice lacking I-opioid receptors. *Brain Research* 2006;1108:176-178.

Cota D, Marsicano G, Lutz B, Vicennati V, Stalla GK, Pasquali R, et al. Endogenous cannabinoid system as a modulator of food intake. *International Journal of Obesity* 2003;27:289-301.

D'Addario C, Micioni Di Bonaventura MV, Puccia M, Romano A, Gaetani S, Ciccocioppo R, et al. Endocannabinoid signaling and food addiction. *Neuroscience & Biobehavioral Reviews* 2014;47:203-224.

Davis JL, Masuoka DT, Gerbrandt LK, Cherkin A. Autoradiographic distribution of L-proline in chicks after intracerebral injection. *Physiology & Behavior* 1979;22:693-695.

Denbow DM. Induction of food intake by a GABAergic mechanism in the turkey. *Physiology & Behavior* 1991;49:485-488.

Denbow DM. Peripheral regulation of food intake in poultry. *Journal of Nutrition* 1994;124:1349S-1354S.

Di Marzo V, Bifulco F, De Petrocellis L. The endocannabinoid system and its therapeutic exploitation. *Nature Reviews Drug Discovery* 2004;3:771-784.

Di Marzo V, Matias I. Endocannabinoid control of food intake and energy balance. *Nature Neuroscience* 2005;8(5):585-589.

Ebenezer IS. The effect of intracerebroventricular administration of baclofen on food intake in rats. *Neuro Report* 1990;1:73-76.

Emadi L, Jonaidi H, Hosseini Amir Abad E. The role of central CB₂ cannabinoid receptors on food intake in neonatal chicks. *Journal of Comparative Physiology A* 2011;197:1143-1147.

Furuse M, Matsumoto M, Saito N, Sugahara K, Hasegawa S. The central corticotropin-releasing factor and glucagon-like peptide-1 in food intake of the neonatal chick. *European Journal of Pharmacology* 1997;339:211-214.

Hassanpour S, Zendeheel M, Babapour V, Charkhkar S. Endocannabinoid and nitric oxide interaction mediates food intake in neonatal chicken. *British Poultry Science* 2015;56(4):443-451.

Hentges ST, Low MJ, Williams JT. Differential regulation of synaptic inputs by constitutively released endocannabinoids and exogenous cannabinoids. *Journal of Neuroscience* 2005;25:9746-9751.



- Honda K. Glucagon-related peptides and the regulation of food intake in chickens. *Animal Science Journal* 2016. DOI:10.1111/asj.12619.
- Irving AJ, Rae MG, Coutts AA. Cannabinoids on the Brain. *Scientific World Journal* 2002;2:632-648.
- Jonaidi H, Abbassi L, Yaghoobi MM, Kaiya H, Denbow DM, Kamali Y, et al. The role of GABAergic system on the inhibitory effect of ghrelin on food intake in neonatal chicks *Neuroscience Letters* 2012;520:82-86.
- Jonaidi H, Babapour V, Denbow DM. GABAergic control of food intake in the meat-type chickens. *Physiology & Behavior* 2002;76:465- 468
- Kangas BD, Delatte MS, Vemuri VK, Thakur GA, Nikas SP, Subramanian KV, et al. Cannabinoid discrimination and antagonism by CB1 neutral and inverse agonist antagonists. *Journal of Pharmacology and Experimental Therapeutics* 2013;344:56-567.
- Menzies JRW, Ludwig M, Leng G. Direct and indirect effects of cannabinoids on in vitro GABA release in the rat arcuate nucleus . *Journal of Neuroendocrinology* 2010;22:585-592.
- Novoseletsky N, Nussinovitch A, Friedman-Einat M. Attenuation of food intake in chicks by an inverse agonist of cannabinoid receptor 1 administered by either injection or ingestion in hydrocolloid carriers. *General and Comparative Endocrinology* 2011;170:522-527.
- Olanrewaju HA, Thaxton JP, Dozier WA, Purswell J, Roush WB, Branton SL. A review of lighting programs for broiler production. *International Journal of Poultry Science* 2006;5(4):301-308.
- Onaivi ES, Carpio O, Ishiguro H, Schanz N, Uhl GR, Benno R. Behavioral effects of CB₂ cannabinoid receptor activation and its influence on food and alcohol consumption. *Annals of the New York Academy of Sciences* 2008;1139:426-433.
- Pagotto U, Marsicano G, Cota D, Lutz B, Pasquali R. The emerging role of the endocannabinoid system in endocrine regulation and energy balance. *Endocrine Reviews* 2006;27(1):73-100.
- Pu S, Jain MR, Horvath TL, Diano S, Kalra PS, Kalra SP. Interactions between neuropeptide Y and gamma-aminobutyric acid in stimulation of feeding:a morphological and pharmacological analysis. *Endocrinology* 1999;140:933-940.
- Qi W, Ding D, Salvi RJ. Cytotoxic effects of dimethyl sulphoxide (DMSO) on cochlear organotypic cultures. *Hearing research* 2008;236:52-60.
- Rey AA, Purrio M, Viveros MP, Lutz B. Biphasic Effects of Cannabinoids in Anxiety Responses: CB₁ and GABA_B Receptors in the Balance of GABAergic and Glutamatergic Neurotransmission. *Neuropsychopharmacology* 2012;37:2624-2634.
- Richards MP. Genetic regulation of feed intake and energy balance in poultry. *Poultry Science* 2003;82:907-916.
- Saito ES, Kaiya H, Tachibana T, Tomonaga S, Denbow DM, Kangawa K, et al. Inhibitory effect of ghrelin on food intake is mediated by the corticotropin-releasing factor system in neonatal chicks. *Regulatory Peptides* 2005;125:201-208.
- Seone JR, Dumont F, Girard CL, Bedard L, Matte JJ. Effects of intraventricular injections of gamma-aminobutyric acid and related substances on feeding behavior in satiated sheep. *Canadian Journal of Physiology and Pharmacology* 1984;62:1296-1209.
- Sharkey KA, Darmani NA, Parker LA. Regulation of nausea and vomiting by cannabinoids and the endocannabinoid system. *European Journal of Pharmacology* 2014;722:134-146.
- Stratford TR, Wirtshafter D. Injections of muscimol into the paraventricular thalamic nucleus, but not mediodorsal thalamic nuclei, induce feeding in rats. *Brain Research* 2013;1490:128-133
- Tajalli S, Jonaidi H, Abbasnejad M, Denbow DM. Interaction between nociceptin/orphanin FQ (N/OFQ) and GABA in response to feeding. *Physiology & Behavior* 2006;89:410-413.
- Tellez R, Gómez-Viquez L, Meneses A. GABA, glutamate, dopamine and serotonin transporters expression on memory formation and amnesia. *Neurobiology of Learning and Memory* 2012;97:189-201.
- Van Tienhoven A, Juhasz LP. The chicken telencephalon, diencephalon and mesencephalon in stereotaxic coordinates. *Journal of Comparative Neurology* 1962;118:185-197
- Volkow ND, Wang GJ, Baler RD. Reward, dopamine and the control of food intake:Implications for obesity. *Trends in Cognitive Sciences* 2011;15(1):37-46.
- Wiley JL, Marusich JA, Zhang Y, Fulp A, Maitra R, Thomas BF, et al. Structural analogs of pyrazole and sulfonamide cannabinoids:effects on acute food intake in mice. *European Journal of Pharmacology* 2012;695:62-70.
- Williams CM, Kirkham TC. Reversal of Δ9-THC hyperphagia by SR141716 and naloxone but not dexfenfluramine. *Pharmacology Biochemistry and Behavior* 2002;71:333-340.
- Zendeheel M, Babapour V, Jonaidi H. Effects of central histamine receptors blockade on GABA (A) agonist-induced food intake in broiler cockerels. *Pakistan Journal of Biological Sciences* 2008;11(3):416-421.
- Zendeheel M, Baghbanzadeh A, Babapour V, Cheraghi J. The effects of bicuculline and muscimol on glutamate-induced feeding behaviour in broiler cockerels. *Journal of Comparative Physiology A* 2009;195:715-720.
- Zendeheel M, Hassanpour S. Ghrelin-induced hypophagia is mediated by the β2 adrenergic receptor in chicken. *The Journal of Physiological Sciences* 2014;64:383-391.