



Revista Facultad Nacional de Agronomía
- Medellín

ISSN: 0304-2847

rfnagron_med@unal.edu.co

Universidad Nacional de Colombia
Colombia

Numpaque, Manuel Alejandro; Gil González, Jesús Humberto; Durango Restrepo, Diego
Luis

Biotransformation of ferulic acid by the phytopathogenic fungi *Colletotrichum acutatum*
and *Lasiodiplodia theobromae*

Revista Facultad Nacional de Agronomía - Medellín, vol. 69, núm. 1, 2016, pp. 7835-7844

Universidad Nacional de Colombia
Medellín, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=179943371009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Biotransformation of ferulic acid by the phytopathogenic fungi *Colletotrichum acutatum* and *Lasiodiplodia theobromae*

Biotransformación del ácido ferúlico con los hongos fitopatógenos
Colletotrichum acutatum y *Lasiodiplodia theobromae*

doi: 10.15446/rfna.v69n1.54751

Manuel Alejandro Numpaque¹, Jesús Humberto Gil González^{1*} and Diego Luis Durango Restrepo¹

ABSTRACT

Key words:

Microbial
transformation
Flavoring
4-vinylguaiacol
Metabolic pathway

The microbial transformation of ferulic acid (FA) offers a cleaner, more economical alternative for the natural production of flavorings and fragrances. In the present study, the biotransformation of FA using the filamentous phytopathogenic fungi *Colletotrichum acutatum* and *Lasiodiplodia theobromae* was researched. Initially, the toxicity of FA against both fungi was evaluated; the FA displayed a moderate toxicity (total inhibition at concentrations $\geq 2000 \text{ mg L}^{-1}$) and apparently a detoxification mechanism was present. Afterwards, the microorganisms were incubated with the substrate at room conditions using a Czapek-Dox culture medium. The results demonstrated that the FA was mainly converted to 4-vinylguaiacol, reaching the highest abundance within the first 48 hours. To a lesser extent, acetovanillone, ethylguaiacol, and vanillin, among others, were produced. Interestingly, the compounds generated in the biotransformation of FA with *C. acutatum* and *L. theobromae* have been used as flavorings. Based on the identified metabolites, a possible metabolic pathway was proposed.

RESUMEN

Palabras claves:

Transformación
microbiana,
Saborizantes
4-vinilguayacol
Ruta metabólica

La transformación microbiana de ácido ferúlico (AF) puede ofrecer una alternativa más limpia y económica para la producción natural de algunos saborizantes y aromas. En el presente estudio, se investigó la biotransformación de AF usando los hongos filamentosos fitopatógenos *Colletotrichum acutatum* y *Lasiodiplodia theobromae*. Inicialmente, se evaluó la toxicidad del AF contra ambos hongos; el sustrato exhibió una toxicidad moderada (inhibición total a concentraciones $\geq 2000 \text{ mg L}^{-1}$) y aparentemente se presentó un mecanismo de desintoxicación. Luego, los microorganismos se incubaron con el sustrato a condiciones ambientales, usando el medio de cultivo Czapek-Dox. Los resultados muestran que el AF es convertido principalmente en 4-vinilguayacol, alcanzando su mayor abundancia dentro de las primeras 48 horas. En menor proporción se producen acetovainillona, etilguayacol, vainillina, entre otros. Interesantemente, los compuestos generados en la biotransformación de AF con *C. acutatum* y *L. theobromae* se han empleado como agentes saborizantes. Con base en los metabolitos identificados, se propone una posible ruta metabólica.

^{1,2,3} Facultad de Ciencias Agrarias - Universidad Nacional de Colombia - A.A. 1779, Medellín, Colombia.

* Corresponding author <jhgilg@unal.edu.co>

Flavorings and fragrances are widely used in the food, drink, and cosmetic industries; however, the majority of flavorings in the global market are obtained through chemical synthesis and less than 5% have a natural origin, generally obtained through plant extraction (Bicas *et al.*, 2010). Currently, chemical synthesis dominates the production of flavorings, using methods that often become incompatible with the environment; furthermore, the low selectivity of the reactions may cause the formation of undesirable product mixtures, which reduces the efficiency of the process and increases the costs of production. According to the European Union Legislature (Xu, *et al.*, 2007), the use of synthetic flavorings is becoming increasingly restricted in food, drinks, and cosmetics; therefore, natural flavorings are appealing to the global market despite their high price. Nevertheless, the production of natural flavorings from direct extractions of botanical materials suffers several problems. Firstly, the concentration of compounds with flavoring or fragrant characteristics can be low in plants, making extraction a costly stage. In addition, agricultural production takes time and the quality of the harvest is influenced by climatic conditions, the availability of soils and plant diseases.

According to the FDA and the European Union Legislature, the products obtained by biotechnological methods can be considered natural when the substrate that is used is natural (Serra *et al.*, 2005). Therefore, biotransformation processes have been suggested as a new environmentally friendly alternative for the production of natural fragrances that are in demand. Ferulic acid (FA), a hydroxycinnamic acid found in the cellular wall of plants, is a natural precursor that plays an important role in the formation of fragrances such as vanillin and some vinylphenols (Hu *et al.*, 2015). The industrial demand for 4-vinylphenols such as 4-vinylcatechol, 4-vinylguaiacol and 2,6-dimethoxy-4-vinylphenol is not met by the availability provided by natural sources (Bernini *et al.*, 2007). This makes FA a highly interesting compound as a substrate for the production of natural flavorings with a commercial value through biotransformation processes. In the literature, there is a significant number of reports on the microorganisms and biotechnical process for the production of flavorings from FA. Some bacterium and yeast genera, such as *Pseudomonas*, *Bacillus*, *Streptomyces*, *Lactobacillus*, *Enterobacter*,

Streptomyces, *Cupriavidus*, *Candida*, *Debaryomyces*, and *Saccharomyces*, among others, have developed multiple routes for the bioconversion of FA into vanillin, vanillic acid, protocatechuic acid, and vanillic alcohol (Mishra *et al.*, 2014; Gallage and Møller, 2015). Nevertheless, the biotransformation of FA by fungi has been little-studied; in particular, the fungi *Schizophyllum commune*, *Aspergillus niger*, *Rhizopus oryzae*, *Sporotrichum thermophile*, and *Pycnoporus cinnabarinus* have been evaluated for the biotransformation of FA (Baquero-Peña *et al.*, 2010; Tsujiyama and Ueno, 2008; Shanker *et al.*, 2007; Topakas *et al.*, 2003; Bonnin *et al.*, 1999). For their part, the phytopathogenic fungi *Lasiodiplodia theobromae* and *Colletotrichum acutatum* have demonstrated high metabolic versatility because they have transformed different phenylpropanoid substrates, resulting in compounds with an aggregate value (Velasco *et al.*, 2007; Velasco *et al.*, 2009; Velasco *et al.*, 2010; Velasco *et al.*, 2012). However, the capacity of both fungi to produce flavoring compounds through biotransformation has been scarcely studied. This article reports for the first time the ability of the fungi *C. acutatum* and *L. theobromae* to transform FA into compounds with an aggregate value with flavoring attributes, such as 4-vinylguaiacol, 4-ethylguaiacol, vanillin, vanillic acid and acetovanillone.

MATERIALS AND METHODS

Materials

Analytical grade solvents: ethyl acetate (EtOAc), methanol (MeOH), *n*-hexane and CHCl₃ were obtained from Merck. The FA standard was acquired from Sigma-Aldrich. In addition, silica gel 60 (0.040-0.063 mm) and Sephadex® LH-20 from Merck and Sigma-Aldrich, respectively, were used. For the culture media, this study employed casein peptone (Merck), alpha-D(+)-glucose anhydrous (Acros Organics), yeast extracts (Oxoid), K₂HPO₄ (Ma-linckrodt Chemical), NaNO₃ (Merck); MgSO₄·7H₂O (Protokimica), and FeSO₄·7H₂O (Carlos Erba). The extracts used for the liquid-liquid extractions were bidistilled before their use.

Separation and analytical methods

Thin layer chromatography (TLC) was carried out using chromatoplates Merck Kieselgel 60 F₂₅₄ 0.25 mm thick and as the mobile phase a *n*-hexane:EtOAc (8:2) mixture. The compounds were visualized under UV radiation at 254 and 365 nm, and spraying with the acetic acid:H₂SO₄:H₂O

mixture (143:28:30) followed by heating (~100 °C, 1 min). Column chromatography (CC) was performed using silica gel 60 or Sephadex® LH-20 as stationary phase. The gas chromatography (GC) analysis used a Hewlett-Packard 6890 chromatograph (Agilent Technologies) interfaced to Agilent HP 5973 Mass Selective Detector in the electronic ionization mode. A DB-35MS column (30 m x 0.25 mm i.d. x 0.25 µm coating thickness) was employed. The chromatographic conditions included the following: column temperature, 50-250 °C at 10 °C min⁻¹; injector temperature, 230 °C; detector temperature, 280 °C; and carrier gas, N₂ at 1 mL min⁻¹. The relative composition of the individual components was determined from the peaks average area. Identification of some metabolites was based on interpretation of their mass spectra, comparison with authentic compounds, and by contrast with the NIST 02 Mass Spectral Library. The nuclear magnetic resonance (NMR) spectra were obtained using deuterated chloroform as a solvent with a Bruker AMX 300 model. The multiplicities were established with the JMOD pulse sequence. The chemical displacements (δ) were expressed in values of ppm and the coupling constant (J) in hertz (Hz).

Toxicity of the ferulic acid (FA) substrate

The toxicity evaluations, referred to as FA's antifungal activity against the filamentous fungi *C. acutatum* and *L. theobromae* (isolated and morphologically characterized by the Laboratorio de Sanidad Vegetal of the Universidad Nacional de Colombia, Medellín), were carried out with the methodology described by Bustillo *et al.* (2003), with some modifications. In Potato Dextrose Agar (PDA) medium at 50 °C, was added enough (FA) till achieve the desired concentrations (100, 500, 1000, 2000, 4000 mg L⁻¹), and then the mixture was poured immediately into 9-cm-diameter Petri dishes. Later, a mycelial plug (6 mm in diameter) cut from the growing edge of 2-day-old culture of each fungus was transferred to the center of Petri dish plates. The cultures were incubated at room temperature and the mycelial growth diameters of the fungi were measured every 12 h for 96 h for *L. theobromae* and every 24 h for 168 h for *C. acutatum*. The measurements were taken in triplicate and each trial had the respective controls (absolute control: PDA medium devoid of FA, and solvent control: ethanol). The toxicity of the FA was expressed as the percentage of mycelial growth inhibition, which was calculated with equation (1):

$$\text{Inhibition(\%)} = \left(1 - \frac{\text{Mean growth of the treatment(mm)}}{\text{Mean growth of the control(mm)}} \right) \times 100 \quad (1)$$

Curves were constructed for the FA toxicity against the microorganisms, which was used to determine an approximate concentration that would inhibit the mycelial growth of the fungi between 80 and 90% at the midpoint of the evaluation period; this value was used as the initial concentration of the substrate in the subsequent biotransformation processes.

Preparation of the pre-inoculums of *C. acutatum* and *L. theobromae*

A Czapek-Dox medium was used (Solution A: glucose 5%, yeast extract 0.1%; Solution B: K₂HPO₄ 0.5%, NaNO₃ 0.2%, MgSO₄·7H₂O 0.05%, FeSO₄·7H₂O 0.001%) in the biotransformations with an inoculum of each fungus of around 1 month of age, previously cultivated on PDA. For each liter of liquid culture medium, a Petri dish was used with the microorganism. The pathogen was inoculated in four 1.0 L Erlenmeyer flasks that contained 500 mL of the established medium. The flasks were agitated (shaker, 120 rpm; Centricol, series 0239, with an incubation chamber) under environmental conditions for 96 h. The mycelia were recovered with filtration, washed with distilled water, and used in the biotransformation at the preparative scale and in the time course experiments.

Biotransformation in the preparative scale and isolation of the metabolites

The mycelia of *C. acutatum* and *L. theobromae*, pre-incubated in 2.0 L of the medium, were transferred under sterile conditions in four, 1.0 L Erlenmeyer flasks that contained 500 mL of the corresponding culture medium and the substrate (FA at 900 mg L⁻¹ for *L. theobromae* and 1400 mg L⁻¹ for *C. acutatum*). The process was carried out at room temperature and under agitation (120 rpm) for 312 and 360 h for *L. theobromae* and *C. acutatum*, respectively. After the incubation period, the culture media and the mycelia were separated through filtration. The filtrate was used to isolate the major metabolic products. For the control, a biotransformation was carried out without the substrate. The resulting filtrate was saturated with NaCl and was extracted with EtOAc (3 x 400 mL). The organic phase was dried with anhydrous sodium sulfate, filtered and evaporated at a reduced pressure in a rotary evaporator. The resulting residue was fractionated

with successive CC, using as stationary phase silica gel (mobile phase: systems of increasing polarity of *n*-hexane-EtOAc) and Sephadex® LH-20 (elution system: mixture of *n*-hexane-CH₂Cl₂-MeOH, 2:1:1 w/v). Three major metabolic products was isolated and denominated as (II), (V) and (VII). A GC-MS analysis of the resulting fractions of the samples also revealed the presence of the compounds designated as (I), (II), (III), (IV), (V) and (VI). The metabolites were identified with spectroscopic and spectrometric methods and through a comparison of spectrums with those obtained for standard samples and/or with the data reported in the database of the NIST Mass Spectral Library, NIST'02 (version 2.0).

Time-course experiments

The material resulting from the pre-inoculation was filtered and the biomass was distributed in 12, 150.0 mL Erlenmeyer flasks with 75 mL of the Czapek-Dox culture medium and the substrate (FA) at a concentration of 900 and 1400 mg L⁻¹ for *L. theobromae* and *C. acutatum*, respectively. The conditions of temperature, agitation and time corresponded to those cited for the preparative biotransformation. Every 24 hours, the content of one Erlenmeyer flask (culture

medium containing the mycelia) was extracted with EtOAc (3 x 50 mL), following the procedure described for the isolation of metabolites. The resulting residue was re-dissolved with 5.0 mL of CHCl₃, filtered with a Whatman microfilter (0.45 µm) and analyzed by TLC and GC-MS. The relative abundance of the products was determined based on the area of the peaks in CG.

RESULTS AND DISCUSSION

FA toxicity for *C. acutatum* and *L. theobromae*

In order to determine the substrate concentration that was suitable for carrying out the biotransformation processes, the antifungal activity of the FA for both phytopathogenic fungi was evaluated. As can be seen in Figure 1, FA reduced the mycelial growth in a dose-dependent manner. *C. acutatum* was almost completely inhibited at the FA concentrations of 2000 and 4000 mg L⁻¹. In the meanwhile, for the interval of 24 to 169 h, at the 100, 500, and 1000 mg L⁻¹ levels, the inhibition of the growth varied throughout a range of 100-21%, 100-45% and 95-75%, respectively. At these concentrations, mycelial growth inhibition was decreased gradually with increasing incubation time, which could be attributed to a detoxification




Figure 1. *In vitro* antifungal activity of the FA against the fungus *C. acutatum*.

mechanism. In general, the results demonstrated a moderate radial growth inhibition of *C. acutatum* by the FA, in comparison with the values reported for other microorganisms. In particular, Sarma and Singh (2003) found that *Sclerotium rolfsii* was completely inhibited at a FA concentration of 1000 mg L⁻¹, while *Aspergillus niger* (Baquero-Peña *et al.*, 2010) and *Rhizopus oryzae* (Shanker *et al.*, 2007) were inhibited at 800 and 500 mg

L⁻¹, respectively. Based on the obtained results, a FA concentration of 1400 mg L⁻¹ was selected to carry out the biotransformation process; this value is intermediate between 2000 (fungal growth limited with inhibition over 95%) and 1000 mg L⁻¹ (inhibition of almost 85% at the midpoint of the evaluation). At this level, enzymatic activity that results in a suitable conversion process for FA by the microorganisms is expected.

Figure 2 shows that the mycelial growth of *L. theobromae* presented 100% inhibition at FA concentrations of 2000 and 4000 mg L⁻¹ during the evaluation time (96 h). In the time interval of 12 to 96 h at the 100, 500 and 1000 mg L⁻¹ levels, the inhibition of the growth varied in ranges of 47-12%, 88-58% and 100-80%, respectively. At these concentrations, there was a decrease in the mycelial inhibition with increase

in time of incubation, which could be attributed to the decrease in the FA toxicity by the microorganism, similar to the findings with *C. acutatum*. The FA concentration selected for the biotransformation with *L. theobromae* was 900 mg L⁻¹; a slightly higher value than the concentration that inhibited the fungus at 80% at the midpoint of the evaluation period. The moderate activity of FA agrees with different studies




Figure 2. *In vitro* antifungal activity of the FA against the fungus *L. theobromae*.

that have demonstrated that this organic acid has little fungitoxicity (Baquero-Peña *et al.*, 2010; Shanker *et al.*, 2007), which is why it has been suggested for possible use as material for biotransformations.

Biotransformation products of FA: isolation and identification

In order to isolate the metabolic products of FA, a preparative biotransformation was carried out in the Czapek-Dox medium using each of the phytopathogenic

fungi. The microorganism was incubated with FA for 336 h; afterwards, the extraction of the metabolites was done with ethyl acetate and the resulting extract was fractionated with size-exclusion chromatography on Sephadex LH-20 (100 x 2 cm) with a mixture of *n*-hexane-CH₂Cl₂-MeOH (50:25:25) as eluent. In total, three metabolic compounds were isolated: 4-vinylguaiaicol (**II**); acetovanillone (**V**) and dihydroferulic acid (**VII**), which were elucidated using ¹H-NMR and MS-EI techniques (Table 1).

Table 1. Products isolated from the biotransformation of the FA.

Metabolite	¹ H NMR (CDCl ₃ , 300 MHz) and MS-EI	Microorganism
4-vinylguaiaicol (II)	¹ H NMR (CDCl ₃ , 300 MHz): δ 3.88 (s, 3H, -OCH ₃), 5.15 (d, 1H, J = 10.8, =CH ₂), 5.61 (d, 1H, J = 17.5, =CH ₂), 5.78 (s, 1H, -OH); 6.67 (dd, 1H, J = 17.5, 10.8, =CH), 6.88-6.96 (m, 3H, H-Ar). MS-EI, m/z [int. rel.]: 150 [100] (M ⁺), 135 [80] (M ⁺ -CH ₃), 107 [42] (M ⁺ -C ₂ H ₄ -CH ₃), 77 [35] (C ₆ H ₅ ⁺). Retention time: Rt = 13.71 min.	<i>C. acutatum</i> <i>L. theobromae</i>
Acetovanillone (V)	¹ H NMR (CDCl ₃ , 300 MHz): δ 2.47 (s, 3H, -CH ₃), 3.90 (s, 3H, -OCH ₃), 6.88 (d, 1H, J = 8.7, H-Ar), 7.45-7.47 (m, 2H, H-Ar). MS-EI, m/z [int. rel.]: 166 [54] (M ⁺), 151 [100] (M ⁺ -CH ₃), 123 [25] (M ⁺ -CO-CH ₃), 108 [6], 93 [3]. Rt = 17.93 min.	<i>C. acutatum</i> <i>L. theobromae</i>
Dihydroferulic acid (VII)	¹ H NMR (CDCl ₃ , 300 MHz): δ 2.60 (t, 2H, J = 7.7, -CH ₂), 2.84 (t, 2H, J = 7.5, -CH ₂), 3.90 (s, 3H, -OCH ₃), 5.10 (s, 1H, -OH), 6.66 (dd, 1H, J = 8.0, 2.0, H-Ar), 6.69 (d, 1H, J = 2.0, H-Ar), 6.81 (d, 1H, J = 8.0, H-Ar).	<i>C. acutatum</i>

The spectroscopic data for the compounds (II), (V), and (VII) agree with the values reported in the literature. Compound (II) presented a molecular formula $C_9H_{10}O_2$ consistent with the molecular ion, $M^+ = 150$ amu. The 1H -NMR spectrum of compound (II) showed the singlet characteristic of the methoxy group (δ 3.88 ppm), a multiplet that corresponded to three aromatic protons (δ 6.88-6.96 ppm), and the two olefinic protons (δ 5.15, 5.61 and 6.67 ppm) with their respective multiplicities, which agrees with the reported by Karmakar *et al.* (2000). In (V), 1H -NMR spectrum showed the presence of two singlets (δ 2.47 and 3.90 ppm) assigned respectively to the methyl and methoxy groups, and three aromatic protons (δ 6.88 and 7.45-7.47 ppm); these characteristics, together with the molecular ion ($M^+ = 166$ amu) and fragmentation pattern in the mass spectrum agree with the reported by Luo *et al.*, 2008. For the compound (VII), the coupling pattern described an ABX system of three aromatic protons; furthermore, there were two up field triplets that corresponded to the methylene groups of the lateral propionic chain and a singlet (3H, δ 3.90 ppm), that corresponded to the methoxy group, agreeing with the report by Saha *et al.* (2003).

Other biotransformation products were detected using TLC and GC-MS analyses and they were identified by comparison with the NIST 2002 mass spectral library. Both *C. acutatum* and *L. theobromae* produced the compounds 4-ethylguaiaicol (I), vanillin (III), and vanillic acid (IV). Only the extract of *L. theobromae* had homovanillic alcohol (VI). The mass fragmentation data of the detected metabolic products corresponded to: 4-ethylguaiaicol (I) ($R_t = 12.07$ min; m/z [int. rel.]: 152[30] (M^+), 137[100] ($M^+ - CH_3$), 122[10] ($M^+ - C_2H_6$), 109[3], 91[3]); vanillin (III) ($R_t = 15.97$ min; m/z [int. rel.]: 152[90] (M^+), 151[100] ($M^+ - H$), 137[7] ($M^+ - CH_3$), 123[15] ($M^+ - CO - H$), 109[16] ($M^+ - CH_2CO - H$), 93[3]; vanillic acid (IV) ($R_t = 17.19$ min; m/z [int. rel.]: 168[77] (M^+), 153[100] ($M^+ - CH_3$), 135[7], 125[43], 110[11], 93[70] ($M^+ - CO_2 - H_2CO - H$)); and homovanillyl alcohol (VI) ($R_t = 18.84$ min; m/z [int. rel.]: 168 (M^+) [32], 137[100] ($M^+ - H_2CO - H$), 122 [11], 107[3], 94 [7]).

Among the isolated and identified compounds produced by both fungi, those that had flavoring properties were notable. The commercial value of the compounds (I) and (II) is almost 40 times higher than that of ferulic acid (Mishra *et al.*, 2014). The compound 4-ethylguaiaicol (I)

possesses a spicy to bland odor; it is responsible for the aroma and flavor of Belgian wheat and German Rauch beers (Mathew and Abraham, 2004; Priefert *et al.*, 2001), it has also been used in soy sauce and wine and as a fragrance in the perfume industry (Krings *et al.*, 2001; Priefert *et al.*, 2001). Vanillin (III) is a highly appreciated aromatic compound in the world and is used as flavoring for food and drinks and is used in pharmaceutical products. Additionally, it has antioxidant, preservative, antimicrobial and antimutagenic properties (Zhao and Moghadasian, 2008; Cerrutti and Alzamora 1996; Shaughnessy *et al.*, 2001). The vanilla flavor, that is to say vanillin, is obtained from the plant *Vanilla planifolia* as gluco-vanillin (Daugisch and Pastore, 2005). A large number of microorganisms has been used for the bioconversion of FA in vanillin, including Gram-negative bacteria of the *Pseudomonas* genus (Civolani *et al.*, 2000; Plaggenborg *et al.*, 2003), actinomycetes of the *Amycolatopsis* and *Streptomyces* genera (Achterholt *et al.*, 2000), Gram-positive bacteria such as *Bacillus subtilis* (Plaggenborget *et al.*, 2001) and *Rhodococcus* sp. (Plaggenborg *et al.*, 2006) and the basidiomycete fungus *Pycnoporus cinnabarinus* (Lesage-Meessen *et al.*, 1996). For its part, acetovanillone (V) has been reported as a minor metabolite of the degradation of ferulic acid by some microorganisms (Krings *et al.*, 2001; Priefert *et al.*, 2001). Furthermore, acetovanillone, or apocynin as it is commonly known, is a medicament isolated from the medicinal root of *Picroriza kurroa* (Lapperre *et al.*, 1999); it has numerous therapeutic applications and is a potential inhibitor of the formation of NADPH and peroxynitrite, which have damaging effects on human tissue (Lafeber *et al.*, 1999; Muijsers *et al.*, 2000).

Time-course experiments for the biotransformation of FA with *L. theobromae* and *C. acutatum*

In order to evaluate the relative abundance of the substrate and the metabolic products over time, the FA was incubated with the microorganisms *L. theobromae* and *C. acutatum* for 312 and 360 h, respectively. Daily, the culture medium of each Erlenmeyer flask was removed and extracted with EtOAc. Subsequently, the extracts were analyzed with TLC and GC. Figure 3 shows how the substrate was converted by *L. theobromae* mainly into the metabolite (II) and other minor products [(I), (III), (V), and (VI)]. The FA was essentially transformed into (II), reaching a relative

abundance of 95% in the first 24 h; afterwards, the concentration slowly decreased to 80% at 312 h. For their part, the compounds (I), (III) and (VI) increased in relative abundance after 24 h, reaching their highest concentration at 72 h. During the biotransformation

(>72 h), the quantities of (III) and (VI) remained approximately constant, under 2 and 4%, respectively; while the compound (V) had a significant increase after 168 h and its relative abundance increased to 13% (312 h).




Figure 3. Evolution over time of the products of the FA biotransformation with *L. theobromae*. The left axis correspond to relative abundance of 4-vinylguaiacol.

Figure 4 demonstrates how *C. acutatum* transformed the substrate into two major metabolites [(I) and (II)] and some minor products [(III), (IV), (V)]. Under the previously described conditions, the transformation of FA mainly produced the compound (II), reaching a relative concentration close to 90% (24 h), which subsequently decreased to 70% (360 h), while (I) reached its maximum

abundance at 48 h (~20%). In addition, (III) appeared after 48 h, reaching its highest concentration at 144 h; this fact coincides with the decrease in (I) and (II). The compound (IV) had an appreciable increase after 144 h and its relative abundance increased until reaching 10% at 360 h. Meanwhile, (V) was produced at low concentrations and its concentration was always below 2%.




Figure 4. Evolution over time of the products of the FA biotransformation with *C. acutatum*. The left axis correspond to relative abundance of 4-vinylguaiacol.

When comparing the progress of the biotransformation, it appeared that both microorganisms transformed the FA mainly into the compound 4-vinylguaiacol (II), with the highest abundance obtained with *L. theobromae*; with this microorganism, the highest proportion of acetovanillone (V) was also reached in the final period of the transformation. The elevated conversion of FA into (II) demonstrated the high efficiency of both fungi for carrying out the non-oxidative decarboxylation process. In the fermentation with *C. acutatum*, the content of 4-ethylguaiacol (I) was higher than that obtained with *L. theobromae*; while homovanillyl alcohol (VI) and vanillic acid (IV) were only detected when using *L. theobromae* and *C. acutatum* as the biocatalyzer, respectively.

Integrated metabolic pathway of FA in *C. acutatum* and *L. theobromae*

A possible metabolic pathway for the biotransformation of FA with the fungi *C. acutatum* and *L. theobromae*

(Figure 5) was established with the structure of the obtained products and the experiments over time.

Both microorganisms have the ability to carry out non-oxidative decarboxylation reactions on FA in order to produce the major compound (II), via ferulic acid decarboxylase (Donaghy *et al.*, 1999). A mechanism for the decarboxylation catalyzed by ferulic acid decarboxylase comprises the enzymatic isomerization of FA to a quinoid intermediate which is then decarboxylated spontaneously (Huang *et al.*, 1993). The decarboxylation of FA can form part of a detoxification system in order to maintain the level of fungitoxic compounds under a threshold concentration; a similar mechanism has been observed in the secondary metabolism of phenylpropenoic acid (Seshime *et al.*, 2005). Enzymatic studies related to the decarboxylation of FA to produce (II) have mainly been carried out with bacteria and yeasts (Mishra *et al.*, 2014; Priefert *et al.*, 2001). High concentrations of 4-vinylguaiacol were obtained by Karmakar *et al.* (2000)




Figure 5. Possible metabolic pathway of FA with the fungi *C. acutatum* and *L. theobromae*.

and Mathew *et al.* (2007) in cultures of *Bacillus coagulans* BK07 (908 mg L⁻¹) and *Debaryomyces hansenii* (1470 mg L⁻¹). The subsequent hydrogenation of (II) with the enzyme vinylphenol reductase (Godoy *et al.*, 2008) resulted in 4-ethylguaiacol (I), which had its highest proportion when *C. acutatum* was used. On the other hand, minor compounds, such as vanillin (III) and vanillic acid (IV), are

probably obtained through β -oxidation of FA that occurs by a mechanism analogous to the β -oxidation of fatty acids (Rosazza *et al.*, 1995). Di Gioia *et al.* (2011) proposed that FA is activated by the enzyme feruloyl-CoA synthetase (EC 6.2.1.34) to produce feruloyl-CoA; subsequently, the thioester is hydrated and cleaved by the enzyme enoyl-CoA hydratase/aldolase (EC 4.2.1.101), resulting in vanillin and

acetyl-CoA. Alternatively, the elimination of one acetate unit from the lateral chain (a retro-aldol elimination) to afford vanillin is one of the more common routes for the metabolism of FA by bacteria, yeasts, and fungi (Rosazza *et al.*, 1995). Afterwards, the enzymatic action of vanillin dehydrogenase on vanillin generates vanillic acid (IV). On the other hand, Priefert *et al.* (2001) indicated that (III) is originated from (II) through hydration, via the intermediate 4-(1'-hydroxy) ethylguaiaicol and the subsequent elimination of the methyl group. In this way, acetovanillone (V) can come from 4-(1'-hydroxy) ethylguaiaicol through the oxidation. Meanwhile, dihydroferulic acid (VII) is originated from the reduction of the double bond C-C of FA. Finally, homovanillyl alcohol (VI) may result from the hydration of (II) or from reduction of (VII).

CONCLUSIONS

The results suggest that the filamentous phytopathogenic fungi *C. acutatum* and *L. theobromae* have the ability to decarboxylize the lateral chain of ferulic acid as the principal metabolic pathway. In the process, 4-vinylguaiaicol is a principal product, which reaches its highest abundance in the first 48 h of the biotransformation. Other minor metabolites, such as acetovanillin, vanillin, and ethylguaiaicol, were also detected. These compounds have great commercial importance in the industries of flavorings and fragrances. For this reason, *C. acutatum* and *L. theobromae* have considerable potential as biocatalyzers for the production of 4-vinylguaiaicol. Nevertheless, further studies are needed to clarify the optimal conditions and the enzymes that are involved.

REFERENCES

- Achterholt, S., H. Priefert and A. Steinbüchel. 2000. Identification of *Amycolatopsis*, sp. strain HR167 genes, involved in the bioconversion of ferulic acid to vanillin. *Applied Microbiology and Biotechnology* 54(6): 799–807. doi:10.1007/s002530000431.
- Baqueiro-Peña, I., G. Rodríguez-Serrano, E. González-Zamora, C. Augur, O. Loera and G. Saucedo-Castañeda. 2010. Biotransformation of ferulic acid to 4-vinyl guaiaicol by a wild and a diploid strain of *Aspergillus niger*. *Bioresource Technology* 101(12): 4721–4724. doi:10.1016/j.biortech.2010.01.086.
- Bernini, R., E. Mincione, M. Barontini, G. Provenzano and L. Setti. 2007. Obtaining 4-vinylphenols by decarboxylation of natural 4-hydroxycinnamic acids under microwave irradiation. *Tetrahedron* 63(39): 9663–9667. doi: 10.1016/j.tet.2007.07.035.
- Bicas, J.L., P. Fontanille, G. M. Pastore and C. Larroche. 2010. A bioprocess for the production of high concentrations of R-(+)- α -terpineol from R-(+)-limonene. *Process Biochemistry* 45(4): 481–486. doi: 10.1016/j.procbio.2009.11.007.
- Bonnin, E., L. Lesage-Meessen, M. Asther and J.F. Thibault. 1999. Enhanced bioconversion of vanillic acid into vanillin by the use of natural cellobiose. *Journal of the Science of Food Agriculture* 79(3): 484–486. doi: 10.1002/(SICI)1097-0010(19990301)79:3<484::AID-JSFA271>3.0.CO;2-H.
- Bustillo, A.J., C.M. García-Pajón, J. Aleu, R. Hernández-Galán and I.G. Collado. 2003. Studies on biotransformation of (\pm)-1-(4'-chlorophenyl)-2-phenylethanol. *Tetrahedron: Asymmetry* 14(23): 3755–3760. doi: 10.1016/j.tetasy.2003.08.026.
- Cerrutti, P. and S.M. Alzamora. 1996. Inhibitory effects of vanillin on some food spoilage yeasts in laboratory media and fruit purées. *International Journal of Food Microbiology* 29(2–3): 379–386. doi:10.1016/0168-1605(95)00026-7.
- Civolani, C., P. Barghini, A.R. Roncetti, M. Ruzzi and A. Schiesser. 2000. Bioconversion of ferulic acid into vanillic acid by means of a vanillate-negative mutant of *Pseudomonas fluorescens* Strain BF13. *Applied and Environmental Microbiology* 66(6): 2311–2317. doi:10.1128/AEM.66.6.2311-2317.2000.
- Daugusch, A. and G. Pastore. 2005. Production of vanillin: a biotechnological opportunity. *Química Nova* 28(4): 642–645. doi: 10.1590/S0100-40422005000400017.
- Di Gioia, D., F. Luziatelli, A. Negroni, A.G. Ficca, F. Fava, and M. Ruzzi. 2011. Metabolic engineering of *Pseudomonas fluorescens* for the production of vanillin from ferulic acid. *Journal of Biotechnology* 156(4): 309–316. doi: 10.1016/j.jbiotec.2011.08.014.
- Donaghy, J.A., P.F. Kelly and A. McKay. 1999. Conversion of ferulic acid to 4-vinyl guaiaicol by yeasts isolated from unpasteurised apple juice. *Journal of the Science of Food and Agriculture* 79(3): 453–456. doi:10.1002/(SICI)1097-0010(19990301)79:3<453::AID-JSFA284>3.0.CO;2-H.
- Gallage, N.J. and B.L. Møller. 2015. Vanillin—bioconversion and bioengineering of the most popular plant flavor and its de novo biosynthesis in the vanilla orchid. *Molecular Plant* 8(1): 40–57. doi: 10.1016/j.molp.2014.11.008.
- Godoy, L., C. Martínez, N. Carrasco and M.A. Ganga. 2008. Purification and characterization of a p-coumarate decarboxylase and a vinylphenol reductase from *Brettanomyces bruxellensis*. *International Journal of Food Microbiology* 127(1–2): 6–11. doi: 10.1016/j.ijfoodmicro.2008.05.011.
- Hu, H., L. Li and S. Ding. 2015. An organic solvent-tolerant phenolic acid decarboxylase from *Bacillus licheniformis* for the efficient bioconversion of hydroxycinnamic acids to vinyl phenol derivatives. *Applied Microbiology and Biotechnology* 99(12): 5071–5081. doi: 10.1007/s00253-014-6313-3.
- Huang, Z., L. Dostal and J.P.N. Rosazza. 1993. Mechanisms of ferulic acid conversions to vanillic acid and guaiaicol by *Rhodotorula rubra*. *The Journal of Biological Chemistry* 268(32): 23954–23958.
- Karmakar, B., R.M. Vohra, H. Nandanwar, P. Sharma, K.G. Gupta and R.C. Sobti. 2000. Rapid degradation of ferulic acid via 4-vinylguaiaicol and vanillin by a newly isolated strain of *Bacillus coagulans*. *Journal of Biotechnology* 80(3): 195–202. doi: 10.1016/S0168-1656(00)00248-0.
- Krings, U., S. Pilawa, C. Theobald and R.G. Berger. 2001. Phenyl propenoic side chain degradation of ferulic acid by *Pycnoporus cinnabarinus*—elucidation of metabolic pathways using [5-2H]-ferulic acid. *Journal of Biotechnology* 85(3): 305–314. doi: 10.1016/S0168-1656(00)00396-5.
- Lafeber, F.P., C.J. Beukelman, E. van den Worm, J.L. van Roy, M.E. Vianen, J. van Roon, H. van Dijk and J.W. Bijlsma. 1999. Apocynin, a plant-derived, cartilage-saving drug, might be useful in the treatment of rheumatoid arthritis. *Rheumatology* 38(11): 1088–1093. doi:10.1093/rheumatology/38.11.1088.
- Lapperre, T.S., L.A. Jiménez, F. Antonicelli, E.M. Drost, P.S. Hiemstra, J. Stolk, W. MacNee and I. Rahman. 1999. Apocynin

increases glutathione synthesis and activates AP-1 in alveolar epithelial cells. *FEBS Letters* 443(2): 235–239. doi: 10.1016/S0014-5793(98)01723-2.

Lesage-Meessen, L., M. Delattre, M. Haon, J.F. Thibault, B.C. Ceccaldi, P. Brunerie and M. Asther. 1996. A two-step bioconversion process for vanillin production from ferulic acid combining *Aspergillus niger* and *Pycnoporus cinnabarinus*. *Journal of Biotechnology* 50(2-3): 107–113. doi: 10.1016/0168-1656(96)01552-0.

Luo, J.R., H.E. Jiang, Y.X. Zhao, J. Zhou and J.F. Qian. 2008. Components of the heartwood of *Populus euphratica* from an ancient tomb. *Chemistry of Natural Compounds* 44(1): 6–9. doi: 10.1007/s10600-008-0003-2.

Mathew, S., T.E. Abraham and S. Sudheesh. 2007. Rapid conversion of ferulic acid to 4-vinyl guaiacol and vanillin metabolites by *Debaryomyces hansenii*. *Journal of Molecular Catalysis B: Enzymatic* 44(2): 48–52. doi: 10.1016/j.molcatb.2006.09.001.

Mathew, S. and T.E. Abraham. 2004. Ferulic acid: an antioxidant found naturally in plant cell walls and feruloyl esterases involved in its release and their applications. *Critical Reviews in Biotechnology* 24(2-3): 59–83. doi: 10.1080/07388550490491467.

Mishra, S., A. Sachan, A.S. Vidyarthi and S.G. Sachan. 2014. Transformation of ferulic acid to 4-vinyl guaiacol as a major metabolite: a microbial approach. *Reviews in Environmental Science and Bio-Technology* 13(4): 377–385. doi: 10.1007/s11157-014-9348-0.

Muijsers, R.B.R., E. van den Worm, G. Folkerts, C.J. Beukelman, A.S. Koster, D.S. Postma and F.P. Nijkamp. 2000. Apocynin inhibits peroxynitrite formation by murine macrophages. *British Journal of Pharmacology* 130(4): 932–936. doi: 10.1038/sj.bjp.0703401.

Plaggenborg, R., J. Overhage, A. Steinbüchel and H. Priefert. 2003. Functional analyses of genes involved in the metabolism of ferulic acid in *Pseudomonas putida* KT2440. *Applied Microbiology and Biotechnology* 61(5-6): 528–535. doi: 10.1007/s00253-003-1260-4.

Plaggenborg, R., A. Steinbüchel and H. Priefert. 2001. The coenzyme A-dependent, non-beta-oxidation pathway and not direct deacetylation is the major route for ferulic acid degradation in *Delftia acidovorans*. *Microbiology Letters* 205(1): 9–16. doi: 10.1111/j.1574-6968.2001.tb10918.x

Plaggenborg, R., J. Overhage, A. Loos, J. Archer, P. Lessard, A. Sinskey and A. Steinbüchel. 2006. Potential of *Rhodococcus* strains for biotechnological vanillin production from ferulic acid and eugenol. *Applied Microbiology and Biotechnology* 72(4): 745–755. doi: 10.1007/s00253-005-0302-5.

Priefert, H., J. Rabenhorst and A. Steinbüchel. 2001. Biotechnological production of vanillin. *Applied Microbiology and Biotechnology* 56(3-4): 296–314. doi: 10.1007/s002530100687.

Rosazza, J.P.N., Z. Huang, L. Dostal, T. Volm and B. Rousseau. 1995. Review: Biocatalytic transformations of ferulic acid: An abundant aromatic natural product. *Journal of Industrial Microbiology and Biotechnology* 15(6): 457–471. doi:10.1007/BF01570016.

Saha, S., R.M. Smith, E. Lenz and I.D. Wilson. 2003. Analysis of a ginger extract by high-performance liquid chromatography coupled to nuclear magnetic resonance spectroscopy using superheated deuterium oxide as the mobile phase. *Journal of Chromatography A* 991(1): 143–150. doi:10.1016/S0021-9673(03)00215-2.

Sarma, B.K. and U.P. Singh. 2003. Ferulic acid may prevent infection of *Cicer arietinum* by *Sclerotium rolfsii*. *World Journal of Microbiology and Biotechnology* 19(2): 123–127. doi: 10.1023/A:1023205522032.

Seshime, Y., P.R. Juvvadi, I. Fujii and K. Kitamoto. (2005). Genomic evidences for the existence of a phenylpropanoid metabolic pathway in *Aspergillus oryzae*. *Biochemical and Biophysical Research Communications* 337(3): 747–751. doi:10.1016/j.bbrc.2005.08.233.

Shanker, K.S., K.H. Kishore, S. Kanjilal, S. Misra, U.S.N. Murthy and R.B.N. Prasad. 2007. Biotransformation of ferulic acid to acetovanillone using *Rhizopus oryzae*. *Biocatalysis and Biotransformation* 25(1): 109–112. doi: 10.1080/10242420601141721.

Serra, S., C. Fuganti and E. Brenna. 2005. Biocatalytic preparation of natural flavours and fragrances. *Trends in Biotechnology* 23(4): 193–198. doi: 10.1016/j.tibtech.2005.02.003.

Shaughnessy, D.T., R.W. Setzer and D.M. DeMarini. 2001. The antimutagenic effect of vanillin and cinnamaldehyde on spontaneous mutation in *Salmonella* TA104 is due to a reduction in mutations at GC but not AT sites. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 480–481(1): 55–69. doi:10.1016/S0027-5107(01)00169-5.

Topakas, E., E. Kalogeris, D. Kekos, B.J. Macris and P. Christakopoulos. 2003. Bioconversion of ferulic acid into vanillic acid by the thermophilic fungus *Sporotrichum thermophile*. *LWT Food Science and Technology* 36(6): 561–565. doi:10.1016/S0023-6438(03)00060-4.

Tsujiyama, S. and M. Ueno. 2008. Formation of 4-vinyl guaiacol as an intermediate in bioconversion of ferulic acid by *Schizophyllum commune*. *Bioscience, Biotechnology and Biochemistry* 72(1): 212–215. doi:10.1271/bbb.60606.

Velasco, R., J.H. Gil, C.M. García and D.L. Durango. 2010. Production of 2-Phenylethanol in the Biotransformation of Cinnamyl Alcohol by the Plant Pathogenic Fungus *Colletotrichum acutatum*. *Vitae* 17(3): 272–280.

Velasco, R., J.H. Gil, C.M. García and D.L. Durango. 2012. Structural modification of trans-cinnamic acid using *Colletotrichum acutatum*. *Revista Facultad de Ingeniería Universidad de Antioquia* 63(1): 20–29.

Velasco, R., D.L. Montenegro, J.F. Vélez, C.M. García and D.L. Durango. 2009. Biotransformación de Compuestos Aromáticos Sustituídos Mediante Hongos Filamentosos Fitopatógenos de los Géneros *Botryodiplodia* y *Colletotrichum*. *Revista de la Sociedad Química de Perú* 75(1): 94–111.

Velasco, R., I.D. Valverde, D.L. Durango and C.M. García. 2007. Biotransformación de los Compuestos 2- Feniletanol y Acetofenona Mediante el Hongo Fitopatógeno *Botryodiplodia theobromae*. *Vitae* 14(2): 43–50.

Xu, P., D. Hua and C. Ma. 2007. Microbial transformation of propenylbenzenes for natural flavour production. *Trends in Biotechnology* 25(12): 571–576. doi:10.1016/j.tibtech.2007.08.011.

Zhao, Z. and M.H. Moghadasian. 2008. Chemistry, natural sources, dietary intake and pharmacokinetic properties of ferulic acid: A review. *Food Chemistry* 109(4): 691–702. doi:10.1016/j.foodchem.2008.02.039.