



Revista Facultad Nacional de Agronomía
- Medellín

ISSN: 0304-2847

rfnagron_med@unal.edu.co

Universidad Nacional de Colombia
Colombia

Maradiaga Rodriguez, Walter Danilo; Pego Evangelista, Adão Wagner; Alves Junior, Jose; Vinicius Honorato, Marcus

Growing of coffee seedlings on different substrates and fertilized with lithothamium

Revista Facultad Nacional de Agronomía - Medellín, vol. 70, núm. 2, 2017, pp. 8177-8182

Universidad Nacional de Colombia

Medellín, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=179951188007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Growing of coffee seedlings on different substrates and fertilized with lithothamnium

Producción de plántulas café cultivado en diferentes tipos de sustratos y fertilizado con lithothamnium

doi: 10.15446/rfna.v70n2.64522

Walter Danilo Maradiaga Rodríguez^{1*}, Adão Wagner Pego Evangelista², Jose Alves Junior² and Marcus Vinicius Honorato²

ABSTRACT

Keywords:

Coffea arabica L.
Filter cake
Cattle manure
Nursery
Fertilizing

The coffee seedling production quality is essential to ensure crop development in field. Thus, the objective of this study was to evaluate the effect of two substrates (cattle manure and filter cake) interacting with four levels of organic fertilizer with lithothamnium in coffee seedlings var. Lempira production. A completely randomized experimental design was used with factorial arrangement of treatments 5 x 2, five levels of lithothamnium (0, 1.75, 3.50, 5.25, and 7 kg m⁻³ of substrate), two substrates (cattle manure and filter cake) and three replicates. The variables evaluated were: plant height, number of leaves, stem diameter, and root length. The treatments produced significant effects on the initial development parameters of the coffee seedlings. The best results in plants grown were obtained in cattle manure substrate with 5.25 kg m⁻³ of lithothamnium, and decreased with levels higher than 1.75 kg m⁻³ of lithothamnium in filter cake substrate.

RESUMEN

Palabras claves:

Coffea arabica L.,
Torta de filtro
Estiércol de corral
Vivero
Fertilización

La producción de plantas vigorosas constituye un requisito indispensable para garantizar el desarrollo de los cultivos en el campo definitivo. Así, el presente trabajo tuvo como objetivo evaluar el efecto de dos sustratos (estiércol de corral y torta de filtro) y cuatro dosis de fertilizante orgánico con lithothamnium en la producción de plántulas de café var. Lempira. Se utilizó un diseño experimental completamente al azar con arreglo factorial de los tratamientos 5 x 2: 5 dosis de lithothamnium (0, 1,75, 3,50, 5,25 y 7 kg m⁻³ de sustrato), dos sustratos (estiércol de corral y torta de filtro) y tres repeticiones. Las variables evaluadas fueron: altura de la planta, número de hojas, diámetro de tallo y crecimiento de raíz. Los tratamientos produjeron efectos significativos en los parámetros de desarrollo inicial de las plántulas de café. Los mejores resultados en plantas cultivadas se obtuvieron en el sustrato de estiércol con 5,25 kg m⁻³ de lithothamnium y disminuyeron con dosis superiores a 1,75 kg m⁻³ de lithothamnium en sustrato de torta de filtro.

¹ Escola de Agronomia. Universidade Federal de Goiás. Caixa Postal 131. CEP 74690-900. Goiânia, GO, Brasil.

² Núcleo de Pesquisas e Recursos Hídricos do Cerrado. Escola de Agronomia. Universidade Federal de Goiás. Caixa Postal 131. CEP 74690-900. Goiânia, GO, Brasil.

* Corresponding author: <wmaradiaga@unag.edu.hn>

In recent years, there has been a significant expansion of the market for organic products (Fibl and Ifoam, 2015). The socio cultural development of modern societies has led to the emergence of new consumption patterns in the food market (Rezende *et al.*, 2010). The awareness of the need to preserve the environment and the growing demands for healthy and quality food have potentiated the organic-ecological production. The high cost of conventional agriculture has led several farmers to rethink their way of producing up to the use of innumerable sources of organic waste as alternatives to fertilization (Rezende *et al.*, 2010).

The need for the coffee sector, to increase the productive efficiency accompanied by reduction of costs of production, for greater competitiveness, causes that the necessity arises to look for new alternatives of production.

It is known that the production of coffee quality seedlings is an important factor, and even limiting the crop productivity. The evaluation of the seedlings quality in the nursery can be a tool to identify their adequate development and if they are healthy with the maximum potential for survival after the transplant to the field (Amaral *et al.*, 2011). Among the factors that interfere in the production of quality seedlings, the substrate and fertilization, can affect the seedlings growth and development, both in the nursery and in the implantation of field crops (Marcuzzo *et al.*, 2005).

Thus, it is fundamental to select the best substrate and fertilization for the optimal seedlings development, especially in the root system development. Although the coffee root system has its developmental characteristics linked primarily to plant genetics, other factors can also modify its spatial distribution, such as soil moisture and the availability of nutrients (Souza *et al.*, 2013).

The search for new agricultural inputs is of great importance for agriculture and environmental sustainability (Evangelista *et al.*, 2013). Thus, it is important to know the effects of organic fertilizers on the availability of nutrients to plants, such as filter cake and lithothamnium. The filter cake is a residue composed of the mixture of ground sugarcane bagasse and sludge from the decantation, being derived from the sugar clarification process. In

each ton of sugarcane, 30 kg to 40 kg of filter cake are produced. It is an organic compound rich in calcium, nitrogen and potassium, with a variable composition, depending on the sugarcane variety and its maturity (González *et al.*, 2014). However, lithothamnium is a product derived from calcareous marine algae, which contains calcium carbonate, magnesium and even more than 20 microelements present in varying amounts, such as Fe, Mn, B, Ni, Cu, Zn, Mo, Se. The main characteristics that potentiate the performance of this product are attributed to the availability of the micronutrients that are adsorbed on the cell walls, being thus easily assimilable by plants and animals and the high porosity of the algae (> 40%), which provides a greater specific surface of action in soil (Dias, 2000).

The use of limestone algae (Lithothamnium), applied in a consortium with filter cake increases the availability of nutrients to the plants, due to the pH correction and the activation of the autotrophic bacteria development responsible for the nitrification process (Oliveira *et al.*, 2009). Studies carried out with this product have shown good results in the formation of yellow passion fruit and papaya seedlings (Mendoza *et al.*, 2006; Hafle *et al.*, 2009), 'Cleopatra' tangerine tree (Cruz *et al.*, 2008) and 'Swingle' citrumelo development (Araújo *et al.*, 2007). However, little is known about the application of Lithothamnium in the production of coffee seedlings.

The objective of this study was to evaluate the effect of two substrates (cattle manure and filter cake) interacting with four levels of organic fertilizer with lithothamnium in coffee seedlings production.

Materials and methods

The work was developed in the experimental area of the School of Agronomy, Federal University of Goiás (UFG), located in the geographical coordinates of 16° 35' South and 49° 16' West and 722 m elevation.

Two types of substrates were tested and five lithothamnium levels. Substrates: (i) conventional substrate composed by ravine soil classified as Oxisol, decomposed cattle manure and sand in proportions: 3:1:1 (weight/weight), adding 1 kg of superphosphate, 3 kg of potassium chloride and 2 kg of limestone per m³ of substrate; (ii) alternative substrate composed by ravine soil, filter cake, in proportions of 4:1

(weight/weight), adding 1 kg of superphosphate, 3 kg of potassium chloride and 2 kg of limestone for m³ of substrate.

The experimental design was completely randomized, with treatments in factorial arrangement 5 x 2, five lithothamnium levels (0, 1.75, 3.50, 5.25 and 7 kg m⁻³ of substrate), two substrates (x and y) and three replicates.

Soil and filter cake physical and chemical attributes are shown in Tables 1 and 2, respectively.

The seedlings were grown in plastic bags with 0.20 m high, 0.11 m wide and 0.07 m diameter, and were placed under a screen of 50% shading to avoid sunburn from excess insolation. The irrigation was performed daily to maintain soil moisture in field capacity.

Table 2. Chemical composition of the filter cake used in the experiment.

Chemical analysis					
Macronutrients	Unit	Value	Micronutrient and others	Unit	Value
P	kg ha ⁻¹	0.28	Al	kg ha ⁻¹	291.74
K	kg ha ⁻¹	90	Zn	kg ha ⁻¹	4.20
Ca	kg ha ⁻¹	1.12	MO	%	16
Mg	kg ha ⁻¹	72.94	Carbono	kg ha ⁻¹	9.28
CTC	-	4.42	pH	-	6.40
Physical analysis					
Sand	g kg ⁻¹	590	Silt	g kg ⁻¹	90
Clay	g kg ⁻¹	320	Bulk density	Mg m ⁻³	1.2

Table 2. Chemical composition of the filter cake used in the experiment.

Element	Unit	Value
N	g kg ⁻¹	13.7
P	g kg ⁻¹	15.7
K	g kg ⁻¹	8.8
Ca	g kg ⁻¹	14
Mg	g kg ⁻¹	4.8
Cu	mg kg ⁻¹	58
Mn	mg kg ⁻¹	431
Zn	mg kg ⁻¹	36.1
OM	kg ⁻¹	29
C/N	-	12.03
pH	-	7.9

The plants morphological characteristics were evaluated monthly from July to November 2015: plant height, leaf number, stem diameter, substrate saline activity, and root length. After 120 days the plants were removed from the bags. Both plant height and root length were measured as soon as they were cut and separated.

Statistical analyzes were performed using SISVAR 3.01 software. Polynomial regression analyzes were applied for fertilization levels, and deriving the equations, the best fertilization level were determined to obtain the maximum growth of coffee seedlings. And, it was use Tukey test ($P>0.05$) to substrates.

Results and discussion

The results of the statistical analysis are presented in Table 3. It is observed that the plant height was influenced by both substrate and organic fertilization (lithothamnium) levels. The root length and saline soil activity were not influenced by the substrate, but by


interaction substrates and lithothamnium levels. The stem diameter and number of leaves were not influenced by the treatments.

Coffee seedlings that were planted with cattle manure had higher plant height than with filter cake (Figure 1).

Table 3. Significant *P*-values of plant height (PIH), stem diameter (StD), number of leaves (NL), and root length (RL) of the coffee seedlings to different substrates and lithothamnium levels.

Variation source	<i>P</i> -value				
	PIH	StD	NL	RL	AS
Substrate	0.7176 *	0.00064 ^{ns}	0.54964 ^{ns}	3.5020 ^{ns}	0.0028 ^{ns}
Fertilization level	0.5325 *	0.07424 ^{ns}	1.24749 ^{ns}	10.075 *	0.3222 *
Substrate x F. Level	0.2878 *	0.08745 ^{ns}	0.64394 ^{ns}	9.6776 *	0.8623 *
Error	0.092	0.06009	3.17315	1.0567	0.1012
CV (%)	4.6	11.4	27.9	6.65	30.42

* Significant at 5% error probability; ^{ns} non-significant by the F test.



*Means followed by different letters differ by Tukey test at 5% error probability.

Figure 1. Plant height mean (PIH) of coffee seedlings developed in two substrates (filter cake and cattle manure)

Braun *et al.* (2009) found similar results. The authors report that the mean values of plant height produced with cattle manure were higher than with commercial substrate (Plantmax®) + soil. This result is probably due to the greater amount of organic material added in the substrate, which can improve infiltration, water retention, aeration, temperature and root penetration (Oliveira *et al.*, 2009).

A quadratic function was better fitted to lithothamnium levels in the cattle manure substrate, and linear function

for the filter cake substrate (Figure 2). The maximum root length of coffee seedlings was obtained in treatment with filter cake fertilized with 1.75 kg of lithothamnium per m⁻³ of substrate. The fertilization with levels higher than 1.75 kg of lithothamnium per m⁻³ of substrate reduced root length. While in cattle manure the maximum root length was with 5.25 kg m⁻³ of lithothamnium reducing at higher levels. According to Evangelista *et al.* (2015) high levels of lithothamnium cause negative effect on plant growth, due to the increase in pH of the soil and inhibition of nutrient absorption by plants. This can be




Figure 2. Regression analysis for the root length of coffee seedlings to different substrates and lithothamnium levels. A) Cattle manure. B) Filter cake.

justified by the possibility that the micronutrients present in lithothamnium, interacting with phosphorus and nitrogen in the filter cake, are given gradually by mineralization and attack of microorganisms in the soil, which causes greater efficiency by the plant (Santos *et al.*, 2010). This may probably be due to smaller doses of lithothamnium, interacting with the phosphorus and the existing nitrogen in the filter cake, are slowly mineralized by the attack of microorganisms in the soil, which causes greater efficiency by the plant (Santos *et al.*, 2010). It is suggested that the application of bioclastic granulate be made along with organic matter, once organic matter enhances chemical and physical properties of the soil (Damatto *et al.*, 2006) in addition to having nutrients in its composition, which are best provided with the application of grained bioclastic (Days, 2000), favoring the coffee growth. Teixeira *et al.* (2015) and Mendoça *et al.* (2006) also evaluate the effect of application of lithothamnium levels and types of substrate on production of *Carica papaya* L. They found that high levels of this fertilizer had negative effects in the, and the authors attributed to a possible toxicity of the substrate.

CONCLUSIONS

The treatments produced significant effects on the initial development parameters of the coffee seedlings. The best results in plants grown were obtained in cattle manure substrate with 5.25 kg m⁻³ of lithothamnium, and decreased with levels higher than 1.75 kg m⁻³ of lithothamnium in filter cake substrate.

REFERENCES

- Amaral J, Martinez H, Laviola B, Fernandez E and Cruz C. 2011. Eficiência de utilização de nutrientes por cultivares de cafeeiro. *Ciência Rural* 41(2): 621-629. doi: 10.1590/S0103-84782011005000027
- Araújo E, Oliveira A, Cavalcante L, Pereira E, Brito N, Neves C and Silva E. 2007. Produção do pimentão adubado com esterco bovino e biofertilizante. *Engenharia Agrícola e Ambiental* 11(5): 466-470. doi: 10.1590/S1415-43662007000500003
- Braun H, Henrique Zonta J, Soares J, Fialho E and Paulucio D. 2009. Desenvolvimento inicial do café conillon (*coffea canephora pierre*) em solos de diferentes texturas com mudas produzidas em diferentes substratos. *Idesia* 27(3): 35-40. doi: 10.4067/S0718-34292009000300006
- Cruz M, Hafle M and Ramos P. 2008. Desenvolvimento do porta-enxerto de tangerineira "Cleópatra". *Revista Brasileira de Fruticultura* 30(2): 471-475. doi: 10.1590/S0100-29452008000200035
- Dias G. 2000. Granulados bioclásticos: algas calcárias. *Revista Brasileira de Geofísica* 18(3): 308-318. doi: 10.1590/S0102-261X200000300008
- Evangelista A, Júnior J and Melo P. 2013. Resposta do cafeeiro à aplicação de níveis de irrigação e adubação com Alfertil. *Revista Brasileira de Engenharia Agrícola e Ambiental* 17(4): 392-396. doi: 10.1590/S1415-43662013000400006
- Evangelista A, Alves J, Casaroli D and Resende F. 2015. Desenvolvimento inicial da mamoneira, girassol e nabo forrageiro adubados com Lithothamnium. *Global Science and Technology* 8(2): 40-48. doi: 10.14688/1984-3801/gst.v8n2p40-48
- Fibl and Ifoam- Institute of Organic Agriculture-International Foundation for Organic Agriculture. En: The World of Organic Agriculture; <https://www.fibl.org/fileadmin/documents/shop/1663-or-ganic-world-2015.pdf>. 306 p. consulta: junho 2016.
- González C, Prado M, Hernández A, Caione G and Selva E. 2014. Uso de torta de filtro enriquecida com fosfato natural e biofertilizantes em Latossolo Vermelho distrófico. *Revista Pesquisa Agropecuária Tropical* 44(2): 135-141. doi: 10.1590/S1983-40632014000200001
- Hafle O, Andrade V, Ramos J, Monterio M and Melo P. 2009. Produção de mudas de mamoeiro utilizando Bokashi e Lithothamnium.

Revista Brasileira de Fruticultura 31(1): 245-251. doi: 10.1590/S0100-29452009000100034

Marcuzzo K, Melo B, Carvalho H, Teodoro E and Alvarenga B. 2005. Desenvolvimento de mudas de cafeeiro em diferentes substratos e doses de fertilizante de liberação gradual. Bioscience Journal, 21(57): 57-63.

Mendonça V, Orbes M, Arruda N, Ramos J, Teixeira G and Antunes H. 2006. Qualidade de mudas de maracujazeiro-amarelo formadas em substratos com diferentes níveis de Lithothamnium. Ciência Agrotecnica 30(5): 900-906. doi: 10.1590/S1413-705420 06000500012

Oliveira A, Oliveira A, Leonardo P, Cruz S and Silva D. 2009. Yield of gherkin in response to doses of bovine manure. Revista Horticulura Brasileira 27(1): 100-102. doi: 10.1590/S0102-053620 09000100020

Rezende R, Junior C, Souza R, Antunes F and Frizzone J. 2010. Crescimento inicial de duas cultivares de cafeeiro em diferentes

regimes hídricos e dosagens de fertirrigação. Revista Engenharia Agrícola 30(3): 447-458. doi: 10.1590/S0100-69162010000300009

Santos H, Tiritan S, Foloni S and Fabris L. 2010. Produtividade de cana-de-açúcar sob adubação com torta de filtro enriquecida com fosfato solúvel. Pesquisa Agropecuaria Tropical 40(4): 454-461. doi: 10.5216/pat.v40i4.7272

Souza R, Helbel Júnior C, Rezende R, Costa A, Freitas P, Tavore V and Maller A. 2013. Características de crescimento inicial de duas cultivares de cafeeiro sob diferentes regimes hídricos e níveis de fertilização NPK. Revista Semina 34(6): 3141. doi: 10.5433/1679-0359.2013v34n6supl1p3141

Teixeira G, Souza H, Mendonça V, Ramos D, Chalfun J, Ferreira A and Melo P. 2015. Produção de mudas de mamoeiro “formosa” em substratos com doses de lithothamnium. Revista da FZVA 16(2): 220-229.