



Revista Brasileira de Ciência do Solo

ISSN: 0100-0683

revista@sbcs.org.br

Sociedade Brasileira de Ciência do Solo
Brasil

Ribeiro da Silva, Ivo; Côrtes Corrêa, Tarcísio Fernando; Ferreira Novais, Roberto; Oliveira Gebrim, Fabrício de; Novais Nunes, Flancer; Silva, Eulene Francisco da; Jot Smyth, Thomas
PROTECTIVE EFFECT OF DIVALENT CATIONS AGAINST ALUMINUM TOXICITY IN SOYBEAN
Revista Brasileira de Ciência do Solo, vol. 32, núm. 5, 2008, pp. 2061-2071
Sociedade Brasileira de Ciência do Solo
Viçosa, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=180214065027>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

PROTECTIVE EFFECT OF DIVALENT CATIONS AGAINST ALUMINUM TOXICITY IN SOYBEAN⁽¹⁾

Ivo Ribeiro da Silva⁽²⁾, Tarcísio Fernando Côrtes Corrêa⁽³⁾, Roberto Ferreira Novais⁽²⁾, Fabrício de Oliveira Gebrim⁽⁴⁾, Flancer Novais Nunes⁽⁴⁾, Eulene Francisco da Silva⁽⁴⁾ & Thomas Jot Smyth⁽⁵⁾

SUMMARY

A large proportion of soybean fields in Brazil are currently cultivated in the Cerrado region, where the area planted with this crop is growing considerably every year. Soybean cultivation in acid soils is also increasing worldwide. Since the levels of toxic aluminum (Al) in these acid soils is usually high it is important to understand how cations can reduce Al rhizotoxicity in soybean. In the present study we evaluated the ameliorative effect of nine divalent cations (Ca, Mg, Mn, Sr, Sn, Cu, Zn, Co and Ba) in solution culture on Al rhizotoxicity in soybean. The growth benefit of Ca and Mg to plants in an acid Inceptisol was also evaluated. In this experiment soil exchangeable Ca:Mg ratios were adjusted to reach 10 and 60 % base saturation, controlled by different amounts of CaCl_2 or MgCl_2 (at proportions from 100:0 up to 0:100), without altering the soil pH level. The low (10 %) and adequate (60 %) base saturation were used to examine how plant roots respond to Al at distinct (Ca + Mg)/Al ratios, as if they were growing in soils with distinct acidity levels. Negative and positive control treatments consisted of absence (under native soil or undisturbed conditions) or presence of lime (CaCO_3) to reach 10 and 60 % base saturation, respectively. It was observed that in the absence of Aluminum, Cu, Zn, Co and Sn were toxic even at a low concentration ($25 \mu\text{mol L}^{-1}$), while the effect of Mn, Ba, Sr and Mg was positive or absent on soybean root elongation when used in concentrations up to $100 \mu\text{mol L}^{-1}$. At a level of $10 \mu\text{mol L}^{-1}$ Al, root growth was only reverted to the level of control plants by the Mg treatment. Higher Tin doses led to a small alleviation of Al rhizotoxicity, while the other cations reduced root growth or had no effect. This is an indication that the Mg effect is ion-specific and not associated to an electrostatic protection mechanism only, since all ions were divalent and used at low concentrations. An increased exchangeable Ca:Mg

⁽¹⁾ Recebido para publicação em outubro de 2006 e aprovado em outubro de 2008.

⁽²⁾ Professor do Departamento de Solos, Universidade Federal de Viçosa – UFV. Av. PH Rolfs s/n, CEP 36570-000 Viçosa (MG). E-mails: ivosilva@ufv.br

⁽³⁾ Graduando em Agronomia, Departamento de Solos, UFV. Bolsista do CNPq.

⁽⁴⁾ Doutorando em Solos e Nutrição de Plantas, Departamento de Solos, UFV.

⁽⁵⁾ Soil Science Department, North Carolina State University, Raleigh, NC, USA

ratio (at constant soil pH) in the acid soil almost doubled the soybean shoot and root dry matter even though treatments did not modify soil pH and exchangeable Al^{3+} . This indicates a more efficient alleviation of Al toxicity by Mg^{2+} than by Ca^{2+} . The reason for the positive response to Mg^{2+} was not the supply of a deficient nutrient because CaCO_3 increased soybean growth by increasing soil pH without inducing Mg^{2+} deficiency. Both in hydroponics and acid soil, the reduction in Al toxicity was accompanied by a lower Al accumulation in plant tissue, suggesting a competitive cation absorption and/or exclusion of Al from plant tissue stimulated by an Mg-induced physiological mechanism.

Index terms: heavy metals, cation amelioration, soil acidity, Ca:Mg ratio

RESUMO: EFEITO PROTETOR DE CÁTIONS DIVALENTES CONTRA A TOXIDEX DE ALUMÍNIO EM SOJA

Uma grande parte da soja no Brasil é cultivada atualmente na região do Cerrado, e é nessa região que ocorre a maior expansão da cultura. Nessa região, a maioria dos solos é ácida e, geralmente, apresenta níveis elevados de alumínio (Al) tóxico. Portanto, é importante compreender como determinados cátions reduzem a rizotoxidez de Al em plantas de soja. Neste estudo foram avaliados, em solução nutritiva, o efeito protetor de nove cátions divalentes (Ca, Mg, Mn, Sr, Sn, Cu, Zn, Co and Ba) contra a rizotoxidez de Al e o efeito positivo do Ca e Mg no crescimento das plantas de soja num solo com elevada saturação por Al. A relação entre Ca e Mg trocáveis foi variada por meio de adição de diferentes quantidades de CaCl_2 ou MgCl_2 (com as proporções variando entre 100:0 até 0:100), sem que ocorresse alteração no pH do solo. Tratamentos testemunhas consistiram de solo sem aplicação ou com aplicação de calcário (CaCO_3). Foi observado que, na ausência do Al, os cátions Cu, Zn, Co e Sn foram tóxicos mesmo em baixas concentrações ($25 \mu\text{mol L}^{-1}$), enquanto Mn, Ba, Sr e Mg em concentrações até $100 \mu\text{mol L}^{-1}$ apresentaram nenhum efeito ou efeito positivo no alongamento radicular da soja. Com $10 \mu\text{mol L}^{-1}$ de Al, apenas no tratamento com Mg as raízes apresentaram recuperação no alongamento para o mesmo patamar que o tratamento controle sem Al. O Sn em doses elevadas proporcionou uma pequena melhora na rizotoxidez de Al, enquanto outros cátions divalentes tiveram nenhum efeito ou reduziram o crescimento radicular. Isso é uma indicação de que o efeito do Mg é específico e não está apenas associado a uma proteção por meio de efeito eletrostático, porque todos os outros cátions usados eram divalentes e foram fornecidos em baixas concentrações. No experimento com solo ácido, observou-se que a redução da relação Ca:Mg trocáveis praticamente dobrou o acúmulo de matéria seca da parte aérea e das raízes, embora os tratamentos não tenham modificados o pH do solo e o Al^{3+} trocável, suportando o maior efeito protetor do Mg contra o Al^{3+} em relação ao Ca. A resposta positiva ao Mg não foi devida ao suprimento do nutriente que se encontrava deficiente, pois a aplicação de CaCO_3 melhorou o crescimento radicular por meio do incremento do pH (e precipitação do Al^{3+}) sem induzir deficiência de Mg. Tanto em condições hidropônicas quanto no solo ácido, a redução na toxidez do Al^{3+} foi acompanhada por menor acúmulo de Al no tecido vegetal, sugerindo a ocorrência de inibição competitiva na absorção, e, ou, exclusão do Al na rizosfera em resposta a um mecanismo fisiológico promovido pelo Mg.

Termos de indexação: metais pesados, efeito protetor de cátions, acidez do solo, relação Ca:Mg.

INTRODUCTION

Soybean is one of the most important export crops grown in Brazil. A vast area of soybean fields is currently cultivated in the Cerrado region, and increases considerably every year. Acid soils with variable charge minerals (kaolinite and oxides) have

acidity levels of more than 50 % of the soils are high and aluminum (Al) levels in the surface and sub-surface layers are toxic (Eswaran, 1997; Lopes, 1983). Grain yield of several crops was shown to be reduced by base saturation lower than 60 % and enchangeable Al^{3+} greater than $0,3 \text{ cmol}_c \text{ dm}^{-3}$ (Nicolodi et al., 2008). This makes liming a key practice of soil fertility management (Lopes, 1996). A better understanding

of related issues and enhanced management will benefit this as well as other regions, since soybean cultivation in acid soils is quickly expanding worldwide.

Although there is significant variation for Al tolerance among soybean genotypes (Sartain & Kamprath, 1978; Spehar, 1994; Menosso et al., 2000; Silva et al., 2000), the species is relatively sensitive to soil acidity in comparison with other crop and tree species (Moyer-Henry et al., 2003). Several studies have reported substantial gains in growth and yield in acid soils treated with lime (Oliveira & Pavan, 1996; Martins et al., 1998; Fageria, 2001). The positive effects of liming on acid soils are numerous, and the reduction of Al toxicity is certainly one of the most beneficial. Knowledge on the amelioration of Al toxicity due to its precipitation as low solubility Al-hydroxides under higher soil pH is not new, but the effects of basic cations (mainly Ca and Mg) on the reduction of the deleterious effect of Al have almost always been overlooked. Experiments in hydroponics have clearly shown that basic cations are able to alleviate Al toxicity in soybean (Alva et al., 1986; Ferrufino et al., 2000; Silva et al., 2001a). In fact, soybean responses to liming in high Al soils with sufficient Ca and Mg availability are usually low or insignificant (Caires et al., 1998, 2003). Conversely, substantial responses to liming have been observed in soils with high Al and lower cation availability (Martins et al., 1998; Fageria, 2001).

Previous research has shown that the ability of cations to reduce Al^{3+} toxicity, except for the small cation H^+ , is directly related to their valences in the order: $\text{H}^+ > \text{C}^{3+} > \text{C}^{2+} > \text{C}^{1+}$ (Kinraide & Parker, 1987; Grauer & Horst, 1992; Kinraide et al., 1992, 1994a; Kinraide, 1998). However, it seems that the amelioration of Al toxicity by cations is not only a matter of a competitive effect with Al by absorption sites at the root surface because cations of a same valence may have a distinct, species-dependent ameliorative effect. For example, it was observed that Ca and Mg confer a similar protection against Al toxicity in wheat (Kinraide, 1998), but there are results in hydroponics indicating that Mg is more efficient than Ca in soybean (Silva et al., 2001a,b). Under field conditions it was found that Mg has a fundamental function against Al toxicity because the response of soybean yield to liming increases as a consequence of greater availability and Mg uptake (Caires et al., 2001). It is currently not known whether other divalent cations have a similar ameliorative effect as Mg^{2+} and whether Mg^{2+} is equally protective in soil as in hydroponic conditions. This study investigates the protective role of physiologically relevant concentrations of several divalent cations against Al^{3+} rhizotoxicity in soybean and evaluates whether the beneficial effects of Mg^{2+} can be extended from hydroponics to soil conditions.

MATERIAL AND METHODS

Experiment 1: Protective effect of divalent cations against Al rhizotoxicity in solution culture

Soybean seeds of the cultivars UFV-16 (Al-tolerant) and Confiança (Al-sensitive) were germinated on $0.01 \text{ mmol L}^{-1} \text{ CaSO}_4$ -soaked germination paper at 25°C for 72 h. Six to eight uniform seedlings of each cultivar were transferred to plastic trays containing 10 L of aerated solution culture with $0.5 \text{ mmol L}^{-1} \text{ CaCl}_2$. The solution pH had previously been adjusted to 4.5 with $0.1 \text{ mmol L}^{-1} \text{ H}_2\text{SO}_4$, to avoid Al precipitation.

Treatments were analyzed in a 8×2 factorial design of eight divalent cations (Mg^{2+} , Zn^{2+} , Cu^{2+} , Mn^{2+} , Sr^{2+} , Sn^{2+} , Co^{2+} and Ba^{2+}), either in the absence or presence of $10 \text{ } \mu\text{mol L}^{-1} \text{ Al}$. Due to the large number of experimental units, the experiment was run independently for the two soybean cultivars. The divalent cations were applied at 0, 25, 50 and $100 \text{ } \mu\text{mol L}^{-1}$. Background Ca levels were varied from 0.4 to 0.5 mmol L^{-1} depending on the concentration of other divalent cations added, in order to maintain a constant ionic strength. The solution pH was adjusted daily to 4.5 by slowly adding $0.1 \text{ mol L}^{-1} \text{ KOH}$ or HCl as required, under continuous stirring. The plants were grown in a greenhouse at $27 \pm 5^\circ\text{C}$. After 16–18 h acclimation to the basal CaCl_2 solution, the primary root length was measured and treatments initiated. Root length was measured again 90 h after treatment initiation. The experiments were arranged in a randomized block design, with three replicates of six seedlings each.

At harvest, roots and shoots were separated, dried for 72 h (70°C) in a forced draft oven, weighed, ground in a Wiley mill with a 1 mm stainless steel sieve and wet-digested with a nitro-perchloric mixture (3:1) according to Sarruge & Haag (1974). After appropriate dilutions the element concentration was determined by ICP-AES.

Parallel experiments were run in hydroponics to evaluate the effect of Ca and Mg on root growth in more detail, under similar conditions as described above. Additional information on treatments is presented in the section Results and Discussion.

Experiment 2: Effect of varying Ca:Mg ratio on soybean growth in an acid soil

In order to test the effectiveness of Ca and Mg as Al toxicity barrier in soil, soybean growth was evaluated in soil samples of the 0–20 cm layer of an Al-toxic sandy-loam Inceptisol with varying Ca and Mg ratios. After air-drying, sub-samples were taken for chemical and physical soil analysis. The soil consists of 18 % clay, 11 % silt, 36 % fine sand and 35 % coarse sand. Under native conditions the soil presented ($\text{cmol}_c \text{ kg}^{-1}$): 0.10 Ca, 0.04 Mg, and 1.00 Al. Except for the nutrients involved in the treatments, all macro and micronutrients were supplied in

amounts to ensure optimal growth. Fourteen treatments were arranged in a 7 x 2 factorial design of seven Ca and, or Mg amendments, at two soil base saturations (10 and 60 %). The seven Ca and Mg treatments consisted of: (1) control with all nutrients (without Ca and Mg). All nutrients except Ca and Mg were supplied in doses and sources according to Novais et al. (1991); (2) soil limed with CaCO_3 to reach 60 % base saturation (3.8 t ha^{-1}); (3) Ca + Mg equivalent to Ca present in the CaCO_3 applied in treatment 1, at 100:0 Ca:Mg; (4) Ca + Mg equivalent to Ca present in the CaCO_3 applied in treatment 1, at 75:25 Ca:Mg; (5) Ca + Mg equivalent to Ca present in the CaCO_3 applied in treatment 1, at 50:50 Ca:Mg; (6) Ca + Mg equivalent to Ca present in the CaCO_3 applied in treatment 1, at 25:75 Ca:Mg, and (7) Ca + Mg equivalent to Ca present in the CaCO_3 applied in treatment 1, at 00:100 Ca:Mg. In treatments 3 to 7 and 8 to 14 Ca and Mg were supplied as chloride salts to avoid changes in soil pH, Al precipitation and Al complexation, thus minimizing the chance for confounding effects. Treatments 8 to 14 were the same as described above, except that a lower lime dose (10 % base saturation) was used. The low (10 %) and adequate (60 %) base saturation were created to examine how plant roots would respond to Al at distinct (Ca + Mg)/Al ratios, as if they were growing in soils with distinct acidity limitations. After treatment application, the soil moisture was adjusted

to field capacity and incubated for two weeks in a greenhouse. Thereafter, it was air-dried, weighed (0.8 kg) and transferred to 1 L plastic pots. Six seeds of soybean cv. UFVS-2001 were sown in each pot and three days after germination, seedlings were thinned to the three most regular in size. Pots were watered twice a day with deionized water in order to maintain moisture at around 80 % of field capacity. Two weeks days after germination the plants (vegetative stage between V_2 and V_3) were removed from pots, separated in shoot and roots, dried at 70°C and then weighed to the fourth decimal place. The material was wet-digested in a nitro-perchloric acid mixture and the Ca, Mg and Al concentrations were measured in the extracts as described above.

RESULTS AND DISCUSSION

Protective effect of Mg and other divalent cations against Al rhizotoxicity

In the first part of this study the effect of several divalent cations on the elongation of soybean roots was evaluated in the absence and presence of Al^{3+} in solution. The differences among divalent cations regarding their effects on root elongation were marked, statistically significant ($p < 0.05$) (Figures 1 and 2). In control solutions without Al^{3+} , cations such as Cu^{2+} ,




Figure 1. Relative root elongation of soybean cultivars UFV-16 and Confiança as a function of increasing doses of divalent cations in a CaCl_2 500 mmol L^{-1} (pH 4.5) basal solution, in the absence of Al. Error bars represent standard error ($n = 3$).




Figure 2. Relative root elongation of soybean cultivars UFV-16 and Confiança as a function of increasing doses of divalent cations in a CaCl_2 500 mmol L^{-1} (pH 4.5) basal solution, in the presence of 10 $\mu\text{mol L}^{-1}$ Al. Error bars represent standard error ($n = 3$).

Zn^{2+} and Co^{2+} were highly toxic to both soybean cultivars, drastically reducing root elongation even at the lowest concentration (Figure 1). Previous studies have also shown that Cu^{2+} and Zn^{2+} at low doses were highly toxic to plant roots (Kinraide et al., 2004; Pedler et al., 2004). Tin was slightly toxic to cv. Confiança only. Although Cu^{2+} and Zn^{2+} are essential elements for plant growth, they are required at low concentrations (less than 10 $\mu\text{mol L}^{-1}$) in solution, while excess supply leads to toxicity, with a consequent reduction of plant growth. Cobalt is only considered a beneficial (non-essential) element for plants, but is essential for N_2 -fixing bacteria (Marschner, 1995). The effect on root elongation of the other cations (Mn^{2+} , Ba^{2+} , Sr^{2+} and Mg^{2+}), mainly the last three, was positive ($p < 0.05$). The positive effect of these cations is possibly related to alleviation of H^+ toxicity (Kinraide, 1991) at low pH levels, as used here, rather than a response to nutrient deficiency in the growth medium. The beneficial effects of Ba^{2+} and Sr^{2+} , which are non-essential elements for plants, seem to support this hypothesis.

The two genotypes differed ($p < 0.05$) in response to solution Al. In solutions containing 10 $\mu\text{mol L}^{-1}$ Al (in 500 $\mu\text{mol L}^{-1}$ CaCl_2 background) and with addition of no other divalent cation, root growth of cv. Confiança was reduced by approximately 75 % and around 55 %

in cv. UFV-16. This differential Al tolerance of soybean genotypes has been reported previously (Sartain & Kamprath, 1978; Spehar, 1994; Menosso et al., 2000; Silva et al., 2000) and is related to the ability of Al-tolerant plants to produce and secrete organic acids, particularly citrate, in the presence of Al (Yang et al., 2000; Menosso et al., 2001; Silva et al., 2001a).

The addition of increasing amounts of Mn^{2+} , Sr^{2+} and Ba^{2+} led to no substantial improvement in root elongation of both soybean cultivars. Cobalt, Zn^{2+} , Cu^{2+} and Sn^{2+} even aggravated the Al rhizotoxicity (Figure 2), probably as a result of their intrinsic toxicity in the absence of Al^{3+} (Figure 1). In contrast, the toxic effect of Al on root growth of both soybean cvs. was eliminated with the addition of as little as 25 $\mu\text{mol L}^{-1}$ Mg^{2+} (Figure 2). The improvement in root elongation by Mg was paralleled by a reduced Al concentration in root tissue (Figure 3h). This was not restricted to Mg since increasing doses of all cations led to a reduced Al concentration in roots, with exception of Sr^{2+} and Ca^{2+} . The former reduced Al accumulation up to a dose of 50 $\mu\text{mol L}^{-1}$ and the latter was not very effective in reducing Al accumulation in the root (Figure 3).

The high Ca^{2+} demand in the growth medium to maintain normal root elongation (Kinraide, 1998;




Figure 3. Aluminum concentration in roots of soybean cvs. UFV-16 and Confiança as a function of increasing doses of divalent cations in a CaCl₂ 500 mmol L⁻¹ (pH 4.5) basal solution, in the presence of 10 μmol L⁻¹ Al. Error bars represent standard error (n = 3).

Sanzonowicz et al., 1998; Silva et al., 2001b) negates a separate analysis of the beneficial effects of supplying a deficient nutrient from others related to the alleviation of Al damage. In the solution without Al, the absence of Ca²⁺ (deionized water, pH 4.5) completely inhibited root elongation, which reached normal growth only when at least 250 μmol L⁻¹ Ca²⁺ was present in the solution (Figure 4a). When Ca²⁺ was supplied at a non-limiting concentration (> 250 μmol L⁻¹; pH 4.5), the presence of 10 μmol L⁻¹ Al limited root elongation by more than 50 % (Figure 4a), which confirms the lower protective effect of Ca²⁺ (Figure 4a) in comparison to Mg²⁺ (Figure 5). In fact, only 500 μmol L⁻¹ Ca²⁺ was sufficient to completely alleviate the deleterious effect of 10 μmol L⁻¹ Al on soybean root growth with 25 μmol L⁻¹ Mg²⁺ in the solution.

It has been shown that strontium is able to substitute for Ca in the dimerization of pectic

compounds of the cell wall *in vitro* (O'Neill et al., 1996). The possibility of using Sr²⁺ in substitution of Ca²⁺ in solution would allow a more accurate quantification of the ameliorative effect of Ca on Al rhizotoxicity. However, Sr²⁺ alone was unable to substitute Ca²⁺, leading to restricted root elongation even in Al-free solution (Figure 4b). The protective effect of cations at low concentrations (micromolar range) against Al rhizotoxicity in soybean therefore seems to be somewhat ion-specific. The mitigating effect of Mg²⁺ is not due to an increase in the ionic strength, which was maintained constant in all treatments. Moreover, since all cations evaluated were divalent, it is unlikely that the ameliorative effect of Mg²⁺ was associated to Al displacement from the root cell surface. The low cation concentrations employed do not support the explanation of a protective action based on the increase of the electrical potential of root cell plasma membrane either (Kinsade et al.,




Figure 4. Root elongation of soybean cultivars UFV-16 and Confiança as a function of increasing Ca (a) or Sr (b) in solution (pH 4.5), in the absence or presence of 10 µmol L⁻¹ Al. Error bars represent standard error (n = 3).

1992; Kinraide, 1994b, 1998). In wheat, Pedler et al. (2004) found that low Mg²⁺ doses were able to completely alleviate Zn²⁺ rhizotoxicity, while Ca amelioration, even at much higher doses, was much weaker. The nature of this protective role of Mg²⁺ is not quite clear at the moment, but the high effectiveness of low Mg²⁺ concentrations indicates the involvement of a biochemical/physiological mechanism. One possibility could be the stimulatory function by Mg²⁺ on the biosynthesis and citric acid secretion by soybean root tips in the presence of Al (Silva et al., 2001c). Since citric acid is a potent Al chelator, it could well be involved in Al detoxification in the rhizosphere or within root tip cells (Ma et al., 2001; Ryan et al., 2001). Due to the ionic radii similarity, an Al uptake blockage due to competitive

transport (Macdiarmid & Gardner, 1996) can not be completely ruled out.

Calcium and magnesium effects on soybean growth in acid soil

We further tested the protective effect of Ca and Mg by growing soybean on an acid, Al-toxic soil under variable Ca²⁺ and Mg²⁺ availability. For the purpose of comparison, the treatments liming (plus all nutrients but Ca²⁺ and Mg²⁺) and control (no liming plus all nutrients, except Ca²⁺ and Mg²⁺) were established (Table 1). A decreasing exchangeable Ca:Mg ratio in the soil from 26 to 0.17 (Table 1) almost doubled soybean shoot and root dry matter f (Figure 6). A similar trend was observed for the treatment with smaller amounts of Ca and Mg application (10 % base saturation). Interestingly, these treatments did not substantially modify the soil pH and exchangeable Al³⁺, thus indicating a more efficient alleviation of Al toxicity by Mg²⁺ than by Ca²⁺. These results also show that the existing exchangeable Ca²⁺ in the soil was not limiting, otherwise no response to Mg²⁺ would have been observed. The positive response to Mg²⁺ was not due to a supply of a deficient nutrient because CaCO₃ increased soybean growth by increasing soil pH and precipitating toxic Al without inducing Mg²⁺ deficiency, as observed for other acid, Mg-deficient soils (Tan et al., 1992).

A low lime dose (10 % saturation) was sufficient to raise soil pH from 4.4 to 5.1 and improved shoot growth. Nevertheless, root growth was still limited under such conditions (Figure 6a). It appears that the remaining 0.6 cmol_c dm⁻³ Al³⁺ in the soil was




Figure 5. Relative root elongation of soybean cultivar Confiança as a function of increasing Ca and Mg in solution (pH 4.5), in the presence of 10 µmol L⁻¹ Al. Error bars represent standard error (n = 3).

Table 1. Selected chemical characteristics of the soil after establishing the treatments

Treatment		pH	Ca ²⁺	Mg ²⁺	Al ³⁺	Ca ²⁺ :Mg ²⁺ ratio
cmol _c dm ⁻³						
Low lime dose (10 % base saturation)						
1	100:0	4.6	0.40	0.05	1.0	8.00
2	75:25	4.8	0.30	0.10	1.0	3.00
3	50:50	4.5	0.30	0.20	0.9	1.50
4	25:75	4.7	0.20	0.20	0.9	1.00
5	0:100	4.7	0.10	0.30	0.9	0.33
6	CaCO ₃	5.1	0.50	0.05	0.6	10.00
Recommended lime dose (60 % base saturation)						
7	100:0	4.2	1.30	0.05	0.9	26.00
8	75:25	4.3	1.10	0.40	0.9	2.75
9	50:50	4.4	0.90	0.80	1.0	1.13
10	25:75	4.4	0.50	1.10	0.9	0.45
11	0:100	4.3	0.20	1.20	0.9	0.17
12	CaCO ₃	5.7	2.60	0.05	0.0	52.00
12	Control (no liming)	4.4	0.10	0.05	0.9	2.00

pH in H₂O (2.5:1). Al, Ca and Mg in the 1 mol L⁻¹ KCl extract.




Figure 6. Dry matter production by soybean cv. UFVS-2001 grown in an acid soil with variable Ca:Mg ratios. (a) 10 % base saturation; (b) 60 % base saturation. Lime treatment received CaCO₃ and all other nutrients, except Mg. Control (Ctrl) treatment received no liming, but the supply of all other nutrients, except Ca and Mg, was ensured. Treatment means followed by the same letter are not significantly different by Tukey's test ($p < 0.05$).

enough to restrict root growth. In fact, Al³⁺ concentration above 0.3 cmol_c dm⁻³ results in lower grain yield for several crops, including soybean (Nicolodi et al., 2008). Contrastingly, root elongation was increased by several orders of magnitude when Mg was added (Figure 6A), even though 0.9 cmol_c dm⁻³ Al³⁺ was still present (Table 1). The better root development at 10 % than that at 60 % base saturation in treatments with chloride salts may be due to chloride toxicity since the soil was a sandy loam and soybean is known to be chloride-sensitive (Pantaloni et al., 1997).

The results also show that soybean plants grow well in a wide range of Ca:Mg ratios, as long as Al³⁺ is absent. This is supported by the fact that plant growth was maximized in limed soil (60 % base saturation) at a Ca:Mg ratio of 52 (Figure 6), whereas at a lower Ca:Mg ratio of 8 or 26 in soils of treatments 1 and 7 in which Al³⁺ was high, both shoot and root growth was reduced. It should be noted that even in the treatment with highest Ca:Mg ratio shoot growth was lower than in limed soil (Figure 6), indicating that alleviation of Al toxicity was not complete, and greater amounts of the cation may be required for full protection. Although the application of greater Ca and Mg amounts (60 % base saturation) led to higher levels of tissue Ca and Mg (compare Figure 7b,c with Figure 7e,f), the responses in shoot and root growth were not as positive as one would expect (compare Figure 6A to Figure 6B). Perhaps, growth was more limited as a result of a higher Al³⁺ concentration in the soil solution due to Ca and Mg displacement of exchangeable Al³⁺, which in turn led to a higher Al uptake (Figure 7). Additionally, the possibility of Cl⁻ toxicity can not be completely ruled out.

The chemical characteristics of the soil used here parallel those of acid subsoils, namely low exchangeable Ca²⁺ and Mg²⁺ and high Al³⁺. Hence, in acid subsoils as those in the Corredor region of




Figure 7. Shoot concentration of Al, Ca and Mg of soybean cv. UFVS-2001 grown in an acid soil with variable Ca:Mg ratios. A-C 10 % base saturation; D-F 60 % base saturation. Lime treatment received CaCO_3 and all other nutrients, except Mg. Control (Ctrl) treatment received no liming, but the supply of all other nutrients, except Ca and Mg, was ensured. Treatment means followed by the same letter are not significantly different by Tukey's test ($\alpha = 0.05$).

Brazil, soybean rooting depth and drought resistance could be maximized by supplying not only a mobile source of Ca^{2+} (Ritchey et al., 1980; Lopes, 1996), but probably also Mg^{2+} . Earlier studies have suggested that a similar soybean growth in acid soil could be obtained with lower lime rates if the amendment contained Mg^{2+} in addition to Ca^{2+} (Muchovej et al., 1986). Furthermore, it has been pointed out that responses to liming in high- Al^{3+} , acid soils seem to be substantial under conditions with low Mg^{2+} , even if Ca^{2+} is not limiting (Caires et al., 2003). Therefore, the importance of a Mg^{2+} source in areas where soybean is frequently used in crop rotations must not be overlooked. The observations that Ca^{2+} and Mg^{2+} from low solubility sources such as carbonate move downward in the soil profile under no till conditions

over time (Caires et al., 2000) further emphasizes the need to use dolomitic limestone, at least until Mg^{2+} levels in the (sub)soils are adequately raised.

ACKNOWLEDGMENTS

The authors are thankful to two anonymous reviewers and the assistant editor for their constructive questions and suggestions. Dr. I. R. Silva is grateful for the financial support from CNPq-Conselho Nacional de Desenvolvimento Científico e Tecnológico through grants 300543/00-0 and PROFIX grant No. 541090/01-1. Finally, Dr. I.R. Silva would like to thank Prof. J.C. Ker for helping to select, classify and collect soil samples in the field.

LITERATURE CITED

- ALVA, A.K.; ASHER, C.J. & EDWARDS, D.G. The role of calcium in alleviating aluminum toxicity. *Aust. J. Agric. Res.*, 37:375-382, 1986.
- CAIRES, E.F.; BLUM, J.; BARTH, G.; GARBUIO, F.J. & KUSMAN, M.T. Alterações químicas do solo e resposta da soja ao calcário e gesso aplicados na implantação do sistema plantio direto. *R. Bras. Ci. Solo*, 27:275-286, 2003.
- CAIRES, E.F.; FONSECA, A.F.; FELDHAUS, I.C. & BLUM, J. Crescimento radicular e nutrição da soja cultivada no sistema plantio direto em resposta ao calcário e gesso na superfície. *R. Bras. Ci. Solo*, 25:1029-1040, 2001.
- CAIRES, E.F.; BANZATO, D.A. & FONSECA, A.F. Calagem na superfície em sistema plantio direto. *R. Bras. Ci. Solo*, 24:161-169, 2000.
- CAIRES, E.F.; CHUEIRI, W.A.; MADRUGA, E.F. & FIGUEIREDO, A. Alterações de características químicas do solo e resposta da soja ao calcário e gesso aplicados na superfície em sistema de cultivo sem preparo do solo. *R. Bras. Ci. Solo*, 22:27-34, 1998.
- ESWARAN, H.; REICH, P. & BEINROTH, F. Global distribution of soils with acidity. In: MONIZ, A.C.; FURLANI, A.M.C.; SHAFFERT, R.E.; FAGERIA, N.K.; ROSELEM, C.A. & CANTARELLA, H., Eds. *Plant-soil interaction at low pH: Sustainable agriculture and forestry production*. Viçosa, MG, Brazilian Soil Science Society, 1997. p.159-164.
- FAGERIA, N.K. Efeito da calagem na produção de arroz, feijão, milho e soja em solo de cerrado. *Pesq. Agropec. Bras.*, 11:1419-1424, 2001.
- FERRUFINO, A.; SMYTH, T.J.; ISRAEL, D.W. & CARTER, T.E. Root elongation of soybean genotypes in response to acidity constraints in a subsurface solution compartment. *Crop Sci.*, 40:413-421, 2000.
- GRAUER, U.E. & HORST, W.J. Modeling cation amelioration of aluminum phytotoxicity. *Soil Sci. Soc. Am. J.*, 56:166-172, 1992.
- KINRAIDE, T.B. & PARKER, D.R. Cation amelioration of aluminum toxicity in wheat. *Plant Physiol.*, 83:546-551, 1987.
- KINRAIDE, T.B. Identity of the rhizotoxic aluminum species. *Plant Soil*, 34:167-178, 1991.
- KINRAIDE, T.B.; RYAN, P.R. & KOCHIAN, L.V. Interactive effects of Al^{3+} , H^+ , and other cations on root elongation considered in terms of cell-surface electrical potential. *Plant Physiol.*, 99:1461-1468, 1992.
- KINRAIDE, T.B. Use of a Gouy-Chapman-Stern model for membrane-surface electrical potential to interpret some features of mineral rhizotoxicity. *Plant Physiol.*, 106:1586-1592, 1994a.
- KINRAIDE, T.B.; RYAN, P.R. & KOCHIAN, L.V. Al^{3+} - Ca^{2+} interactions in aluminum rhizotoxicity-Evaluating the Ca^{2+} -displacement hypothesis. *Planta*, 192:104-109, 1994b.
- KINRAIDE, T.B. Three mechanisms for the calcium alleviation of aluminum toxicity. *Plant Physiol.*, 118:513-520, 1999.
- KINRAIDE, T.B.; PEDLER, J.F. & PARKER, D.R. Relative effectiveness of calcium and magnesium in the alleviation of rhizotoxicity in wheat induced by copper, zinc, aluminum, sodium, and low pH. *Plant Soil*, 259:201-208, 2004.
- LOPES, A.S. Soils under cerrado: A success story in soil management. *Better Crops Int.*, 10:9-15, 1996.
- LOPES, A.S. Solos sob "Cerrado": Características, propriedades e manejo. Piracicaba, Instituto do Potassa & Fosfato, Instituto Internacional da Potassa, 1983. 162p.
- MA, J.F.; DELHAIZE, E. & RYAN, P.R. Aluminum tolerance in plants and the complexing role of organic acids. *Trends Plant Sci.*, 6:273-278, 2001.
- MACDIARMID, C.W. & GARDNER, R.C. Al toxicity in yeast-a role for Mg? *Plant Physiol.*, 112:1101-1109, 1996.
- MARSCHNER, H. Mineral nutrition of higher plants. San Diego, Academic Press, 1995. 889p.
- MARTINS, O.C.; NOVAIS, R.F.; ALVAREZ V., V.H.; BARROS, N.F. & RIBEIRO, A.C. Respostas à aplicação de diferentes misturas de calcário e gesso em solos, II. Crescimento de Raízes, absorção de nutrientes e produtividade de soja. *R. Ceres*, 45:451-466, 1998.
- MENOSSO, O.G.; COSTA, J.A.; ANGHINONI, I. & BOHNEN, H. Crescimento radicular e produção de ácidos orgânicos em cultivares de soja com diferentes tolerâncias ao alumínio. *Pesq. Agropec. Bras.*, 11:1339-1345, 2001.
- MENOSSO, O.G.; COSTA, J.A.; ANGHINONI, I. & BOHNEN, H. Tolerância de genótipos de soja ao alumínio em solução. *Pesq. Agropec. Bras.*, 11:2157-2166, 2000.
- MOYER-HENRY, K.A.; SILVA, I.R.; JOHANNES, E.; ALLEN, N.S.; GOLDFARB, B.; JENNETTE, M.W. & RUFTY, T.W. An evaluation of aluminum tolerance in loblolly pine. In: *ANNUAL MEETING OF THE AMERICAN SOCIETY OF PLANT BIOLOGY*, 2003, Honolulu. *Plant Biology* 2003. Honolulu, 2003.
- MUCHOVEJ, R.M.C.; BORGES, A.C.; NOVAIS, R.F. & THIÉBAUT, J.T.L. Effect of liming level and Ca:Mg ratios on yield, nitrogen content and nodulation of soybeans grown in acid cerrado soil. *J. Soil Sci.*, 37:235-240, 1986.
- NICOLODI, M.; ANGHINONI, I. & GIANELLO, C. Indicadores da acidez do solo para recomendação de calagem no sistema plantio direto. *R. Bras. Ci. Solo*, 32:237-247, 2008.
- NOVAIS, R.F.; NEVES, J.C.L. & BARROS, N.F. Ensaio em ambiente controlado. In: *EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA - EMBRAPA. Métodos de pesquisa em fertilidade do solo*. Brasília, 1991. p.189-251.
- OLIVEIRA, E.L. & PAVAN, M.A. Control of soil acidity in no-tillage system for soybean production. *Soil Till. Res.*, 38:47-57, 1996.
- O'NEILL, M.A.; WARRENFELTZ, D.; KATES, K.; PELLERIN, P.; DOCO, T.; DARVILL, A.G. & ALBERSHEIM, P. Rhamnogalacturonan-II, a pectic polysaccharide in the walls of growing plant cell, forms a dimer that is covalently cross-linked by a borate ester. In vitro conditions for the formation and hydrolysis of the dimer. *J. Biol. Chem.*, 271:22222-22228, 1996.

- PANTALONE, V.R.; KENWORTHY, W.J.; SLAUGHTER, L.H. & JAMES, B.R. Chloride tolerance in soybean and perennial *Glycine* accessions. *Euphytica*, 97:235-239, 1997.
- PEDLER, J.F.; KINRAIDE, T.B. & PARKER, D.R. Zinc rhizotoxicity in wheat and radish is alleviated by micromolar levels of magnesium and potassium in solution culture. *Plant Soil.*, 259:191-199, 2004.
- RITCHEY, K.D; SOUSA, D.M.G.; LOBATO E. & CORREA O. Calcium leaching to increase rooting depth in a Brazilian Savannah Oxisol. *Agron. J.*, 72:40-44, 1980.
- RYAN, P.R.; DELHAIZE, E. & JONES, D.L. Function and mechanism of organic anion exudation from plant roots. *Ann. Rev. Plant Physiol. Plant Molec. Biol.*, 52:527-560, 2001.
- SANZONOWICZ, C.; SMYTH, T.J. & ISRAEL, D.W. Hydrogen and Al inhibition of soybean root extension from limed soil into acid subsurface solutions. *J. Plant Nutr.*, 21:387-403, 1998.
- SARTAIN, J.B. & KAMPRATH, E.J. Aluminum tolerance in soybean cultivars based on root elongation in solution culture compared with growth in acid soil. *Agron. J.*, 70:17-20, 1978.
- SARRUGE, J.R. & HAAG, H.P. Análises químicas em plantas. Piracicaba, Escola Superior de Agricultura Luiz de Queiroz, 1974. 56p.
- SILVA, I.R.; SMYTH, T.J.; ISRAEL, D.W.; CARTER, T.E.; RAPER T.E. & RUFTY, T.W. Differential aluminum tolerance in soybean: An evaluation of the role of organic acids. *Physiol. Plant.*, 112, 200-210, 2001a.
- SILVA, I.R.; SMYTH, T.J.; CARTER, T.E. & RUFTY, T.W. Altered aluminum sensitivity in soybean genotypes in the presence of magnesium. *Plant Soil*, 230:223-230, 2001b.
- SILVA, I.R.; SMYTH, T.J.; ISRAEL, D.W.; CARTER, T.E.; RAPER, T.E. & RUFTY, T.W. Magnesium ameliorates aluminum rhizotoxicity in soybean by increasing citric acid production and exudation by roots. *Plant Cell. Physiol.*, 42:546-554, 2001c.
- SILVA, I.R.; SMYTH, T.J.; MOXLEY, D.F.; ALLEN, N.S. & RUFTY, T.W. Aluminum accumulation at nuclei of cells in the root tip: Fluorescent detection using lumogallion and confocal laser scanning microscopy. *Plant Physiol.*, 123:543-552, 2000.
- SILVA, V.; MOTTA, A.C.V.; MELO, V.F. & LIMA, V.C. Variáveis de acidez em função da mineralogia da fração argila do solo. *R. Bras. Ci. Solo*, 32:551-559, 2008.
- SPEHAR, C.R. Aluminum tolerance of soya bean genotypes in short term experiments. *Euphytica*, 76:73-80, 1994.
- TAN, K.; KELTJENS, W.G. & FINDENEGG, G.R. Aluminum toxicity with sorghum genotypes in nutrient solutions and its amelioration by magnesium. *J. Plant Nutr. Soil Sci.*, 155:81-86, 1992.
- YANG, M.Z.; SIVAGURU, M.; HORST, W.J. & MATSUMOTO H. Aluminum tolerance is achieved by exudation of citric acid from soybean roots. *Physiol. Plant.*, 110:72-77, 2000.