



Revista Brasileira de Ciência do Solo

ISSN: 0100-0683

revista@sbcs.org.br

Sociedade Brasileira de Ciência do Solo  
Brasil

Costa do Nascimento, Carlos Antonio; Vitti, Godofredo Cesar; de Abreu Faria, Leticia; Cerqueira Luz,  
Pedro Henrique; Latanze Mendes, Fernanda

AMMONIA VOLATILIZATION FROM COATED UREA FORMS

Revista Brasileira de Ciência do Solo, vol. 37, núm. 4, julio-agosto, 2013, pp. 1057-1063

Sociedade Brasileira de Ciência do Solo

Viçosa, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=180228128022>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

## Comissão 3.2 - Corretivos e fertilizantes

### AMMONIA VOLATILIZATION FROM COATED UREA FORMS<sup>(1)</sup>

Carlos Antonio Costa do Nascimento<sup>(2)</sup>, Godofredo Cesar Vitti<sup>(3)</sup>, Letícia de Abreu Faria<sup>(4)</sup>, Pedro Henrique Cerqueira Luz<sup>(5)</sup> & Fernanda Latanze Mendes<sup>(6)</sup>

#### SUMMARY

Nitrogen fertilization is a major component of the cost of agricultural production, due to the high cost and low efficiency of fertilizers. In the case of urea, the low efficiency is mainly due to losses by volatilization, which are more pronounced in cultivation systems in which plant residues are left on the soil. The objective of this work was to compare the influence of urea coated with sulfur or boric acid and copper sulfate with conventional N fertilizers on N volatilization losses in sugar cane harvested after stubble burning. The sources urea, sulfur-coated urea, urea coated with boric acid and copper sulfate, as well as nitrate and ammonium sulfate, were tested at amounts containing N rates of 120 kg ha<sup>-1</sup> N. The integration of new technologies in urea fertilization can reduce N losses by volatilization. These losses were most reduced when using nitrate and ammonium sulfate. The application of a readily acidified substance (boric acid) to urea was more efficient in reducing volatilization losses and nutrient removal by sugar cane than that of a substance with gradual acidification (elemental sulfur).

**Index terms:** elemental sulfur, boric acid, copper sulfate, sugarcane, harvest without burning.

#### RESUMO: VOLATILIZAÇÃO DE AMÔNIA ORIUNDA DE UREIA EM RAZÃO DE SEUS RECOBRIMENTOS

*A adubação nitrogenada é um dos principais componentes na formação do custo de produção, por causa do elevado preço e da baixa eficiência dos fertilizantes. No caso da ureia, a baixa eficiência deve-se principalmente às perdas por volatilização, que se intensifica em*

---

<sup>(1)</sup> Paper from the Master's thesis in Soil Science and Plant Nutrition (ESALQ/USP) of the first author, funded by FAPESP. Received for publication on November 30, 2011 and approved on April 26, 2013.

<sup>(2)</sup> PhD candidate in Soil Science and Plant Nutrition Graduate Program - ESALQ-USP. Av. Pádua Dias, 11. Postal Box 9. CEP 13418-900 Piracicaba (SP). E-mail: cactagro@yahoo.com.br

<sup>(3)</sup> Professor at Soil Science Department, ESALQ-USP. E-mail: gcvitti@usp.br

<sup>(4)</sup> PhD candidate in Soil Science and Plant Nutrition Graduate Program - ESALQ-USP, FAPESP grantee. E-mail: evazoot@yahoo.com.br

<sup>(5)</sup> Professor at Animal Science Department FZEA-USP. Av. Duque de Caxias, 254. CEP 13635-900 Pirassununga (SP). E-mail: phcerluz@usp.br

<sup>(6)</sup> Post doctoral candidate in Soil Science and Plant Nutrition Graduate Program - ESALQ-USP. E-mail: fernanda\_latanze@yahoo.com.br

*sistemas de cultivo com preservação de restos culturais sobre o solo. Objetivou-se com este trabalho comparar a influência da ureia recoberta com enxofre, com ácido bórico e sulfato de cobre e com fertilizantes nitrogenados convencionais sobre as perdas de nitrogênio (N) por volatilização, em cana-de-açúcar colhida sem queima prévia da palhada. As fontes estudadas para fornecer dose de N igual a 120 kg ha<sup>-1</sup> foram: ureia, ureia revestida com enxofre, ureia revestida com ácido bórico e sulfato de cobre, nitrato e sulfato de amônio. A agregação de novas tecnologias à ureia resultou na redução das perdas de N por volatilização de amônia. Entretanto, as menores perdas de N por volatilização de amônia foram encontradas com uso de nitrato e sulfato de amônio. A adição de substância prontamente acidificante (ácido bórico) à ureia foi mais eficiente na diminuição das perdas por volatilização e na exportação de nutriente pela cana-de-açúcar, em relação à adição de substância com ação gradual de acidificação (enxofre elementar).*

*Termos de indexação: enxofre elementar, ácido bórico, sulfato de cobre, cana-de-açúcar, sem queima.*

## INTRODUCTION

Environmental laws encourage the adoption of harvest forms of sugarcane that dispense the previous burning of the crop stubble, which reflects back on the fertilization practices, not only in terms of amounts but also of sources and application forms. Commonly, sugar cane fertilizers are water-soluble and consequently readily available to the crop. Among the most typical nitrogen fertilizers are urea, sulfate and ammonium nitrate (Cantarella et al., 2007, Vitti et al., 2008).

Nitrogen (N) is a very dynamic element in the soil and as such can get lost by one or several processes, e.g., volatilization, leaching, denitrification, runoff, and crop removal (De Datta, 1981).

The losses of ammonia nitrogen (NH<sub>3</sub>-N) from urea applied to the surface are significant in soils with tropical and subtropical climate, due to the climatic conditions such as high temperatures, winds and high relative air humidity (Oliveira et al., 2007).

The straw left on the soil surface in the green cane system facilitates ammonia (NH<sub>3</sub>) losses by evaporation from amidic ammonia sources not incorporated into the soil (Trivelin et al., 1998). In a study by Cantarella et al. (1999), 12 and 30 %, respectively, of the NH<sub>3</sub>-N of the urea fertilizer applied on sugarcane straw, at 50 and 100 kg ha<sup>-1</sup> N, was lost by volatilization. Costa et al. (2003) and Vitti et al. (2007a) reported losses of more than 30 % of the N applied to straw cane.

The growth and quality of plants are mainly due to the amount of water and fertilizer. It is therefore very important to improve the use of water and fertilizer nutrients. However, about 40-70, 80-90 and 50-70 % of the applied amount of N, P and K fertilizer, respectively, is normally lost to the environment and cannot be absorbed by plants, causing great economic and resource losses, aside from environmental pollution (Lara Cabezas et al., 1999; Trivelin et al., 2002; Costa et al., 2003; Wu & Liu, 2008).

Fertilizers with higher efficiency represent a new trend to maintain fertilizer consumption and minimize environmental pollution (Du et al., 2008; Tang et al., 2009; Machado et al., 2011).

With the aim of increasing the use efficiency of urea, several studies tested various polymers with different principles of action, urease and nitrification inhibitors (Cantarella et al., 2008; Bhatia et al., 2010), rock dusts and organic compounds (Gioacchini et al., 2006; Kaoosi, 2007), micronutrients and some metals (Grohs et al., 2011), and sulfur (Girardi & Mourão Filho, 2004; Shaviv, 2005).

The purpose of this study was to compare the influence of urea coated with sulfur or with boric acid and copper sulfate, and of conventional nitrogen fertilizers on nitrogen losses by volatilization in sugarcane harvested without previous stubble burning.

## MATERIAL AND METHODS

The experiment was conducted in the sugarcane region of Piracicaba, São Paulo (South Latitude: 22° 40'; West Longitude: 47° 53') in a ratoon cane plantation harvested without burning, on sandy soil (chemical properties see table 1). The experimental design in randomized blocks consisted of six treatments: CT (control), UR (urea), AN (ammonium nitrate), UBC (urea coated with boric acid and copper sulfate), AS (ammonium sulfate), and SU (sulfur-coated urea), with four replications. All treatments were fertilized with N, K<sub>2</sub>O and P<sub>2</sub>O<sub>5</sub> corresponding to 120, 100 and 30 kg ha<sup>-1</sup>, respectively.

The NH<sub>3</sub>-N losses by volatilization were recorded (2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 22 days) after N fertilization. To estimate N losses by ammonia volatilization from soil after N fertilization, we used the direct method with semi-open collectors (Nömmik, 1973) in a dynamic mode, i.e., changing the chamber of the collectors to the subsequent bases (PVC rings by which collectors are fixed to the ground) and the

integration of correction factors proposed by Lara Cabezas et al. (1999). At each base, 15 g of N was applied in the fertilizer forms under study, in an amount proportional to the diameter of the collectors (15 cm).

Two pieces of foam (density 0.02 g cm<sup>-3</sup>) were fixed within each semi-open collector, soaked in 50 mL of phosphoric acid solution (0.25 mol L<sup>-1</sup>) and glycerin (3 % v/v) prior to installation, to absorb the ammonia. The upper foam was fixed (31 cm above the ground) to capture atmospheric ammonia, and to isolate the lower foam (15.5 cm above the ground), which has the function of capturing NH<sub>3</sub>-N from the soil surface (Lara Cabezas & Trivelin, 1990). For each sampling period, the foam soaked in phosphoric acid was replaced.

The upper foams were discarded and the ones beneath stored under refrigeration (5 °C) until analysis. Each foam was placed on a porcelain funnel equipped with a vacuum pump, and the foam was washed with approximately 400 mL of deionized water. From the solution extracted from each foam, a 50 mL aliquot was taken for NH<sub>3</sub>-N distillation and determination by the Kjeldahl method. At the time of fertilizer application and collector installation, the moisture content of the soil and straw were 6 and 20 %, respectively.

The volatilized ammonia nitrogen was calculated with the volumes of sulfuric acid used for titration in the blank tests and samples, according to equation (1).

$$N = (V_s - V_b) \times f \quad (1)$$

Where N = nitrogen captured in the collector, V<sub>s</sub> = volume of acid used for sample titration, V<sub>b</sub> = volume of acid spent for titration of the control; and f = 0.7 (factor of acid molarity, the volume of the aliquot and of the extract from the flask). Subsequently, the results were corrected according to the equations proposed by Lara Cabezas et al. (1999). The daily and accumulated volatilization rates were calculated as the sum of daily losses.

After the analysis of variance and confirmation of the significance of the F test, the means were partitioned by Tukey's test at a significance level of 5 %, using the statistical software SAS version 9.01 (SAS, 2004).

## RESULTS AND DISCUSSION

N losses by volatilization during the 22-day evaluation did not exceed 10 % of the applied N, regardless of the source (Figure 1). This was probably

a result of the high precipitation (> 50 mm) on the first days after fertilization (Figure 2). According to Lara Cabezas et al. (1997), when there is enough water to dilute the hydroxyl (OH<sup>-</sup>) concentration around the urea granules produced in the hydrolysis reaction, the urea is not only leached into the soil, but ammonia volatilization is also reduced. According to Cantarella et al. (1999), the incorporation of urea to a depth of ≥ 5 cm is sufficient to control the NH<sub>3</sub> loss.

Cantarella et al. (1999) observed that the rain maintained the level of volatilization low and concluded that loss rates were more pronounced when rainfall was insufficient to leach urea into the soil. Costa et al. (2003) also observed a reduction in the NH<sub>3</sub>-N loss rate after precipitation events.


These losses peaked between 15 and 16 days after fertilization, due to a lack of moisture to hydrolyze urea, because there was no rain until the 12<sup>th</sup> day. The process of urea hydrolysis started after rain fell on the 14<sup>th</sup>. According to Cantarella (2007), the losses can peak even on the first day if the soil is covered with plant residues. However, discrepant results of the duration of this period are reported in the literature: Gava (1999) observed the greatest losses on the 11<sup>th</sup> day while Costa et al. (2003), Pereira et al. (2009) and Tasca et al. (2011) mentioned a peak in the first six days. This variation is due to many factors that influence volatilization losses, among them moisture (Da Ros et al., 2005).

Nitrogen losses by volatilization were statistically higher when uncoated urea (UR) was used. Urea coating causes a statistically significant reduction of NH<sub>3</sub>-N losses in relation to UR, and losses from sulfur-coated urea (SU) were higher than from urea coated with boric acid and copper sulphate (UBC). However, regardless of the coating material, volatilization losses were not reduced to the level of losses from soils fertilized with nitrate (AN) and ammonium sulfate (AS) as nitrogen sources (Figure 1). The NH<sub>3</sub>-N losses from the different sources in the evaluation period can thus be ranked as follows: UR>US>UBC>AN=AS.


The higher losses from conventional urea (UR) in relation to the other sources are due to the action of urease, since the conditions for the action of the enzyme in the soil, microorganisms and plant residues were favorable. The influence of the enzyme on volatilization losses is greater when urea is applied on a soil surface covered with crop residues in an environment with insufficient moisture for fertilizer incorporation (Figure 1).

**Table 1. Soil chemical properties prior to the experiment**

| Depth | pH (CaCl <sub>2</sub> ) | MO                 | P                       | S | K <sup>+</sup> | Ca <sup>2+</sup> | Mg <sup>2+</sup>                   | Al <sup>+3</sup> | H+Al | CTC  | V  | m  |
|-------|-------------------------|--------------------|-------------------------|---|----------------|------------------|------------------------------------|------------------|------|------|----|----|
| cm    |                         | g dm <sup>-3</sup> | — mg dm <sup>-3</sup> — |   |                |                  | mmol <sub>c</sub> dm <sup>-3</sup> |                  |      |      | %  |    |
| 0-10  | 4.7                     | 13                 | 6                       | 9 | 1.0            | 10               | 5                                  | 3                | 31   | 47.4 | 34 | 16 |



**Figure 1. Concentration of accumulated NH<sub>3</sub>-N at each sampling after fertilization. UR: conventional urea, AS: ammonium sulfate, AN: ammonium nitrate, UBC: urea coated with boric acid and copper sulfate; SU: sulfur-coated urea. Means followed by the same letter do not differ statistically at 5 % probability by the Tukey test.**



**Figure 2. Daily precipitation and volatilization losses of NH<sub>3</sub>-N as a function of days after fertilization. UR: conventional urea, AS: ammonium sulfate, AN: ammonium nitrate, UBC: urea coated with boric acid and copper sulfate; SU: sulfur-coated urea.**

The straw left on the soil surface in the “green cane” harvest system facilitates NH<sub>3</sub> volatilization losses from amidic and ammoniacal fertilizers not incorporated into the soil (Trivelin et al., 1998) since apart from hampering the incorporation of urea into the soil, straw contains high amounts of urease. Cantarella et al. (1999) reported NH<sub>3</sub>-N volatilization losses of 12 and 30 %, respectively, from urea applied on sugarcane straw at 50 and 100 kg ha<sup>-1</sup> N, while Costa et al. (2003) and Vitti et al. (2007a) observed losses of more than 30 % of the N applied on cane residues.

With the use of sulfur-coated urea (SU), NH<sub>3</sub>-N losses were reduced by 50 % compared to conventional urea (UR), due to the physical protection of the layer of elemental sulfur in the SU.

Volatilization began to increase after a rainfall of only 2 mm (Figure 2), which was probably not enough rain to solubilize the elemental sulfur coating. In a study with urea treated with a urease inhibitor, Cantarella et al. (2008) observed the beginning of volatilization after a rainfall of 2.6 mm, in an experiment with sugarcane harvested without previous stubble burning.

The coating with elemental sulfur may also have acted as a barrier against the ureolytic action of residues on the ground. Vitti et al. (2007b) emphasized that the waste on the soil increases volatilization losses when urea is the N source, since the straw not only contributes to the amount of urease, but also forms a physical barrier preventing the soil-fertilizer contact, which reduces NH<sub>4</sub><sup>+</sup> adsorption to soil colloids. According to Silva et al. (1995), an increased contact between fertilizer and soil particles increases NH<sub>4</sub><sup>+</sup> adsorption to negatively charged soil, hampering the transformation into NH<sub>3</sub>, the form in which N is lost by volatilization.

Although the losses by UR and SU volatilization were statistically different, after the beginning of volatilization, a rainfall of over 50 mm reduced volatilization losses to almost zero. One can therefore not infer that coating with elemental sulfur would have the same effect under loss-prone conditions over a longer period (Figure 1). The dissolution of urea in soil solution occurs after the hydrolytic and microbial degradation of the protective sulfur coating, facilitated by micropores and imperfections, cracks and incomplete coating, and urea unprotected by sulfur is subject to the same losses (Shaviv, 2005; Cantarella, 2007; Trenkel, 2010).

In the plots with UBC as N source, NH<sub>3</sub>-N volatilization losses were on average 75 and 52 % lower than when using UR and SU, respectively. The reason is the reductive effect of boric acid on the pH around the urea granule, thus reducing the conversion of NH<sub>4</sub><sup>+</sup> to NH<sub>3</sub>. According to Cantarella (2007), the reaction of urea hydrolysis consumes H<sup>+</sup> protons and raises the pH around the particles, inducing a H<sup>+</sup> deficit that hampers the persistence of N in the form of NH<sub>4</sub><sup>+</sup>. However, the hydrolysis of UBC generates protons by the dissociation of H<sub>3</sub>BO<sub>3</sub>, thus meeting the H<sup>+</sup> demand for the maintenance of N in its reduced form.

The reduction of N volatilization losses by acidification of the medium has been studied for many years, mainly based on mixtures of urea with other acidifying N sources (Lara Cabezas et al., 1997; Costa et al., 2003; Lara Cabezas & Souza, 2008). However, because the ammonium sulfate is hygroscopic, mixing with urea tends to reduce the critical moisture and flowability of the mixture, and depending on the exposure time of the mixture in the field, can affect its application (Vitti et al., 2002). Furthermore, the unit price of N will increase with the mixture of



sources, since the concentration of N in ammonium sulfate is lower than in urea.

Thus, it is more feasible to use compounds such as boric acid as acidifying agents in the region around the granules, because this acid releases  $H^+$  when in contact with water. To provide the same amount of protons as by the  $H_3BO_3$  system, a three-fold amount of ammonium sulfate would be required. Sulfate is less efficient, due to the need of converting  $SO_4^{2-}$  to  $H_2SO_4$  for posterior donation of protons from the system.

Another explanation for the lower N volatilization from UBC is the toxicity caused by heavy metals such as Cu, Zn and Ag, affecting the urease activity in the soil. Chaperon & Sauvé (2007) demonstrated a reduction of urease activity in the soil due to the increase of metal concentration in the soil, including Cu.

Stuczynski et al. (2003) reported an inhibitory effect of Zn on the activity of a group of soil enzymes, especially of urease. Tabatabai (1977) reported an inhibitory effect of  $Cu^{+2}$  on urease activity in the soil, and that this effect was only lower than that of  $Ag^+$  and  $Hg^+$ . With the application of UBC, Grohs et al. (2011) obtained a reduction of approximately 50 % of ammonia volatilization compared to UR, in tilled soil.

Almost no  $NH_3$ -N volatilization losses were observed from plots fertilized with ammonium nitrate (AN) (Figure 2) and ammonium sulfate (AS) as N source, which can be explained by the chemical species in which the nutrient was applied to the crop. Similar results were obtained by Cantarella et al. (2003), Vitti et al. (2005), and Lara Cabezas & Souza (2008).

For the occurrence of volatilization losses, N must be in the form of  $NH_3$  which, under normal temperature and pressure conditions, is in gaseous form due to its low boiling point ( $-38^\circ C$ ). Ammonia is only formed when the pH of the medium is high, due to the removal of  $H^+$  protons from ammonium. This situation only occurs when fertilizers are applied to soil with a basic pH (a rare case in tropical soils) or when the dissolution of the particular nitrogen source favors the increase of  $OH^-$  around the granule (Cantarella, 2007).

When  $NH_4^+$  in the forms of AN and AS is applied to aerated soil, it tends to be rapidly oxidized to nitrate in soil with low available C, incorporated into microbial biomass, as in the case of soil where sugarcane is harvested without previous burning, or absorbed by plants. The  $NO_3^-$  is reduced, incorporated into the soil biomass or absorbed by plants (Cantarella, 2007). It is worth noting that in the transformation chain of nitrogen in tropical soil, both N sources resulted in the formation of only negligible amounts of ammonia, contrary to when urea was applied (Vitti et al., 2005; Lara Cabezas & Souza, 2008).

## CONCLUSIONS

1. In this study, the integration of new technologies in urea fertilizers resulted in the reduction of nitrogen losses by volatilization. These losses were lowest with the use of nitrate and ammonium sulfate.

2. The application of a readily acidified substance (boric acid) to urea was more efficient in reducing volatilization losses and nutrient removal by sugar cane than that of a substance with gradual acidification (elemental sulfur).

## ACKNOWLEDGEMENTS

The authors thank the São Paulo Research Foundation (FAPESP) for financial support in the development of the Project and the National Council of Scientific and Technological Development (CNPq) for the scholarship.

## LITERATURE CITED

- BHATIA, A.; SASMAL, S.; JAIN, N.; PATHAK, H.; KUMAR, R. & SINGH, A. Mitigating nitrous oxide emission from soil under conventional and no-tillage in wheat using nitrification inhibitors. *Agric. Ecosyst. Environ.*, 136:247-253, 2010.
- CANTARELLA, H.; MATTOS JUNIOR, D. de; QUAGGIO, J.A. & RIGOLIN, A.T. Fruit yield of Valencia sweet orange fertilized with different N sources and the loss of applied N. *Nutr. Cycl. Agroecosys.*, 67:215-223, 2003.
- CANTARELLA, H. Nitrogênio. In: NOVAIS, R.F.; ALVAREZ V., V.H.; BARROS, N.F.; FONTES, R.L.F.; CANTARUTTI, R.B. & NEVES, J.C.L., eds. *Fertilidade do solo*. Viçosa, MG, Sociedade Brasileira de Ciência do Solo, 2007. p.422-423.
- CANTARELLA, H.; ROSSETO, R.; BARBOSA, W.; PENNA, M.J. & RESENDE, L.C.L. Perdas de nitrogênio por volatilização da amônia e resposta da cana-de-açúcar à adubação nitrogenada, em sistema de colheita de cana sem queima prévia. In: CONGRESSO NACIONAL DA STAB, 7., Londrina, 1999. *Anais...* Londrina, STAB, 1999. p.82-87.
- CANTARELLA, H.; TRIVELIN, P.C.O.; CONTIN, T.L.M.; DIAS, F.L.F.; ROSSETTO, R.; MARCELINO, R.; COIMBRA, R.B. & QUAGGIO, J.A. Ammonia volatilization from urease inhibitor-treated urea applied to sugarcane trash blankets. *Sci. Agríc.*, 65:397-401, 2008.
- CHAPERON, S. & SAUVÉ, S. Toxicity interaction of metals (Ag, Cu, Hg, Zn) to urease and dehydrogenase activities in soils. *Soil Biol. Bioci.*, 39:2329-2338, 2007.

- COSTA, M.C.G.; VITTI, G.C. & CANTARELLA, H. Volatilização de N-NH<sub>3</sub> de fontes nitrogenadas em cana-de-açúcar colhida sem despalha a fogo. *R. Bras. Ci. Solo*, 27:631-637, 2003.
- DA ROS, C.O.; AITA, C. & GIACOMINI, S.J. Volatilização de amônia com aplicação de uréia na superfície do solo, no sistema plantio direto. *Cienc. Rural*, 35:799-805, 2005.
- DE DATTA, S.K. Principles and practices of rice production. New York, John Wiley, 1981. 618p.
- DU, C.; TANG, D.; ZHOU, J.; WANG, H. & SHAVIV, A. Prediction of nitrate release from polymer-coated fertilizers using an artificial neural network model. *Biosyst. Eng.*, 99:478-486, 2008.
- GAVA, G.J.C. Utilização do nitrogênio da uréia (15N) e da palhada (15N) por soqueira de cana-de-açúcar no manejo sem despalha a fogo. Piracicaba, Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo, 1999. 81p. (Dissertação de Mestrado)
- GIOACCHINI, P.; NICOLA, A.R.; MONTECCHIO, D.; MARZAORI, C. & CIAVATTA, C. Dynamics of mineral nitrogen in soils treated with slow-release fertilizers. *Commun. Soil Sci. Plant Anal.*, 37:1-12, 2006.
- GIRARDI, E.A. & MOURÃO FILHO, F.A.A. Crescimento inicial de laranjeira 'Valência' sobre dois porta-enxertos em função da adubação nitrogenada no plantio. *R. Bras. Frutic.*, 26:117-119, 2004.
- GROHS, M.; MARCHESAN, E.; SANTOS, D.E.; MASSONI, P.F.S.; SARTORI, G.M.S. & FERREIRA, R.B. Resposta do arroz irrigado ao uso de inibidor de urease em plantio direto e convencional. *Ci. Agrotec.*, 35:336-345, 2011.
- KAOOSI, M. Effects of zeolite application on rice yield, nitrogen recovery, and nitrogen use efficiency. *Commun. Soil Sci. Plant Anal.*, 38:69-76, 2007.
- LARA CABEZAS, W.A.R.; KORNDÖRFER, G.H. & MOTTA, S.A. Volatilização de N-NH<sub>3</sub> na cultura de milho: I. Efeito da irrigação e substituição parcial da ureia por sulfato de amônio. *R. Bras. Ci. Solo*, 21:481-487, 1997.
- LARA CABEZAS, W.A.R. & TRIVELIN, P.C.O. Eficiência de um coletor semi-aberto estático na quantificação de N-NH<sub>3</sub> volatilizado da ureia aplicada ao solo. *R. Bras. Ci. Solo*, 14:345-352, 1990.
- LARA CABEZAS, W.A.R. & SOUZA, M.A. Volatilização de amônia, lixiviação de nitrogênio e produtividade de milho em resposta à aplicação de misturas de ureia com sulfato de amônio ou com gesso agrícola. *R. Bras. Ci. Solo*, 32:2331-2342, 2008.
- LARA CABEZAS, W.A.R.; TRIVELIN, P.C.O.; BENDASSOLLI, J.A. & GASCHO, G.J. Calibration of a semi-open static collector for determination of ammonia volatilization from nitrogen fertilizers. *Commun. Soil Sci. Plant Anal.*, 30:389-406, 1999.
- MACHADO, D.L.M.; LUCENA, C.C.; SANTOS, D.; SIQUEIRA, D.L.; MATARAZZO, P.H.M. & STRUIVING, T.B. Slow-release and organic fertilizers on early growth of Rangpur lime. *R. Ceres*, 58:359-365, 2011.
- NÔMMIK, H. The effect of pellet size on the ammonia loss from urea applied to forest soil. *Plant Soil*, 39:309-318, 1973.
- OLIVEIRA, P.P.A.; TRIVELIN, P.C.O. & OLIVEIRA, W.S. Balanço de nitrogênio (<sup>15</sup>N) da ureia nos componentes de uma pastagem de capim-marandu sob recuperação em diferentes épocas de calagem. *R. Bras. Zootec.*, 36:1982-1989, 2007.
- PEREIRA, H.S.; LEÃO, A.F.; VERGINASSI, A. & CARNEIRO, M.A.C. Ammonia volatilization of urea in the out-of-season corn. *R. Bras. Ci. Solo*, 33:1685-1694, 2009.
- SAS Institute Inc. SAS OnlineDoc® 9.1.3. Cary, 2004.
- SHAVIV, A. Controlled release fertilizers. In: IFA INTERNATIONAL WORKSHOP ON ENHANCED-EFFICIENCY FERTILIZERS, 2005, Frankfurt. Proceedings... Paris, International Fertilizer Industry Association, 2005. p.1-15.
- SILVA, A.J.; LIMA JUNIOR, M.A.; FRAGA, V.S. & FERREIRA, N.M.C. Perdas de amônia por volatilização proveniente da uréia aplicada a solos dos trópicos úmidos. *R. Bras. Ci. Solo*, 19:141-144, 1995.
- STUCZYNSKI, T.I.; McCARTY, G.W. & SIEBIELEC, G. Response of soil microbiological activities to cadmium, lead, and zinc salt amendments. *J. Environ. Qual.*, 32:1346-1355, 2003.
- TABATABAI, M.A. Effects of trace elements on urease activity. *Soil Biol. Bioc.*, 9:9-13, 1977.
- TANG, C.Y.; KWON, Y.N. & LECKIE, J.O. Effect of membrane chemistry and coating layer on physiochemical properties of thin film composite polyamide RO and NF membranes: II. Membrane physiochemical properties and their dependence on polyamide and coating layers. *Desalination*, 242:168-182, 2009.
- TASCA, F.A.; ERNANI, P.R.; ROGERI, D.A.; GATIBONI, L.C. & CASSOL, P.C. Volatilização de amônia do solo após a aplicação de ureia convencional ou com inibidor de uréase. *R. Bras. Ci. Solo*, 35:493-502, 2011.
- TRENKEL, M.E. Slow and controlled-release and stabilized fertilizers: An option for enhancing nutrient use efficiency in agriculture. Paris, International Fertilizer Industry Association, 2010. 167p.
- TRIVELIN, P.C.O.; BENDASSOLLI, J.A.; OLIVEIRA, M.W. & MURAOKA, T. Potencialidade da mistura de aquamônia com vinhaça na fertilização de canaviais colhidos sem despalha a fogo. Parte II: Perdas por volatilização de amônia e recuperação do <sup>15</sup>N aplicado ao solo. *STAB*, 16:23-29, 1998.
- TRIVELIN, P.C.O.; OLIVEIRA, M.W.; VITTI, A.C.; GAVA, G.J.C. & BENDASSOLLI, J.A. Perdas do nitrogênio da ureia no sistema solo-planta em dois ciclos de cana-de-açúcar. *Pesq. Agropec. Bras.*, 37:193-201, 2002.
- VITTI, A.C.; TRIVELIN, P.C.O.; GAVA, G.J.C.; FRANCO, H.C.J.; BOLOGNA, I.R. & FARONI, C.E. Produtividade da cana-de-açúcar relacionada à localização de adubos nitrogenados aplicados sobre os resíduos culturais em canavial sem queima. *R. Bras. Ci. Solo*, 31:491-498, 2007b.

- VITTI, A.C.; TRIVELIN, P.C.O.; GAVA, G.J.C.; PENATTI, C.P.; BOLOGNA, I.R.; FARONI, C.E. & FRANCO, H.C.J. Produtividade da cana-de-açúcar relacionada ao nitrogênio residual da adubação e do sistema radicular. *Pesq. Agropec. Bras.*, 42:249-256, 2007a.
- VITTI, A.C.; CANTARELLA, H.; TRIVELIN, P.C.O. & ROSSETTO, R. Nutrição e adubação: Nitrogênio. In: DINARDO-MIRANDA, L.L.; VASCONCELOS, A.C.M. & LANDELL, M.G.A., orgs. *Cana-de-açúcar*. Campinas, IAC, 2008. p.239-269.
- VITTI, G.C.; TAVARES JÚNIOR, J.E. ; LUZ, P.H.C.; FAVARIN, J.L. & COSTA, M.C.G. Influência da mistura de sulfato de amônio com ureia sobre a volatilização de nitrogênio amoniacal. *R. Bras. Ci. Solo*, 26:663-671, 2002.
- VITTI, A.C.; TRIVELIN, P.C.O.; GAVA, G.J.C. & PENATTI, C.P. Produtividade da cana-de-açúcar relacionada à localização de adubos nitrogenados sobre a palha. *STAB*, 23:6-9, 2005.
- WU, L. & LIU, M. Preparation and properties of chitosan-coated NPK compound fertilizer with controlled-retention. *Carb. Poly.*, 72:240-247, 2008.