



Revista Brasileira de Ciência do Solo

ISSN: 0100-0683

revista@sbcs.org.br

Sociedade Brasileira de Ciência do Solo
Brasil

Hentz, Paulo; Corulli Corrêa, Juliano; Fontaneli, Renato Serena; Rebelatto, Agostinho; da
Silveira Nicoloso, Rodrigo; Neves Semmelmann, Claudio Eduardo
Poultry Litter and Pig slurry Applications in an Integrated Crop-Livestock System
Revista Brasileira de Ciência do Solo, vol. 40, 2016, pp. 1-12
Sociedade Brasileira de Ciência do Solo
Viçosa, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=180249980070>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Poultry Litter and Pig slurry Applications in an Integrated Crop-Livestock System

Paulo Hentz⁽¹⁾, Juliano Corulli Corrêa^{(2)*}, Renato Serena Fontaneli⁽³⁾, Agostinho Rebelatto⁽¹⁾, Rodrigo da Silveira Nicoloso⁽²⁾ and Claudio Eduardo Neves Semmelmann⁽¹⁾

⁽¹⁾ Instituto Federal Catarinense, Campus Concórdia, Concórdia, Santa Catarina, Brasil.

⁽²⁾ Empresa Brasileira de Pesquisa Agropecuária, Embrapa Suínos e Aves, Concórdia, Santa Catarina, Brasil.

⁽³⁾ Empresa Brasileira de Pesquisa Agropecuária, Embrapa Trigo, Passo Fundo, Rio Grande do Sul, Brasil.

ABSTRACT: Organic fertilizers derived from poultry litter and pig slurry are alternatives to mineral fertilizers in increasing soil nutrient availability. The aim of this study was to evaluate soil response, through characterization of organic C and available N, P, and K contents, and corn yield response to increasing amounts of poultry litter, pig slurry, and mineral fertilizers in an integrated crop-livestock production system (ICL) from 2011 to 2013. The experimental design consisted of randomized blocks in a 4 × 3 + 1 factorial arrangement with four replicates. The treatments consisted of four types of fertilizer, two organic (poultry litter and pig slurry) and two mineral, balanced with the same amounts of N, P and K as the organic fertilizers, one of which corresponded to the levels in the pig slurry (M1) and the other to the levels in the poultry litter (M2) in combination with three increasing application rates of N (100, 200, and 300 kg ha⁻¹ N) and control without fertilizer. For two years after implementing the ICL system, the application of the different rates of N using organic (pig slurry and poultry litter) or mineral (M1 and M2) fertilizers increased corn yields and K and P availability in the soil; these results were accompanied by small changes in organic C and total N content. There are similar efficiencies between the treatments pairs (pig slurry/M1 and poultry litter/M2).

Keywords: mineral fertilizers, organic carbon, nitrogen.

* **Corresponding author:**

E-mail: juliano.correa@embrapa.br

Received: September 1st, 2015

Approved: December 14, 2015

How to cite: Hentz P, Corrêa JC, Fontaneli RS, Rebelatto A, Nicoloso RS, Semmelmann CEN. Poultry Litter and Pig slurry Applications in an Integrated Crop-Livestock System. Rev Bras Cienc Solo. 2016;40:e0150072.

Copyright: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided that the original author and source are credited.



INTRODUCTION

The practice of fertilization, when necessary, can result in a yield up to three times greater than when not adopting the practice (IFDC, 2012) and is responsible for preserving 67 million hectares in Brazil (FAO, 2013). Such activity helps explain Brazil's position as the world's fourth largest consumer of mineral fertilizers; however, 75 % of these products are imported (ANDAs, 2013), which places under risk on agribusinesses, which account for 27 % of the Gross Domestic Product.

The development of sustainable production technologies aimed at improving nutrient availability in soil-plant systems following fertilizer applications is important in this scenario (Fancelli, 2010). Furthermore, research is needed on reuse of the nutrients available in organic waste (Grohskopf et al., 2015) and the use of highly productive conservation systems, such as integrated crop-livestock production systems (ICL), which place particular importance on soil fertility (Tormena et al., 2012).

To ensure that fertilization using organic fertilizers is sustainable and provides adequate nutrient availability to the soil-plant system, technical recommendation criteria are needed. Specific recommendations include the development of critical P limits (Gatiboni et al., 2015) to obtain the expected benefits from the chemical, physical, and biological properties of the soil and subsequent crop yield (Mafra et al., 2014; Grave et al., 2015). The concept has been adopted that organic fertilizers help promote system sustainability and should not be regarded as a potential source of environmental pollutants (Lourenzi et al., 2013).

Fertilization practices using organic fertilizers in ICL systems often result in economic and environmental gains because the diversity of the integrated system is enhanced such that new nutrient-cycling pathways are created and new ecosystem processes emerge (Anghinoni et al., 2011). Our knowledge of fertilization in ICL systems is based on mineral fertilizers and can be used to predict the increased export of nutrients generated by high-yield potential (Costa et al., 2015). However, organic fertilizers make different nutrients available to plants than mineral fertilizers (Scherer and Nesi, 2009; Scherer et al., 2010), and fertilizer recommendation studies should therefore be conducted on organics in ICL systems.

The hypothesis of this study is based on the premise that organic fertilizer is equal to or better than the mineral fertilizers in ICL systems when the same criteria for N, P, and K applications to the soil are adopted. Thus, the aim of this study was to evaluate soil response, based on the organic C content and N, P, and K availability in the soil, and corn yield response to different amounts of poultry litter, pig slurry, and mineral fertilizers in an ICL system.

MATERIALS AND METHODS

The experiment was conducted during the 2011-2013 growing seasons at the Instituto Federal Catarinense - IFC (Santa Catarina Federal Institute), Concordia *Campus*, in the municipality of Concordia, with geographical coordinates of latitude 27° 12' 0.08" and longitude 52° 4' 58.22" and an altitude of 569 m. The production system adopted was ICL, which included the cultivation of Syngenta Celeron LT hybrid corn (*Zea mays* L.) intercropped with brachiaria (*Urochloa brizantha* (Hochst. ex A. Rich) RD Webster), cultivar MG-5, during the summer and dual-purpose rye (*Secale cereale* L.), cultivar BRS Serrano, during the winter.

The climate is classified as humid subtropical (Cfa) according to the Köppen classification system, where the colder months (June and July) have mean temperatures of approximately 15 °C, and the mean annual temperature is 23 °C. The rains are regular and well distributed with a total annual rainfall greater than 1,500 mm, sufficient to ensure there are no water deficits. The relief is predominantly undulating with an 8 % slope.

Daily data for the maximum and minimum temperatures and rainfall throughout the two years of the experiment were collected at the weather station of Embrapa Suínos e Aves and are presented in figure 1. It should be noted that the rainfall and temperature conditions were adequate for crop development.

The soil in the experimental area was a *Nitossolo Vermelho Eutrófico* according to the Brazilian Soil Classification System (Santos et al., 2013). Soil chemical and physical properties are shown in table 1. The experimental area was previously managed with corn crops in the summer and black oats and turnips for vegetative cover in the winter from 1994 to 2011. During this period, there were two lime applications per year with 5 Mg ha^{-1} of dolomitic limestone, organic fertilization applications with pig slurry at $50 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$, based on the normative instructions of the state environmental agency of Santa Catarina (FATMA), and mineral fertilizer applications based on crop needs as defined by soil analysis; crop yields were also determined. Even with medium to high nutrient levels in the soil, increased production levels are expected from the amounts of organic and mineral fertilizers added to the soil to achieve the yield thresholds proposed by the study.

In setting up the experiment, cover crops were desiccated by applying glyphosate herbicide ($2,160 \text{ g ha}^{-1} \text{ a.i.}$), and then rye was planted. This agricultural practice was repeated 14 days prior to sowing the winter crops in 2011 and 2012, and prior to the 2011/12 and 2012/13 summer harvests.

The experiment was conducted under field conditions using a randomized block design with four replicates in a $4 \times 3 + 1$ factorial design, consisting of four fertilizers, three different amounts of N, and a control treatment without fertilization. The fertilizer treatments




Figure 1. Rainfall and maximum and minimum temperature recorded in the experiment during the 2011/2013 crop seasons in Concordia, SC, Brazil. Temperature and rainfall were reported as the average and sum of values registered during consecutive 10 days periods, respectively.

consisted of two organic fertilizers (poultry litter and pig slurry), two mineral fertilizers (M1 and M2) and a control (no fertilizer) in combination with 100, 200, and 300 kg ha⁻¹ N applied in each treatment. The experimental units were formed by 5 × 5 m plots (25 m²) with 2.5 m between blocks.

The fertilizer was applied on the surface next to the plant row for both the winter and summer crops. The pig slurry and poultry litter were obtained from production systems at the IFC, Concórdia Campus. The chemical properties of each organic fertilizer for each crop are shown in table 2 and were analyzed according to official methods (Rice et al., 2012) to determine the N, P, and K content.

Information on the N, P, and K concentrations in the organic fertilizer were used to establish the mineral fertilizer formulations from the following sources: urea for N, triple superphosphate for P, and potassium chloride for K, such that the M1 treatment contained the same levels of these nutrients as the pig slurry, and the M2 treatment, the same as the poultry litter (slurry/M1 and litter/M2).

Table 1. Soil chemical and physical properties prior to fertilizer application in 2011

Property	0.00-0.05 m	0.05-0.10 m	0.10-0.20 m	0.20-0.40 m
N (mg kg ⁻¹)	1,885	1,707	1,543	1,203
Cu (mg kg ⁻¹)	4.7	5.46	4.4	3.9
K (mg kg ⁻¹)	590	406	346	232
Mg (mg kg ⁻¹)	588	488	508	452
Ca (mg kg ⁻¹)	1,676	1,344	1,912	1,496
H+Al (mmol _c kg ⁻¹)	57	60	58	54
CTC (mmol _c kg ⁻¹)	205	178	205	172
V (%)	72	66	72	69
P (mg kg ⁻¹)	100	80	70	24
Zn (mg kg ⁻¹)	5.1	4.4	3.6	1
Organic C (g kg ⁻¹)	18	17	17	14
pH(H ₂ O)	5.8	5.6	5.5	5.3
Clay (g kg ⁻¹)	680	680	700	700

N, Cu, K, Mg, Ca, H+Al, P, Zn, Organic C, pH(H₂O): in according to Tedesco et al. (1995). Clay: in according to Claessen (1997).

Table 2. Nitrogen, phosphorus, and potassium contents in poultry litter (PL) and pig slurry (PS) and phosphorus and potassium inputs in cropping systems

	Content of nutrients in fertilizer residue			P input			K input		
	N	P	K	N application rate (kg ha ⁻¹)					
				100	200	300	100	200	300
Rye crop 2011									
	g kg ⁻¹ (PL) or g L ⁻¹ (PS)			kg ha ⁻¹					
Poultry litter	20.3	11.6	26.9	57	114	171	132	265	397
Pig slurry	4.20	0.42	2.43	10	20	30	58	116	174
Maize crop 2011/12									
Poultry litter	20.7	13.4	12.8	64	128	192	62	124	186
Pig slurry	3.72	1.43	1.54	39	78	117	41	82	123
Rye crop 2012									
Poultry litter	19.3	36.1	46.2	190	380	570	242	484	726
Pig slurry	2.9	0.85	1.14	29	58	87	38	76	114
Maize crop 2012/13									
Poultry litter	21.8	13.3	13.4	61	122	183	61	122	183
Pig slurry	3.85	0.99	4.23	26	52	78	109	218	327

Determinations carried according to Rice et al (2012).

Dual-purpose BRS Serrano rye was used for pasture formation during the winter at a sowing density of 250-350 viable seeds m^{-2} (40 to 60 kg ha^{-1}), with a 0.17 m spacing between the rows. Syngenta single hybrid Celeron TL corn was the summer crop, with a 0.80 m row spacing and eight seeds per meter, intercropped with brachiaria MG-5, which was manually seeded between rows at a density of 4 kg ha^{-1} .

Soil samples were taken from the 0.00-0.05, 0.05-0.10, 0.10-0.20, and 0.20-0.40 m layers at the end of each summer growing season. Three simple samples were taken at random using a Dutch auger, one from the row and two from between the rows, to form a composite sample. The composite samples were analyzed to determine the C, N, P, and K content, as described by Tedesco et al. (1995).

In both crop years, the corn crop in each plot was harvested manually from two 2-m-long rows that were 0.8 m apart, for a total of 3.2 m^2 . Manual husking, weighing, and drying; separation of the chaff and grain; and determination of the weight of the harvested grain for per-hectare grain-yield calculations, with subsequent corrections for 13 % humidity, were then performed.

The model used to analyze the variance inherent in a randomized block design considered the following factors: fertilizer, application rate, and blocks. After confirming the significance of the reference variable using the F test, means values of the treatments were compared using Student's t-test at a 5 % of probability. The linear and quadratic effects between the N levels were also tested in each fertilizer using the GLM procedure in SAS.

RESULTS AND DISCUSSION

Fertilization with organic and mineral fertilizers with increasing amounts of N in an integrated crop-livestock production system (ICL) system increased organic C content for the M2 at every soil layer, and at the application rate of 300 kg ha^{-1} N, the C content in the M2 was higher than in the poultry litter treatment at the 0.00-0.05, 0.05-0.10, and 0.10-0.20 m layers, higher than M1 at the 0.05-0.10 m layer, and higher than the pig slurry treatment at the 0.20-0.40 m layer (Table 3). A trend toward an increase in the organic C content was observed for the M1 and slurry treatments in the 0.20-0.40 m layer. Scherer et al. (2010), after 15 to 25 years of successive applications of pig slurry in a no-tillage system, observed that the organic C content in the soil did not change in *Neossolo Litólico*, *Cambissolo Háplico*, and *Latossolo Vermelho*.

Two years after implementation of the ICL system, higher organic C content found in soils at the 300 kg ha^{-1} levels of N in the M2 treatment was converted into organic C input, based on soil bulk density, reaching the order of 1.2 $\text{Mg ha}^{-1} \text{yr}^{-1}$ in the 0.00-0.05 and 0.05-0.10 m surface layers, 2.4 $\text{Mg ha}^{-1} \text{yr}^{-1}$ in the 0.10-0.20 m layer, and 6 $\text{Mg ha}^{-1} \text{yr}^{-1}$ in the 0.20-0.40 m layer. These results are consistent with reports by Lovato et al. (2004) and Leite et al. (2009), who observed a mean input of 4.1 $\text{Mg ha}^{-1} \text{yr}^{-1}$ of organic C in the surface layer and 10.4 $\text{Mg ha}^{-1} \text{yr}^{-1}$ in the 0.10-0.20 m layer in a no-tillage system.

Soil under ICL potentially serves as a sink of atmospheric C and favors accumulation of soil organic matter (SOM) (Nicoloso et al., 2008). Input of C in the ICL system is provided by high production of plant residues (plant residues on the soil surface and roots in the soil profile) in both crops and pastures compared to conventional grain-only production systems (Souza et al., 2010; Anghinoni et al., 2011; Loss et al., 2012). Importantly, soils with high clay content have high capacity to store C because they have a higher specific surface area and better aggregation (Six et al., 2000).

The application of organic and mineral fertilizers in the ICL systems did not significantly change the total N content in the soil, with a trend toward decrease observed up to levels of 111 kg ha^{-1} in the M2 treatment in the 0.05-0.10 m layer and a trend toward linear increase observed for the pig slurry treatment in the 0.20-0.40 m layer. No differences

were observed among treatments for the same application rate for each particular layer (Table 4). These results can be explained by a soil reservoir that contains from 1 to 10 Mg ha⁻¹ of organic N in the 0.00-0.20 m layer depending on the type of soil (Phelan, 2009). The initial N levels in the soil (Table 1) were such that the treatment where no N was applied was just as effective as the other treatments; that is, the N was derived from the SOM and not exclusively supplied by the fertilizer.

The increasing levels of N in the system from the organic and mineral fertilizers were directly related to available P content (Table 5) due to the input of 121, 242, and 363 kg ha⁻¹ P in the form of poultry litter and 49, 98, and 196 kg ha⁻¹ P in the form of pig slurry in 2011 and 251, 502, and 753 kg ha⁻¹ P in the form of poultry litter and 55, 110, and 165 kg ha⁻¹ P in the form of pig slurry in 2012, with the same order of magnitude between litter/M2 and manure/M1 (Table 2).

The M2 treatment showed a trend toward an increase in available P content in all layers, which was linear for the 0.00-0.05 m layer and quadratic for the other layers; in contrast, the M1 treatment showed a trend toward a linear increase only for the 0.20-0.40 m layer. Among the organic fertilizers, the poultry litter showed a trend toward a linear increase in P in the 0.00-0.05 and 0.05-0.10 m layers and a trend toward a quadratic increase in the 0.20-0.40 m layer, whereas the manure showed a trend toward a linear increase in the 0.05-0.10 m layer and a quadratic increase in the deeper layers (Table 5). This significant input of P into the system increased the degree of saturation of the soil solution and promoted precipitation of orthophosphates with Fe, Al, and Ca ions for the formation of amorphous minerals with varying degrees of solubility (Reddy et al., 2005).

Table 3. Soil organic carbon content (OC) according to organic and mineral fertilizer application in integrated crop-livestock systems in 2013

ICL	N application rate (kg ha ⁻¹)				Equation
	0	100	200	300	
OC (0.00-0.05 m)					
	g kg ⁻¹				
Poultry litter	19	20	20	20 b	$\bar{y} = 20.0$
Pig slurry	19	19	19	21 b	$\bar{y} = 20.0$
M1	19	21	21	21 b	$\bar{y} = 21.0$
M2	19	19	20	23 a	$\hat{y} = 20 - 0.014^* x + 0.00009^* x^2 \quad R^2 = 0.97$
OC (0.05-0.10 m)					
Poultry litter	18	20	19	18 b	$\bar{y} = 18.6$
Pig slurry	18	18	20	20 ab	$\bar{y} = 18.9$
M1	18	18	20	19 b	$\bar{y} = 18.5$
M2	18	19	19	21 a	$\hat{y} = 17.8 + 0.013^{**} x \quad R^2 = 0.78$
OC (0.10-0.20 m)					
Poultry litter	17	18	18	16 b	$\bar{y} = 16.9$
Pig slurry	17	18	18	19 a	$\bar{y} = 17.8$
M1	17	15	17	19 a	$\bar{y} = 17.2$
M2	17	16	19	20 a	$\hat{y} = 15.9 + 0.013^{**} x \quad R^2 = 0.79$
OC (0.20-0.40 m)					
Poultry litter	13	15	16	16 ab	$\bar{y} = 15.0$
Pig slurry	13	17	17	15 b	$\hat{y} = 13.3 + 0.044^{**} - 0.0001^{**} x^2 \quad R^2 = 0.99$
M1	13	14	15	18 a	$\hat{y} = 12.8 + 0.015^{**} x \quad R^2 = 0.92$
M2	13	14	14	18 a	$\hat{y} = 12.8 + 0.015^{**} x \quad R^2 = 0.75$

Means followed by different letters in columns at the same depth are different according to Student's t-test ($p \leq 0.05$). Significance level ** ($p \leq 0.01$) and * ($p \leq 0.05$). M1: mineral treatment contained the same levels of these nutrients as the pig slurry, and the M2 mineral treatment, the same as the poultry litter.

Table 4. Total nitrogen content (TN) in soil according to organic and mineral fertilizer application in integrated crop-livestock systems in 2013

ICL	N application rate (kg ha ⁻¹)				Equation
	0	100	200	300	
TN (0.00-0.05 m)					
	g kg ⁻¹				
Poultry litter	1.9	2.0	2.3 a	1.9	$\bar{y} = 2.0$
Pig slurry	1.9	1.9	1.8 b	1.9	$\bar{y} = 1.9$
M1	1.9	1.8	1.9 b	1.8	$\bar{y} = 1.8$
M2	1.9	1.7	1.8 b	2.1	$\bar{y} = 1.9$
TN (0.05-0.10 m)					
Poultry litter	1.7	1.8	1.7	1.7	$\bar{y} = 1.7$
Pig slurry	1.7	1.7	1.6	1.8	$\bar{y} = 1.7$
M1	1.7	1.7	1.7	1.8	$\bar{y} = 1.7$
M2	1.7	1.6	1.7	1.9	$\hat{y} = 1.6 - 0.002^* x + 0.000009^* x^2$ R ² = 0.99
TN (0.10-0.20 m)					
Poultry litter	1.6	1.7	1.4	1.5	$\bar{y} = 1.6$
Pig slurry	1.6	1.6	1.6	1.7	$\bar{y} = 1.6$
M1	1.6	1.6	1.6	1.6	$\bar{y} = 1.6$
M2	1.6	1.6	1.6	1.7	$\bar{y} = 1.6$
TN (0.20-0.40 m)					
Poultry litter	1.4	1.6	1.5	1.5	$\bar{y} = 1.5$
Pig slurry	1.4	1.5	1.5	1.8	$\hat{y} = 1.38 + 0.001^* x$ R ² = 0.71
M1	1.4	1.6	1.5	1.6	$\bar{y} = 1.5$
M2	1.4	1.5	1.5	1.7	$\bar{y} = 1.5$

Means followed by different letters in columns at the same depth are different according to Student's t-test (p<0.05). Significance level ** (p≤0.01) and * (p≤0.05). M1 mineral treatment contained the same levels of these nutrients as the pig slurry, and the M2 mineral treatment, the same as the poultry litter.

Table 5. Phosphorus content in soil according to organic and mineral fertilizer application in integrated crop-livestock systems in 2013

ICL	N application rate (kg ha ⁻¹)				Equation
	0	100	200	300	
P (0.00-0.05 m)					
	mg kg ⁻¹				
Poultry litter	64.3	101.0 a	124.0 a	140.0 a	$\hat{y} = 70.33 + 0.25^{**} x$ R ² = 0.97
Pig slurry	64.3	59.4 b	72.5 b	85.1 b	$\bar{y} = 70.3$
M1	64.3	52.3 b	60.8 b	63.4 b	$\bar{y} = 60.2$
M2	64.3	69.8 b	85.6 b	133.0 a	$\hat{y} = 54.99 + 0.22^{**} x$ R ² = 0.84
P (0.05-0.10 m)					
Poultry litter	61.3	76.9 a	108.0	111.0 a	$\hat{y} = 62.33 + 0.18^{**} x$ R ² = 0.92
Pig slurry	61.3	58.6 ab	89.4	79.4 b	$\hat{y} = 59.39 + 0.08^{*} x$ R ² = 0.56
M1	61.3	44.6 b	99.4	49.4 c	$\bar{y} = 63.7$
M2	61.3	45.0 b	85.0	112.0 a	$\hat{y} = 57.78 - 0.13^{*}x + 0.001^{*} x^2$ R ² = 0.91
P (0.10-0.20 m)					
Poultry litter	48.1	55.0	54.4 b	49.1 b	$\bar{y} = 51.6$
Pig slurry	48.1	61.9	72.5 a	50.9 b	$\hat{y} = 46.67 + 0.28^{**}x - 0.0009^{**} x^2$ R ² = 0.89
M1	48.1	41.3	51.1 b	56.0 b	$\bar{y} = 49.1$
M2	48.1	52.0	32.9 c	108.0 a	$\hat{y} = 54.01 - 0.38^{*}x + 0.002^{**} x^2$ R ² = 0.79
P (0.20-0.40 m)					
Poultry litter	31.6	42.7 b	53.0 a	44.6	$\hat{y} = 30.65 + 0.19^{**}x - 0.0005^{**} x^2$ R ² = 0.93
Pig slurry	31.6	41.0 b	53.6 a	43.8	$\hat{y} = 30.27 + 0.19^{**}x - 0.0005^{**} x^2$ R ² = 0.87
M1	31.6	38.7 b	39.1 b	43.1	$\hat{y} = 32.83 + 0.03^{**} x$ R ² = 0.89
M2	31.6	60.6 a	35.0 b	39.4	$\hat{y} = 35.78 + 0.18^{*}x - 0.00062^{*} x^2$ R ² = 0.30

Means followed by different letters in columns at the same depth are different according to Student's t-test (p<0.05). Significance level ** (p≤0.01) and * (p≤0.05). M1 mineral treatment contained the same levels of these nutrients as the pig slurry, and the M2 mineral treatment, the same as the poultry litter.

Management practices that include organic fertilizers have positive effects on nutrient availability, making nutrients available to plants and microorganisms in the first year after application (Schomberg et al., 2009), and these practices are especially effective at making available those nutrients that are preferentially displaced through the process of diffusion, such as with P (Dao, 2014). Successive applications of organic fertilizers at amounts that exceed crop demand can lead to P movement in the soil profile (Hesketh and Brookes, 2000). Cassol et al. (2012) applied 200 m³ ha⁻¹ of manure and observed an increase in the extractable P content in the deeper layers, up to 0.40 m, showing evidence of the transfer of this element into deeper layers in a *Latossolo Vermelho*.

Thus, the results reported herein suggest that organic P in the poultry litter and in the pig slurry, due to higher soil inputs, allowed for different P availability dynamics compared to the paired mineral fertilizers, thus contributing more effectively to P availability in the soil.

The application of increasing amounts of N from organic and mineral fertilizers in the ICL system increased K content in the soil (Table 6), explained by the input of 194, 389, and 583 kg ha⁻¹ K in the form of poultry litter and 99, 198, and 297 kg ha⁻¹ K in the form of pig slurry in 2011 and by the input of 303, 606, and 909 kg ha⁻¹ K from poultry litter and 147, 294, and 441 kg ha⁻¹ K from pig slurry in 2012 (Table 2). The same amounts of K were applied through mineral fertilizers (M1 and M2) in the form of potassium chloride.

The treatments that significantly contributed to K content in the soil showed a trend toward a linear increase from poultry litter and pig slurry and a trend toward a quadratic increase from M1 in the 0.00-0.05 m layer, a trend toward a quadratic increase from M2 in the 0.05-0.10 and 0.20-0.40 m layers, and a trend toward a quadratic increase from manure and a linear increase from M1 in the 0.10-0.20 m layer (Table 6). Among the fertilizers at the same application rate, increased availability of K in the M2 and poultry litter treatments was present because of the higher initial levels of this nutrient applied to the system compared to the M1 and manure treatments (Table 2).

Table 6. Potassium content in soil according to organic and mineral fertilizers application in integrated crop-livestock systems in 2013

ICL	N application rate (kg ha ⁻¹)				Equation
	0	100	200	300	
K ₂ O (0.00-0.05 m)					
	mmol _c kg ⁻¹				
Poultry litter	14	16 b	19 a	20 a	$\hat{y} = 14.3 + 0.02^{**} x$ R ² = 0.95
Pig slurry	14	10 c	10 c	9 c	$\hat{y} = 13.3 - 0.01^{**} x$ R ² = 0.80
M1	14	9 c	13 b	14 b	$\hat{y} = 13 - 0.04^{*} x + 0.00014^{*} x^2$ R ² = 0.54
M2	14	20 a	15 b	21 a	$\bar{y} = 17.5$
K ₂ O (0.05-0.10 m)					
Poultry litter	14	11 ab	20 a	14 b	$\bar{y} = 14.9$
Pig slurry	14	8 b	12 b	10 c	$\bar{y} = 11.2$
M1	14	10 b	14 b	13 b	$\bar{y} = 12.9$
M2	14	14 a	12 b	19 a	$\hat{y} = 15 - 0.05^{**} x + 0.0002^{**} x^2$ R ² = 0.83
K ₂ O (0.10-0.20 m)					
Poultry litter	11	13 ab	12 ab	10 b	$\bar{y} = 11.5$
Pig slurry	11	7 c	8 c	10 b	$\hat{y} = 10 - 0.04^{**} x + 0.0001^{**} x^2$ R ² = 0.96
M1	11	10 b	14 a	14 a	$\hat{y} = 10.30 + 0.01^{*} x$ R ² = 0.74
M2	11	14 a	10 bc	14 a	$\bar{y} = 12.4$
K ₂ O (0.20-0.40 m)					
Poultry litter	9	11 b	13 c	10 c	$\bar{y} = 10.8$
Pig slurry	9	8 c	11c	9 c	$\bar{y} = 9.1$
M1	9	7 c	17 b	20 a	$\bar{y} = 13.2$
M2	9	16 a	22 a	14 b	$\hat{y} = 8.7 + 0.12^{**} x - 0.0004^{**} x^2$ R ² = 0.89

Means followed by different letters in columns at the same depth are different according to Student's t-test (p<0.05). Significance level ** (p≤0.01) and * (p≤0.05). M1 mineral treatment contained the same levels of these nutrients as the pig slurry, and the M2 mineral treatment, the same as the poultry litter.

Increased K levels in the soil were observed by Scherer and Nesi (2009) when poultry litter or pig slurry was used, which added 117 and 55 kg ha⁻¹ yr⁻¹ of the nutrient, respectively. After 19 applications of manure, Lourenzi et al. (2013) observed an input of 303, 606, and 1212 kg ha⁻¹ of K, and showed that the addition of increasing amounts of this organic fertilizer results in increases in the levels of K available in the 0.00-0.10 m layer. After 20 years of applying manure to an *Argissolo Vermelho*, Scherer et al. (2010) found increased levels of available K in both the surface and deeper layers. Similar results were observed by Adeli et al. (2008) and Ceretta et al. (2010), who emphasized that care that should be taken in the application of manure due to the mobility of K through percolation and runoff.

In comparisons of the organic and mineral fertilizers in the pairs (litter/M2 and manure/M1), increased P availability in the organic treatments and, conversely, increased K availability in the mineral treatments were often observed (Tables 5 and 6).

The application of increasing amounts of N from organic and mineral fertilizers in the ICL system increased corn yield in the 2011/12 and 2012/13 harvests (Table 7). These results can primarily be explained by increased P and K availability in the soil (Tables 5 and 6).

During the 2011/12 harvest, corn yields in the ICL system showed a linear increase for poultry litter and pig slurry and a quadratic increase for M1 and M2, where mineral fertilizers were superior to organic fertilizers at 100 and 200 kg ha⁻¹ of N. As of 200 kg ha⁻¹ of N, the pig slurry fertilizer was superior to the poultry litter fertilizer, and at 300 kg ha⁻¹ of N, it was similar to the mineral fertilizers (Table 7).

A direct relationship between poultry litter and pig slurry fertilization and corn plant yields due to the availability of N and P in the system when using this agricultural practice were reported by Oliveira et al. (2009) and Novakowski et al. (2013). Hanish et al. (2009) also reported a positive effect of poultry litter fertilization on corn yield when they compared the use of poultry litter to topdressing with natural urea.

Table 7. Maize yield according to application rates of nitrogen with organic and mineral fertilizers in integrated crop-livestock systems in 2011/2012 and 2012/2013 harvests

ICL	N application rate (kg ha ⁻¹)				Equation
	0	100	200	300	
Crop 2011/2012					
	Mg ha ⁻¹				
Poultry litter	6,184	8,567 b	8,698 c	11,153 b	$\hat{y} = 6,394 + 15.0^{**} x \text{ } R^2 = 0.91$
Pig slurry	6,184	8,410 b	11,753 b	14,629 a	$\hat{y} = 5,942 + 28.7^{**} x \text{ } R^2 = 0.99$
M1	6,184	12,666 a	14,925 a	14,118 a	$\hat{y} = 6,242 + 80.7^{**} x - 0.182^{**} x^2 \text{ } R^2 = 0.99$
M2	6,184	11,516 a	14,219 a	15,621 a	$\hat{y} = 6,250 + 60.5^{**} x - 0.098^{*} x^2 \text{ } R^2 = 0.99$
Crop 2012/2013					
Poultry litter	3,327	6,485 ab	7,382	7,812	$\hat{y} = 3,416 + 34.8^{**} x - 0.07^{*} x^2 \text{ } R^2 = 0.99$
Pig slurry	3,327	6,904 a	7,702	8,826	$\hat{y} = 3,482 + 35.7^{**} x - 0.06^{*} x^2 \text{ } R^2 = 0.97$
M1	3,327	7,940 a	8,859	8,646	$\hat{y} = 3,455 + 53.1^{**} x - 0.120^{**} x^2 \text{ } R^2 = 0.98$
M2	3,327	5,378 b	8,250	9,065	$\hat{y} = 3,491 + 20.1^{**} x \text{ } R^2 = 0.96$
Total grain yield in the system					
Poultry litter	9,511	15,051 b	16,079 c	18,964 b	$\hat{y} = 9,830 + 49^{**} x - 0.07^{**} x^2 \text{ } R^2 = 0.92$
Pig slurry	9,511	15,313 b	19,455 b	23,454 a	$\hat{y} = 9,587 + 59^{**} x - 0.04^{**} x^2 \text{ } R^2 = 0.91$
M1	9,511	20,605 a	23,783 a	22,763 a	$\hat{y} = 9,697 + 134^{**} x - 0.30^{**} x^2 \text{ } R^2 = 0.93$
M2	9,511	16,893 b	22,469 ab	24,686 a	$\hat{y} = 9,434 + 90^{**} x - 0.13^{**} x^2 \text{ } R^2 = 0.96$

Means followed by different letters in columns at the same depth are different according to Student's t-test ($p < 0.05$). Significance level ^{**} ($p \leq 0.01$) and ^{*} ($p \leq 0.05$). M1 mineral treatment contained the same levels of these nutrients as the pig slurry, and the M2 mineral treatment, the same as the poultry litter.

In the second growing season (2012/13 harvest), the use of increasing amounts of N showed a quadratic increase in corn yield from poultry litter, pig slurry and M1, and a linear increase from M2, whereas, at the 100 kg ha⁻¹ rate of N application, the pig slurry and M1 treatments were superior to the M2 treatment (Table 7).

The nutrients supplied by pig slurry and the other favorable chemical, physical, and biological effects that manure has on the soil generally increase corn grain yields (Ceretta et al., 2005; Scherer, 2011). According to Sartor et al. (2012), corn grain production increases at a linear rate with the application of manure, with an increase of 59.9 kg ha⁻¹ for each m³ of manure applied.

When the positive corn yield results for both harvest years are combined and expressed as total production of the ICL system, the need for fertilizer recommendations to achieve high production levels is more evident (Table 7). Chantigny et al. (2007), in a study analyzing the application of 130 kg ha⁻¹ of N to corn crops in the form of ammonium nitrate and pig slurry over 4 years, achieved an average of 9.3 Mg ha⁻¹, with mean yields that were 22 and 4 % higher than the control and mineral fertilizer, respectively.

CONCLUSIONS

Two years after implementing the integrated crop-livestock production system, the application of different amounts of N using organic (pig slurry and poultry litter) or mineral (M1 and M2) fertilizers increased corn yields and soil K and P availability with few changes in organic C and total N content.

There are similar efficiencies between the treatments pairs (pig slurry/M1 and poultry litter/M2).

REFERENCES

- Adeli A, Bolster CH, Rowe DE, McLaughlin MR, Brink GE. Effect of long-term swine effluent application on selected soil properties. *Soil Sci.* 2008;173:223-35. doi:10.1097/ss.0b013e31816408ae
- Anghinoni I, Moraes A, Carvalho PCF, Souza ED, Conte O, Lang CR. Benefícios da integração lavoura-pecuária sobre a fertilidade do solo em sistema plantio direto. In: Fonseca AF, Caires EF, Barth G, editors. *Fertilidade do solo e nutrição de plantas no sistema plantio direto*. Ponta Grossa: 2011. p.272-309.
- Associação Nacional para Difusão de Adubos - ANDA. Estatísticas [internet]. São Paulo, SP [Acesso: 10 Set 2013]. Disponível em: <http://www.anda.org.br/index.php?mpg=03.01.00&ver=por/>.
- Cassol PC, Costa AC, Ciprandi O, Pandolfo CM, Ernani PR. Disponibilidade de macronutrientes e rendimento de milho em Latossolo fertilizado com dejetos suíno. *Rev Bras Cienc Solo.* 2012;36:1911-23. doi:10.1590/S0100-06832012000600025
- Ceretta CA, Basso CJ, Pavinato PS, Trentin EF, Girotto E. Produtividade de grãos de milho, produção de matéria seca e acúmulo de nitrogênio, fósforo e potássio na rotação aveia preta/milho/nabo forrageiro com aplicação de dejetos líquidos de suínos. *Cienc Rural.* 2005;35:1287-95. doi:10.1590/S0103-84782005000600010
- Ceretta CA, Girotto E, Lourenzi CR, Trentin G, Vieira RCB, Brunetto, G. Nutrient transfer by runoff under no tillage in a soil treated with successive applications of pig slurry. *Agric Ecosyst Environ.* 2010;139:689-99. doi:10.1016/j.agee.2010.10.016
- Chantigny MH, Angers DA, Rochette P, Belanger G, Masse D, Cote D. Gaseous nitrogen emissions and forage nitrogen uptake on soils fertilized with raw and treated pig slurry. *J Environ Qual.* 2007;36:1864-72. doi:10.2134/jeq2007.0083
- Claessen MEC, organizador. *Manual de métodos de análise de solo*. 2ª ed. Rio de Janeiro: Centro Nacional de Pesquisa de Solos; 1997.

- Costa NR, Andreotti M, Lopes KSM, Yokobatake KLA, Ferreira JP, Pariz CM, Bonini CSB, Longhini VZ. Atributos do solo e acúmulo de carbono na integração lavoura-pecuária em sistema plantio direto. *Rev Bras Cienc Solo*. 2015;39:852-63. doi:10.1590/01000683rbc20140269
- Dao TH. Landscape-scale geographic variations in microbial biomass and enzyme-labile phosphorus in manure-amended Hapludults. *Biol Fertil Soils*. 2014;50:155-67. doi:10.1007/s00374-013-0844-0
- Fancelli AL. Boas práticas para uso eficiente de fertilizantes na cultura do milho. Piracicaba: International Plant Nutrition Institute; 2010.
- Food and Agriculture Organization of the United Nations - FAO. Relatório Anual da Organização das Nações Unidas para a Alimentação e Agricultura. 2013. [Accessed on: 24 Nov 2015]. Available at: <http://sna.agr.br/roberto-rodrigues-sobre-o-relatorio-anual-da-fao-tecnologia-agricola-brasileira-aumentou-a-produtividade-por-area-e-preservou-67-milhoes-de-hectares/>.
- Gatiboni LC, Smyth TJ, Schmitt DE, Cassol PC, Oliveira CMB. Soil phosphorus thresholds in evaluating risk of environmental transfer to surface waters in Santa Catarina, Brazil. *Rev Bras Cienc Solo*. 2015;4:1225-33. doi:10.1590/01000683rbc20140461
- Grave RA, Nicoloso RS, Cassol PC, Aita C, Corrêa JC, Costa MD, Fritz DD. Short-term carbon dioxide emission under contrasting soil disturbance levels and organic amendments. *Soil Till Res*. 2015;146:184-92. doi:10.1016/j.still.2014.10.010
- Grohskopf MA, Cassol PC, Correa JC, Mafra MSH, Panisson J. Organic nitrogen in a typic hapludox fertilized with pig slurry. *Rev Bras Cienc Solo*. 2015;1:127-39. doi:10.1590/01000683rbc20150080
- Hanish AL, Fonseca JA, Almeida, E. Efeito do uso de diferentes estratégias de manejo agroecológico no desempenho produtivo da cultura do milho. *Rev Bras Agroecol*. 2009;2:1631-4.
- Hesketh N, Brookes PC. Development of indicator risk of phosphorus leaching. *J Environ Qual*. 2000;29:105-10. doi:10.2134/jeq2000.00472425002900010013x
- International Fertilizer Development Center - IFDC. 2012. [Accessed on: 24 Nov 2015]. Available at: <http://rootsforgrowth.com/food-nutrition-security>.
- Leite LFC, Cardoso MJ, Costa DB, Freitas RCA, Ribeiro VQ, Galvão SRS. Estoques de C e de N e produtividade do milho sob sistemas de preparo e adubação nitrogenada em um Latossolo Vermelho-Amarelo do cerrado piauiense. *Cienc Rural*. 2009;39:2460-6. doi:10.1590/S0103-84782009000900012
- Loss A, Pereira MG, Giacomini SG, Perin A, Anjos LHC. Carbon and nitrogen content and stock in no-tillage and crop-livestock integration systems in the Cerrado of Goiás State, Brazil. *J Agric Sci*. 2012;4:96-105. doi:10.5539/jas.v4n8p96
- Lourenzi CR, Ceretta CA, Silva LS, Girotto E, Lorensini F, Tiecher TL, Conti L, Trentin G, Brunetto, G. Nutrientes em camadas de solo submetido a sucessivas aplicações de dejetos líquidos de suínos e sob plantio direto. *Rev Bras Cienc Solo*. 2013;1:157-67. doi:10.1590/S0100-06832013000100016
- Lovato T, Mielniczuk J, Bayer C, Vezzani F. Adição de carbono e nitrogênio e sua relação com os estoques no solo e com o rendimento do milho em sistema de manejo. *Rev Bras Cienc Solo*. 2004;1:175-87. doi:10.1590/S0100-06832004000100017
- Mafra MSH, Cassol PC, Albuquerque JA, Correa JC, Grohskopf MA, Panisson J. Acúmulo de carbono em Latossolo adubado com dejetos líquidos de suínos e cultivado em plantio direto. *Pesq Agropec Bras*. 2014;8:630-37. doi:10.1590/S0100-204X2014000800007
- Nicoloso RS, Lovato T, Amado TJC, Bayer C, Lanza Nova ME. Balanço do carbono orgânico no solo sob integração lavoura-pecuária no Sul do Brasil. *Rev Bras Cienc Solo*. 2008;32:2425-33. doi:10.1590/S0100-06832008000600020
- Novakowski JH, Sandini IE, Falbo MK, Moraes A. Fertilization with broiler litter in the production of organic corn in integrated crop-livestock. *Semina: Cienc Agron*. 2013;4:1663-72. doi:10.5433/1679-0359.2013v34n4p1663
- Oliveira FA, Cavalcante LF, Silva IF, Pereira WE, Oliveira JC, Costa Filho JF. Crescimento do milho adubado com nitrogênio e fósforo em um Latossolo Amarelo. *Rev Bras Cienc Agrár*. 2009;3:238-44. doi:10.5039/agraria.v4i3a1

- Phelan PL. Ecology-based agriculture and the next Green Revolution. Is modern agriculture exempt from the laws of ecology? In: Bohlen P, House G, editors. Sustainable agroecosystem management. Boca Raton: CRC Press; 2009.
- Reddy KR, Wetzel RG, Kadlec RH. Biogeochemistry of phosphorus in wetlands. In: Sims JT, Sharpley AN, editors. Phosphorus: agriculture and the environment. Madison: American Society of Agronomy; 2005.
- Rice EW, Baird RB, Eaton AD, Clesceri LS. Standard methods for the examination of water and wastewater. 22nd.ed. Washington, DC: American Public Health Association; 2012.
- Santos HG, Jacomine PKT, Anjos LHC, Oliveira VA, Oliveira JB, Coelho MR, Lumbreiras JF, Cunha TJF. Sistema brasileiro de classificação de solos. 3ª ed. Rio de Janeiro: Embrapa Solos; 2013.
- Sartor LR, Assmann AL, Assmann TS, Bigolin PE, Miyazawa M, Carvalho PCF. Effect of swine residue rates on corn, common bean, soybean and wheat yield. Rev Bras Cienc Solo. 2012;2:661-9. doi:10.1590/S0100-06832012000200035
- Scherer EE. Efeito de fontes orgânicas e mineral de nitrogênio sobre produção de milho e propriedades químicas do solo sob sistema plantio direto. Rev Agropec Catarinense. 2011;24:71-6.
- Scherer EE, Nesi CN. Características químicas de um Latossolo sob diferentes sistemas de preparo e adubação orgânica. Bragantia. 2009;3:715-21. doi:10.1590/S0006-87052009000300019
- Scherer EE, Nesi CN, Massotti Z. Atributos químicos do solo influenciados por sucessivas aplicações de dejetos suínos em áreas agrícolas de Santa Catarina. Rev Bras Cienc Solo. 2010;5:1375-83. doi:10.1590/S0100-06832010000400034
- Schomberg HH, Wietholter S, Griffin TS, Reeves DW, Cabrera ML, Fisher DS, Endale DM, Novak JM, Balkcom KS, Raper RL, Kitchen NR, Locke MA, Potter KN, Schwartz RC, Truman CC, Tyler DD. Assessing indices for predicting potential nitrogen mineralization in soils under different management systems. Soil Sci Soc Am J. 2009;73:1575-86. doi:10.2136/sssaj2008.0303
- Six J, Elliott ET, Paustian K. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. Soil Biol Biochem. 2000;32:2099-103. doi:10.1016/S0038-0717(00)00179-6
- Souza ED, Costa SEVGA, Anghinoni I, Lima CVS, Carvalho PCF, Martins AP. Biomassa microbiana do solo em sistema de integração lavoura-pecuária em plantio direto, submetido a intensidades de pastejo. Rev Bras Cienc Solo. 2010;34:79-88. doi:10.1590/S0100-06832010000100008
- Tedesco MJ, Gianello C, Bissani CA, Bohnen, H, Volkweiss SJ. Análises de solo, plantas e outros materiais. Porto Alegre: Universidade Federal do Rio Grande do Sul; 1995.
- Tormena CA, Betioli Junior E, Petean LP, Alves SJ. Atributos físicos de um Latossolo Vermelho distroférrico em sistema de integração lavoura-pecuária. Rev Bras Cienc Solo. 2012;36:389-400. doi:10.1590/S0100-06832012000200008