



Agronomía Colombiana

ISSN: 0120-9965

agrocolfabog@gmail.com

Universidad Nacional de Colombia
Colombia

Casas-Leal, Nelson Enrique; Vallejo-Cabrera, Franco Alirio; Estrada-Salazar, Edgar Iván
Mechanisms of resistance to *Neoleucinodes elegantalis* (Guenée) in wild germplasm of the genus
Solanum
Agronomía Colombiana, vol. 31, núm. 2, mayo-agosto, 2013, pp. 153-160
Universidad Nacional de Colombia
Bogotá, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=180328569003>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's homepage in redalyc.org

Mechanisms of resistance to *Neoleucinodes elegantalis* (Guenée) in wild germplasm of the genus *Solanum*

Mecanismos de la resistencia a *Neoleucinodes elegantalis* (Guenée) en germoplasma silvestre del género *Solanum*

Nelson Enrique Casas-Leal¹, Franco Alirio Vallejo-Cabrera¹, and Edgar Iván Estrada-Salazar¹

ABSTRACT

Several phytosanitary problems affect tomato (*Solanum lycopersicum*) crops, one of the most important being the tomato fruit borer, which has caused losses of up to 70% in areas of the Valle del Cauca department (Colombia). To find resistance mechanisms to this pest, plants of three wild introductions of *Solanum* (PI134417, PI134418, LA1264) and the commercial cultivar Unapal-Maravilla were planted in a screenhouse. A completely randomized design was used with eight replicates. Five releases of *N. elegantalis* were carried out and the response was evaluated by the antixenosis or no preference test. Five racemes of each introduction were also suspended from the top of a field cage (1.8 x 1.5 x 1.5 m) to evaluate their response to six releases of the pest. For the two experiments, both an analysis of variance and analysis of means were performed. The average oviposition per plant was significantly higher in Unapal-Maravilla as compared with the wild introductions, and the average number of eggs per fruit was significantly lower in LA1264. The insects laid more eggs on the fruit surface (>70%). Glandular trichomes types I, IV and VI found in the wild introductions and their associated chemicals had an antibiotic and antixenotic effect on *N. elegantalis*.

Key words: tomatoes, *Solanum habrochaites*, defense mechanisms, trichomes.

RESUMEN

Diversos problemas fitosanitarios afectan el cultivo de tomate (*Solanum lycopersicum*), siendo el pasador del fruto uno de los más importantes, causando pérdidas de hasta el 70% en zonas del departamento del Valle del Cauca (Colombia). Para encontrar mecanismos de resistencia a esta plaga, se sembraron plantas de tres introducciones silvestres de *Solanum* (PI134417, PI134418, LA1264) y un cultivar comercial Unapal-Maravilla en una casa de malla circular, utilizando un diseño completamente al azar con ocho repeticiones. Se realizaron cinco liberaciones de *N. elegantalis* y la respuesta se evaluó mediante la prueba de antixenosis o de no preferencia. Adicionalmente, cinco racimos de cada introducción fueron suspendidos de la parte superior de una jaula (1,8 x 1,5 x 1,5 m) para evaluar su respuesta a seis liberaciones de la plaga. Ambos ensayos se sometieron a análisis de varianza y análisis de medias. Unapal-Maravilla presentó un promedio de ovoposición por planta significativamente mayor que las introducciones silvestres, y LA1264 presentó un número promedio de huevos por fruto significativamente menor. Los insectos depositaron más huevos en la superficie del fruto (>70%). Los tricomas glandulares tipos I, IV y VI presentes en introducciones silvestres y sus sustancias químicas asociadas ejercieron un efecto antixenótico y antibiótico en *N. elegantalis*.

Palabras clave: tomate, *Solanum habrochaites*, mecanismos de defensa, tricomas.

Introduction

The tomato (*Solanum lycopersicum* L.) is one of the most important vegetables in Colombia and the world, in terms of fresh consumption and industrial purposes. One of the main problems affecting this crop, however, is its high susceptibility to pests. In Colombia, four species of insect pests severely limit tomato production: fruit borer *Neoleucinodes elegantalis*, citrus gall midge *Prodiplipsis longifila*, whitefly *Bemisia tabaci* and tomato leaf miner *Tuta absoluta*. Crop losses of up to 70%, attributable to *N. elegantalis*, have been reported in the department of Valle del Cauca and, as this pest remains inside the fruit throughout its larval stage, it

is very difficult to control. Furthermore, all tomato cultivars used in Colombia are susceptible to this species. The problem is so serious that for production to be economically viable, producers must resort to extreme control practices such as the indiscriminate use of highly toxic, residual pesticides, which greatly increases production costs and poses risks not only for the environment but also for the health of those using these products and consumers.

The development of cultivars genetically resistant to this pest is a realistic, efficient and long-term management alternative. To date, however, few studies have been conducted in Colombia that can serve as a basis to develop tomato

Received for publication: 8 February, 2013. Accepted for publication: 5 June, 2013.

¹ Faculty of Agricultural Sciences, Universidad Nacional de Colombia. Palmira (Colombia). favallejoc@unal.edu.co

cultivars resistant to *N. elegantalis* and, as a result, the information available is insufficient to begin the selection of resistant tomato genotypes. This study therefore aimed to study the response of insect resistance mechanisms present in three wild introductions of the genus *Solanum*, for subsequent breeding into cultivated tomatoes.

Materials and methods

Location

The study was carried out at the Experimental Center of the Universidad Nacional de Colombia–Palmira campus (CEUNP), located in the municipality of Candelaria, department of Valle del Cauca ($03^{\circ}25'25.3''$ N, $76^{\circ}25'47.8''$ W) at an altitude of 972 m a.s.l. and with an average temperature of 26°C and average relative humidity of 76%.

Evaluation under screenhouse conditions

The wild introductions *Solanum habrochaites* var. *glabratum* PI134417 and PI134418 and introduction *S. habrochaites* var. *typicum* LA1264, all identified as highly resistant, were used, as well as the *Solanum lycopersicum* commercial cultivar Unapal-Maravilla, identified as very susceptible based on assessments made by Restrepo *et al.* (2003).

A completely randomized design was used with the four introductions, eight plants per introduction. Five racemes were evaluated per plant. The plants were grown in 25-L plastic buckets, inside a circular screenhouse, with a 9 m diameter and 2.5 m in height, covered with muslin cloth. Plants of each introduction were randomly distributed in a circle, equidistant from the center of the screenhouse. Every week, plants were removed from the screenhouse for 8 h to facilitate pollination of the wild introductions by insects. The racemes formed were covered with muslin-cloth bags. One two-stemmed plant was used per bucket, and no insecticides were applied after the onset of fructification to avoid altering insect behavior.

Five releases of adult *N. elegantalis* were carried out according to the procedure described by Casas and Estrada (2005) each consisting of three females and one male, in the center of the screenhouse, to allow free choice of host plants. These releases were carried out when the diameter was greater than 1 cm for half of the raceme fruits of all plants of the four introductions.

Five days after each release, the racemes were carefully cut to avoid the release of the eggs laid by the insects, stored in a Bellotti-type containment tray, and placed in the laboratory to carry out measurements aided by a microscope. The

variables evaluated were: number of oviposited fruits per plant, number of eggs per fruit, number of eggs per plant, number of eggs per raceme, location of laid eggs and presence of trichomes on the fruit.

The data corresponding to the variable oviposited fruit per plant were transformed using arcsine and those of variables number of eggs per fruit and number of eggs per plant, using. Descriptive analysis was performed as well as analysis of variance using the GLM procedure of SAS®. Means comparison was performed using the LSD test, with a significance level of $P \leq 0.05$.

Evaluation in field cage conditions

To confirm the results obtained in the screenhouse, further testing was conducted in a field cage measuring 1.5 x 1.5 x 1.8 m, covered with muslin cloth. Fruit racemes of uniform development and number of fruits of each of the four introductions were hung inside the cage (Fig. 1).

A completely randomized design was used with the same four introductions, six racemes per introduction. Six releases of adult *N. elegantalis* were carried out, each with 12 females and three males. Racemes were evaluated 5 d after each release following the same procedure as in the screenhouse. The racemes were classified by position on the plant as follows: level A (low racemes: 1, 2 and 3) and level B (middle or upper racemes: 4 and 5).

The following variables were measured: oviposition per raceme, average number of eggs per raceme, total number of eggs per raceme and location of laid eggs. The data corresponding to the variable oviposition per raceme were transformed using arcsine and that of the variables average number of eggs per raceme and total eggs per raceme, using.

Analysis of variance was performed using the GLM procedure of SAS®, and mean comparison was performed using the LSD test, with a significance level of $P \leq 0.05$.

Results and discussion

Evaluation under screenhouse conditions

Number of oviposited fruit per plant

Significant differences were detected between introductions regarding number of oviposited fruits per plant. Unapal-Maravilla presented the highest oviposition per plant as compared with PI134417, PI134418 and LA1264. Likewise, LA1264 recorded an average significantly lower than those of Unapal-Maravilla, PI134417 and PI134418. There were




FIGURE 1. Fruit racemes of four introductions of *Solanum* evaluated in field cage conditions: (A) Unapal-Maravilla, (B) PI134417, (C) PI134418 and (D) LA1264.

no significant differences between the introductions of *S. habrochaites* var. *glabratum* PI134417 and PI134418 (Tab. 1).

Based on the number of oviposited fruits per plant, the high susceptibility of cultivated Unapal-Maravilla tomatoes and the high resistance of wild introductions to *N. elegantalis* were confirmed. In the case of Unapal-Maravilla, 70.0% of the fruits were oviposited, compared with only 7.3% of the fruits of *S. habrochaites* var. *glabratum* PI134417 and 6.8% of those of PI134418. Introduction *S. habrochaites* var. *typicum* LA1264 presented the lowest oviposition value, with only 0.5% of its fruits presenting eggs. The extremely

low oviposition of *N. elegantalis* in wild introductions, as compared with cultivated tomatoes, suggests that they have an antixenotic effect on the tomato fruit borer.

Number of eggs per fruit

Significant differences were found between the introductions regarding the number of eggs per fruit. LA1264 presented a significantly lower value than Unapal-Maravilla, PI134417 and PI134418. There were no significant differences between Unapal-Maravilla, PI134417 and PI134418 (Tab. 1). It should be taken into account that the number of eggs per fruit is conditioned by the natural reproductive activity of the insect, which is impossible to control.

TABLE 1. Oviposited fruits per plant, number of eggs per fruit and number of eggs per plant of *Neoleucinodes elegantalis* in different introductions of *Solanum* sp. under screenhouse conditions.

Variable	Introduction	Mean	cv (%)	Pr >F
Oviposited fruit per plant (%)	Unapal-Maravilla	70.0 a		
	PI134417	7.3 b	13.86	<0.0001**
	PI134418	6.8 b		
	LA1264	0.5 c		
Number of eggs per fruit	Unapal-Maravilla	3.1 a		
	PI134417	2.9 a	21.33	<0.0065*
	PI134418	3.2 a		
	LA1264	2.4 b		
Number of eggs per plant	Unapal-Maravilla	153.0 a		
	PI134417	82.0 b	38.68	<0.0002**
	PI134418	70.0 b		
	LA1264	12.0 c		

Means with the same letter were not significantly different ($P \leq 0.05$), using the LSD test. cv = Coefficient of variation.




FIGURE 2. Location of eggs laid by the tomato fruit borer *Neoleucinodes elegantalis* in (A) Unapal-Maravilla, (B) PI134417, (C) PI134418 and (D) LA1264.

TABLE 2. Location of eggs laid by *Neoleucinodes elegantalis* on four *Solanum* introductions under screenhouse conditions.

Introduction	Location of eggs laid	
	Fruit (%)	Calyx and peduncle (%)
Unapal-Maravilla	75.8	24.2
PI134417	63.4	36.6
PI134418	71.4	28.6
LA1264	66.7	33.3

Number of eggs per plant

The number of eggs per plant varied significantly between introductions. Unapal-Maravilla recorded a value significantly higher than the averages of PI134417, PI134418 and LA1264. Similarly, LA1264 presented a value significantly lower than those of Unapal-Maravilla, PI134417 and PI134418. There were no significant differences between PI134417 and PI134418 (Tab. 1).

Kennedy (2003) reported that *S. habrochaites* var. *glabratum* and *S. habrochaites* var. *typicum* have high levels of resistance to a large number of arthropod species. The results of this study verify the resistance of wild *Solanum* sp. introductions to the tomato fruit borer *N. elegantalis*.

Location of laid eggs

The location of *N. elegantalis* eggs laid on the four *Solanum* introductions is shown in both Fig. 2 and Tab. 2, with 69.3% of the eggs being laid on the fruit and the remaining 30.7% on the calyx and peduncle.

These results show that *N. elegantalis* prefers to lay eggs on the fruit surface rather than other structures such as the calyx and peduncle, regardless of the type of tomato (cultivated or wild). In a few cases, eggs were found in small leaflets connected to the raceme, in raceme rachis and even in leaf folioles near the racemes.

Number of eggs per raceme. Fig. 3 shows the percentage of oviposition in the first five racemes, with susceptible





FIGURE 3. Oviposition of *Neoleucinodes elegantalis* per raceme in four *Solanum* introductions.


introduction Unapal-Maravilla presenting the highest number of eggs. The apparent increase in eggs laid on racemes 4 and 5 of wild introductions can be attributed to the low availability of racemes and fruits of the susceptible introduction.

A significant increase in oviposition per raceme was observed in wild introductions, especially in PI134417 and PI134418, increasing from 3.5% in the lower racemes to 12% in middle racemes. In the case of LA1264, the difference in raceme position was not significant, oviposition being below 1% in both.

When only a low number of fruits was available on the susceptible introduction Unapal-Maravilla, the insects were forced to oviposit on the fruits of wild introductions, although these were not the preferred host plants. This situation can be attributed to the instinct of survival and self-preservation of the species in conditions of confinement. Restrepo *et al.* (2007) reported the presence of several eggs on fruits of wild introductions under field conditions, thus converting them into host plants of this insect; the fruits, however, showed no damage.




A



B

FIGURE 4. *Neoleucinodes elegantalis* eggs attached to type 1 trichomes of wild *Solanum* introductions: (A) PI134417 and (B) PI134418.



A



B


FIGURE 5. Antibiotic reaction on neonate larvae of *Neoleucinodes elegantalis* found on type one trichomes of wild *Solanum* introduction fruits: (A) larvae that died near the chorions and (B) larvae that died trying to escape from the fruit surface.



A




B




C


FIGURE 6. Antibiotic and antixenotic reaction of type IV and VI trichomes on the tomato fruit borer *Neoleucinodes elegantalis*: (A) PI134417, (B) PI134418 and (C) LA1264.



A



B




C


FIGURE 7. Antibiotic reaction on neonate larvae of *Neoleucinodes elegantalis* found on and within the fruit of wild introductions of *Solanum*: (A) larvae that died trying to penetrate the fruit, (B) living larvae penetrating the fruit and (C) larvae that died inside the fruit.



A



B



C

FIGURE 8. Trichomes present in wild genotypes of the genus *Solanum*: (A) fruit of the introduction PI134418, (B) type 1 glandular trichomes and (C) type IV and VI glandular trichomes.

While there is evidence of the presence of different mechanisms that allow wild introductions to defend themselves against different pests, ranging from physical (different types of trichomes) to chemical barriers (toxic substances in the trichomes and release of volatile repellents), these did not prevent adult female *N. elegantalis* from ovipositing on wild introductions under controlled conditions, especially on *S. habrochaites* var. *glabratum* PI134417 and PI134418, although to a much lesser degree than on cultivated tomato Unapal-Maravilla.

Van Schie *et al.* (2007) stated that *S. lycopersicum* plants produce and liberate a variety of volatile organic compounds, mainly terpenes present in their trichomes. Despite this, the present study demonstrated that, under controlled conditions, the cultivated tomato Unapal-Maravilla was highly susceptible to *N. elegantalis* attack.

Presence of trichomes in fruit and oviposition

Most of the eggs laid on the wild introductions attached to type 1 trichomes of the fruit (Fig. 4). This behavior probably resulted from the attempt to avoid the fruit surface, where type IV and VI trichomes are found, largely responsible for the high levels of resistance in these introductions as mentioned by Simmons and Gurr (2005), Kennedy (2003), De Souza *et al.* (2002) and Aragão *et al.* (2000).

These results validate the studies conducted by Kennedy (2003) and Simmons *et al.* (2004), who put forth that both glandular and non-glandular trichomes are essential plant defense components of the genus *Solanum*.

The presence of eggs attached to type 1 trichomes of the fruit in wild *Solanum* introductions suggests an antixenotic effect on adult *N. elegantalis* because it alters the insect's reproductive behavior when compared with its behavior on the susceptible introduction Unapal-Maravilla, where oviposition occurs on the fruit surface. The larvae hatching from eggs laid on this type of trichomes, after descending and reaching the fruit surface where type IV and VI trichomes are found, died quickly and very near the chorions (Fig. 5A). Some larvae were also observed trying to escape from the fruit surface, again ascending the type I trichomes, where they died shortly afterwards (Fig. 5B). These reactions of neonate larvae of *N. elegantalis* suggest a possible antibiotic effect.

Regarding the antibiotic and antixenotic effect of type I, IV and VI glandular trichomes as well as non-glandular trichomes present in wild introductions, Aragão *et al.* (2000) asserted that non-glandular trichomes can act directly on the insects, affecting oviposition, feeding, locomotion and

behavior by density and size. Simmons and Gurr (2005) stated that type VI glandular trichomes of *S. habrochaites* var. *typicum* contain several sesquiterpenes, including zingiberene, which is predominant in this type of trichome.

Kennedy (2003) indicated that the leaves of *S. habrochaites* var. *glabratum* PI134417 were lethal to multiple pest species due to the presence of toxic substances such as methylketones (2-tridecanone and 2-undecanone) in type VI glandular trichomes. The ketones make up 90% of the content of the gland head of type VI glandular trichomes in PI134417.

Eggs laid directly on the fruit surface were observed in the wild introductions PI134417, PI134418 and LA1264, but in smaller amounts. Larvae hatched from these eggs died a few millimeters from the chorions (Fig. 6), indicating an alteration in insect biology related to a possible antibiotic effect on the first development stage of the borer.

Type IV and VI trichomes of the three wild introductions PI134417, PI134418 and LA1264 trapped the neonate larvae of *N. elegantalis*, limiting their movements and causing their death by action of the secondary metabolites they contain (Fig. 6). Simmons and Gurr (2005) indicated that when the contents of glandular trichomes are released and come into contact with the pest, they can cause irritation, trap the larvae and even kill them.

Peter and Shanower (1998) found that tomato trichomes produce large amounts of 2-tridecanone, 2-undecanone and a viscous mixtures of acyl sugars. They also release volatile chemicals such as (E)-betafarnesene and nepetalactone.

Aragão *et al.* (2000) stated that glandular trichomes secrete terpenes, alkaloids, phenolics and other substances that act as olfactory or taste repellents. Similarly, Simmons and Gurr (2005) indicated that the exudates of these trichomes can deter or repel pests, suggesting an important antixenotic effect in response to chemicals present in trichome exudates, such as the acyl sugars in type IV trichomes that have a deterrent effect on insects.

Several of the larvae that were able to initially overcome the barriers of type I, IV and VI glandular trichomes as well as the effect of their chemical components tried to penetrate the fruit, but died in the attempt (Fig. 7A). Living larvae were observed penetrating the fruits (Fig. 7B), but soon after penetration were found dead inside the fruit (Fig. 7C), slightly damaging the affected fruits, mainly the mesocarp tissue. These reactions on insect biology suggest a possible antibiotic effect on *N. elegantalis*.

TABLE 3. Oviposited fruits, average number of eggs per raceme and number of *Neoleucinodes elegantalis* eggs oviposited on 10 racemes of *Solanum* under cage conditions.

Variable	Introduction	Mean	cv (%)	Pr >F
Oviposited fruit (%)	Unapal-Maravilla	36.3 a		
	PI134417	6.9 b	22.6	<0.0024**
	PI134418	3.0 b		
	LA1264	4.1 b		
Average number of eggs per raceme	Unapal-Maravilla	8.3 a		
	PI134417	4.3 b	47.8	<0.0011**
	PI134418	5.8 b		
	LA1264	2.0 b		
Total number of eggs laid on 10 racemes	Unapal-Maravilla	100 a		
	PI134417	13 b	68.6	<0.0200*
	PI134418	2 b		
	LA1264	29 b		

*Means with the same letter were not significantly different ($P \leq 0.05$) using the LSD test.

Simmons and Gurr (2005) stated that antibiosis is probably conferred by the chemical components of trichome exudates. Two types of methylketones (2-tridecanone and 2-undecanone) are present in type VI trichomes of *S. habrochaites* var. *glabratum*, and are associated with many adverse effects on different Lepidopterans and Hemipterans. They also indicated that the trichomes associated with adverse effects on pests (mortality, survival and entrapment) are type IV and VI glandular trichomes present in the wild introductions used in this study.

The adverse reactions observed in insect reproductive behavior and biology demonstrate a possible antibiotic and antixenotic effect on *N. elegantalis*, mainly attributed to the presence of type IV and VI glandular trichomes in fruits of wild introductions PI134417, PI134418 and LA1264.

In the case of Unapal-Maravilla, the female insects laid most of the eggs on the fruit surface and, to a lesser extent, on the calyx. Eggss howed no alteration in their development and larvae easily penetrated the fruit and developed normally.

The fruits of the wild introductions PI134417, PI134418 and LA1264 showed type I, IV and VI glandular trichomes (Fig. 8), which are the main defense against different pests, including the tomato fruit borer *N. elegantalis*.

Cultivated tomatoes present type I, III, V, VI and VII glandular trichomes on the fruit, young stems, shoot sand flower buds (Luckwill, 1943). They also contain steroidial glycoalkaloids, such as α -tomatine, that act as toxic defense compounds. These are found throughout the plant, including fruits and roots. However, despite having a great diversity of trichomes, Unapal-Maravilla is highly susceptible to *N. elegantalis* attack. Some researchers, such as Williams et

TABLE 4. Location of *Neoleucinodes elegantalis* eggs laid on four introductions of *Solanum* under field cage conditions.

Introduction	Location of eggs laid	
	Fruit (%)	Calyx and peduncle (%)
Unapal-Maravilla	70.0	30.0
PI134417	61.5	38.5
PI134418	100.0	0.0
LA1264	72.4	27.6

al. (1980), have indicated that the glandular trichomes of cultivated tomatoes have concentrations of 2-tridecanone and 2-undecanone 72 times lower than those found in wild introductions of *S. habrochaites*.

The different reactions observed in neonate larvae of *N. elegantalis* suggest that the type I, IV and VI glandular trichomes present in the wild introductions exert a strong antibiotic effect on neonate larvae of this pest, causing 100% mortality because no larvae of advanced instars or living larvae were found after two days of observation.

Evaluation in field cage conditions

The analysis of variance for the variables percentage of oviposition, average number of eggs per raceme and number of eggs per raceme showed significant differences among the introductions. Values for Unapal-Maravilla were significantly higher than those for PI134417, PI134418 and LA1264. No significant differences were observed between the values of wild introductions PI134417, PI134418 and LA1264 (Tab.3).

Tab. 3 shows the susceptibility of the cultivated tomato Unapal-Maravilla and the resistance of the wild introductions to the tomato fruit borer *N. elegantalis*. In

Unapal-Maravilla, 36.3% of the fruits were oviposited as compared with only 6.9% in *S. habrochaites* var. *glabratum* PI134417 and 3.0% in PI134418. In the case of *S. habrochaites* var. *typicum* LA1264, 4.1% of its fruit demonstrated oviposition. The notable difference in percentage of oviposition in wild introductions as compared with cultivated tomatoes suggests an antixenotic effect of these introductions, following a trend similar to that observed under the screenhouse conditions.

As observed in the screenhouse trial (Tab. 1), the cultivated tomato Unapal-Maravilla differed significantly from the wild introductions regarding percentage of oviposited fruit as well as total number of eggs laid on 10 racemes, with 76% of the oviposition occurring on the fruit and the remaining 24% on the calyx and peduncle. These results are similar to those observed under the screenhouse conditions where oviposition on the fruit was 69.3% and 30.7% on the other plant structures (Tab. 4).

Just as in the screenhouse, the female insects preferred to oviposit on the fruit surface rather than on other structures such as the calyx and peduncle.

In introduction PI134418, all oviposited eggs were on the fruit, whereas oviposition on the fruit was 70.0% in Unapal-Maravilla, 61.5% in PI134417 and 72.4% in LA124.

No eggs of the tomato fruit borer *N. elegantalis* were laid on the shoots accompanying the racemes, probably because of the rapid deterioration of plant material promoted by tissue dehydration. Despite this fact, the fruits of the evaluated racemes had a good appearance when evaluations were done 5 days after cutting, with racemes looking similar to those of whole plants in the field.

The reactions on the reproductive behavior and biology of the neonate larvae of this pest observed in the field cage conditions were similar to those observed in the screenhouse trials, confirming the apparent antibiotic and antixenotic effects on *N. elegantalis*, mainly due to the presence of type IV and VI glandular trichomes in the fruits of the wild *Solanum* introductions PI134417, PI134418 and LA1264.

Literature cited

Aragão, C.A., D.B. França, F. Rodrigo, and B.R. Gandolfi. 2000. Tricomas foliares em tomateiro com teores contrastantes do aleloquímico 2-tridecanona. *Sci. Agric.* 57(4), 813-816.

Casas, N.E. and E.I. Estrada. 2005. Estudios preliminares sobre la utilización de ultrasonido en el control del pasador del fruto *Neoleucinodes elegantalis* Guenée en el cultivo del tomate *Lycopersicum esculentum* Mill. M.Sc. thesis. Faculty of Agricultural Sciences, Universidad Nacional de Colombia, Palmira, Colombia.

De Souza, C., A. Viana-Bailez, and J. Blackmer. 2002. Toxicity of tomato allelochemicals to eggs and neonate larvae of *Neoleucinodes elegantalis* (Guenée) Service (Lep: Crambidae). p. 295. In: Abstracts, Entomology International Congress Proceedings. Agricultural Research Service, Washington DC.

Kennedy, G.G. 2003. Tomato, pests, parasitoids, and predators: trophic interactions involving the genus *Lycopersicon*. *Annu. Rev. Entomol.* 48, 51-72.

Luckwill, L.C. 1943. The genus *Lycopersicon*: An historical, biological, and taxonomic survey of the wild and cultivated tomatoes. *Aberd. Univ. Stud.* 120, 1-44.

Peter, A.J. and T.G. Shanower. 1998. Plant glandular trichomes chemical factories with many potential uses. *Resonance* 3, 41-45.

Restrepo, E.F., F.A. Vallejo, and M. Lobo. 2003. Estudios básicos para iniciar la producción de cultivares de tomate *Solanum lycopersicum* L. con resistencia al pasador del fruto *Neoleucinodes elegantalis* (guenée). Ph.D. thesis. Faculty of Agricultural Sciences, Universidad Nacional de Colombia, Palmira, Colombia.

Restrepo, E., F.A. Vallejo, and M. Lobo. 2007. Producción de poblaciones segregantes resistentes al pasador del fruto a partir de cruzamientos entre tomate y accesiones silvestres de *Lycopersicum* spp. *Acta Agron.* 56(1), 1-6.

Simmons, A.T., G.M. Gurr, D. McGrath, P.M. Martin, and H.I. Nicol. 2004. Entrapment of *Helicoverpa armigera* (Hübner) (Lep: Noctuidae) on glandular trichomes of *Lycopersicon* species. *Aust. J. Entomol.* 43, 196-200.

Simmons, A.T. and G.M. Gurr. 2005. Trichomes of *Lycopersicon* species and their hybrids: Effects on pests and natural enemies. *Agric. For. Entomol.* 7(4), 265-276.

Van Schie, C.C., M.A. Haring, and R.C. Schuurink. 2007. Tomato linalool synthase is induced in trichomes by jasmonic acid. *Plant Mol. Biol.* 64, 251-263.

Williams, W.G., G.G. Kennedy, E.T. Yamamoto, J.D. Thacker, and J. Bordner. 1980. 2-tridecanone: a naturally occurring insecticide from the wild tomato *Lycopersicon hirsutum* f. *glabratum*. *Science* 207(4433), 888-889.