



Agronomía Colombiana

ISSN: 0120-9965

agrocolfabog@gmail.com

Universidad Nacional de Colombia
Colombia

Morillo C., Yacenia; Cruz Morillo C., Ana; Muñoz F., Jaime Eduardo; Ballesteros P., William; González, Alonso

Molecular characterization of 93 genotypes of cocoa (*Theobroma cacao* L.) with random amplified microsatellites RAMs

Agronomía Colombiana, vol. 32, núm. 3, 2014, pp. 315-325

Universidad Nacional de Colombia
Bogotá, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=180333786003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Molecular characterization of 93 genotypes of cocoa (*Theobroma cacao* L.) with random amplified microsatellites RAMs

Caracterización molecular con microsatélites amplificados al azar (RAMs) de 93 genotipos de cacao (*Theobroma cacao* L.)

Yacenia Morillo C.¹, Ana Cruz Morillo C.², Jaime Eduardo Muñoz F.³, William Ballesteros P.⁴, and Alonso González⁵

ABSTRACT

Random amplified microsatellite (RAMs) markers six were used to characterize 93 genotypes of cocoa in Tumaco (Colombia). Hundred twenty seven bands were generated. The number of polymorphic loci varied between 11 and 25 for the AG and TG primers, respectively. This study differentiated the 93 genotypes into six groups with a 0.53 similarity, 0.28 mean heterozygosity (He) for the population, and 0.12 ± 0.02 genetic differentiation coefficient or Fst. A significant level of genetic diversity was evident in the *T. cacao* genotypes. This resource would benefit selection programs of individual trees or plant breeding programs. The genotypes clustered in a large proportion in accordance with the collection zone. This characteristic was associated with collection zones and along the rivers in the municipality of Tumaco. The RAM technique proved to be a useful tool for the determination of genetic diversity in *Theobroma* species.

Key words: *Malvaceae*, molecular markers, heterozygosity, clonal multiplication, genetic diversity.

RESUMEN

Seis marcadores microsatélites (RAMs) fueron utilizados para caracterizar 93 genotipos de cacao de las zonas productoras del municipio de Tumaco (Colombia). Se generaron 127 bandas. El número de loci polimórfico varió entre 11 y 25 para los cebadores AG y TG, respectivamente. Este estudio diferenció los 93 genotipos en seis grupos con una similaridad de 0,53, una heterocigosidad promedio (He) para la población de 0,28, y un coeficiente de diferenciación genética o Fst de $0,12 \pm 0,02$. Se evidenció un nivel importante de diversidad genética en los genotipos de *T. cacao*. Este recurso beneficiaría los programas de selección de árboles individuales o un programa de fitomejoramiento. Los genotipos se agruparon en *clusters* en gran proporción de acuerdo a su lugar de colecta. Esta característica fue asociada con zonas localizadas en las veredas ubicadas a lo largo de los ríos del municipio de Tumaco. La técnica RAMs demostró ser una herramienta útil para la determinación de la diversidad genética en especies de *Theobroma*.

Palabras clave: *Malvaceae*, marcadores moleculares, heterocigosidad, multiplicación clonal, diversidad genética.

Introduction

The exact origin of *Theobroma cacao* L. remains shrouded in mystery and it is hard to define, due to the vastness of its geographical distribution and the bio-diverse landscape (Pound, 1945; Bartley, 2005; Motamayor *et al.*, 2008). It is thought to have originated in the lowland rainforests of northeastern South America and in the southeast of Mexico (Dias, 2001). However, although the origin and domestication are controversial and widely debated, the Upper Amazon Basin is the most accepted origin given its large morphological (Pound, 1945; Bartley, 2005) and allelic diversity (Zhang *et al.*, 2012). The borders between Brazil, Peru and the Southern of southern Colombia encompass

the highest genetic diversity of this tree species (Thomas *et al.*, 2012).

Climatic models indicate that, 21,000 years ago, at the peak of the last ice age, tropical zones provided a refuge for species, including *T. cacao*. It is assumed that, from these zones, humans carried *T. cacao* to Mexico, where the selection of grains started. A better understanding about the spatial and temporal distribution of the genetic diversity of *T. cacao* (Zhang *et al.*, 2006) and the adaptability of this species to environmental changes (Pautasso, 2009) is fundamental to the development of a set of protocols to face the challenges of markets and future climates.

Received for publication: 28 October, 2014. Accepted for publication: 27 November, 2014.

Doi: 10.15446/agron.colomb.v32n3.46879

¹ Faculty of Agricultural Sciences and Natural Resources, Universidad de los Llanos. Villavicencio (Colombia).

² Faculty of Agricultural Sciences, Universidad Pedagógica y Tecnológica de Colombia (UPTC). Tunja (Colombia).

³ Faculty of Agricultural Sciences, Universidad Nacional de Colombia. Palmira (Colombia).

⁴ Faculty of Agricultural Sciences, Universidad de Nariño. Pasto (Colombia).

⁵ Tibaitata Research Center, Corporación Colombiana de Investigación Agropecuaria (Corpoica). Bogotá (Colombia). agonzalezma@corpoica.org.co

Molecular studies have confirmed the broad genetic diversity of the species, placing the Nacional cocoa variety in a group separate from the other three (Crouzillat *et al.*, 2000). Recent studies suggest the existence of 10 genetic groups, providing a better understanding of the amplitude and structure of *T. cacao* genetic diversity in the current populations. Motamayor *et al.* (2008) found little genetic diversity in the accessions of Central America and broad diversity in the Upper Amazon. This finding coincides with theories on the central origin of *T. cacao* (Chessman, 1944). The confirmation of these theories would bring beneficial implications for the genetic improvement and conservation of *T. cacao*.

Theobroma cacao has been characterized with different molecular markers (Lanaud *et al.*, 1999). The genetic diversity distribution of the 936 *T. cacao* genotypes was analyzed with 96 microsatellites in Latin America (Thomas *et al.*, 2012). The results confirm the high diversity of *T. cacao* in the Upper Amazon. These areas encompass the south of Peru to the Ecuadorian Amazon and the borders between Colombia, Peru and Brazil. They suggested that *T. cacao* was already widely distributed in the western Amazon before glaciation started.

Recent studies have reported high genetic diversity on farms in Honduras and Nicaragua (Ji *et al.*, 2012). This research was based on 70 polymorphic markers of only one nucleotide (SNP) and analyzed 84 traditional varieties and 31 clones from international collections. The results indicated a high genetic diversity in the traditional varieties with an appealing potential for further studies on intra-population variation.

The genetic diversity of 77 accessions of the 'Criollo' cocoa, collected in the Mayas Mountains of Belize, was analyzed using microsatellite markers (Motilal *et al.*, 2010). They identified 11 distinctive genotypes, which positioned the 'Criollo' cocoa within the germplasm of Belize. On the Ecuadorian Coast, 332 *T. cacao* accessions were characterized with 60 microsatellite molecular markers (AKA Simple Sequence Repeat, SSRs) (Loor, 2007). The results showed different heterozygosity levels. The genotypes with more homozygosity were genetically closer to the native Ecuadorian cocoa, which is known as 'Nacional' cocoa.

Restriction fragment length polymorphism (RFLPs) and random amplified polymorphic DNA (RAPDs) molecular markers were used by Lerceteau *et al.* (1997) in INIAP, Ecuador. Their results indicated high variability in the

degree of heterozygosity, which included genotypes with low levels for this characteristic.

Marcano *et al.* (2009) used genetic mapping to analyze characteristics important for *T. cacao* cultivation and also detected a degree of variation or polymorphism in the phenotypically distinct samples, as in the formation of two genetic groups. The first included 'Trinitarian' a Trinitarian origin and the second 'Trinitarian' included Trinitarian and 'Forastero' clones.

Amplified fragment length polymorphism (AFLPs) molecular markers were used by Palacio (2007) to evaluate 13 clones in Colombia. His results showed 250 developed markers.

The molecular markers known as RAMs are efficient for measuring the genetic diversity in plants and animals and differences between families, between species and within a species, allowing for the selection of fixed regions within the DNA molecule for different studies (Muñoz *et al.*, 2008). The number of detectable polymorphisms is theoretically unlimited and allows for the analysis of the information that is both expressed and unexpressed (Mahuku *et al.*, 2002). This methodology is feasible for small laboratories at a low cost and with simple equipment, with no requirement for previous sequence knowledge, radioactive isotope use, or population studies (Hantula *et al.*, 1996).

Farmers from Tumaco have traditionally grown different cocoa genotypes without technological management. Despite the fact that these genotypes have an aromatic profile with floral notes, highly appreciated in the cocoa market; its yields are very low. For this reason, their persistence is at risk because of cultivars with high productivity (CCN clone for Ecuador) but with lower quality. To face these challenges, this study aimed to obtain the first approximation of the genetic diversity of *T. cacao* in the municipality of Tumaco, in southern Colombia, using random amplified microsatellites (RAMs).

Materials and methods

Collection of *T. cacao* genotypes

The collection of *T. cacao* genotypes was carried out in the municipality of Tumaco (Colombia) on farms located in San Luis Robles, Rio Rosario, Rio Mejicano, Rio Chagüi and Mascarey (Municipio de Tumaco, 2006) (Fig. 1). The gathering was based on the morphologically characterization study of the elite *T. cacao* trees by Ballesteros (2011). According to Vallejo and Peña (2006), the zone

TABLE 1. *Theobroma cacao* genotypes used for the molecular characterization with RAM molecular markers from Tumaco (Colombia).

Genotype	Location	Collection point	Latitude	Longitude	Genotype	Location	Collection point	Latitude	Longitude
1	Isla Grande	Rosario	1.67626	-78.6466	105	Iscuande	Rosario	1.688694	-78.658806
3	Isla Grande	Rosario	1.65943	-78.63496	109	Iscuande	Rosario	1.687472	-78.655833
4	Isla Grande	Rosario	1.65933	-78.63499	110	Isla Grande	Rosario	1.6665	-78.640722
5	Isla Grande	Rosario	1.65922	-78.63515	111	Quebrada Arriba	San Luis Robles	1.678389	-78.688972
6	Isla Grande	Rosario	1.663694	-78.640944	112	Quebrada Arriba	San Luis Robles	1.67825	-78.689194
7	Isla Grande	Rosario	1.66387	-78.64067	113	Quebrada Arriba	San Luis Robles	1.678194	-78.689111
8	Carretera Rosario	San Luis Robles	1.6652	-78.67163	114	Pacora	Chagüi	1.7805	-78.485306
9	Carretera Rosario	San Luis Robles	1.66534	-78.67111	2CN51	Piñal Dulce	San Luis Robles	1.675503	-78.71379
10	Carretera Rosario	San Luis Robles	1.66544	-78.67094	CCN 51	Corpoica	Tumaco	1.8166	-78.7666
13	Monte Bonito	San Luis Robles	1.67275	-78.70705	CR01	La Lomita	San Luis Robles	1.696267	-78.6813
14	Piñal Dulce	San Luis Robles	1.6681	-78.7163	CR02	La Lomita	San Luis Robles	1.696233	-78.68125
21	La Sirena	Chagüi	1.732639	-78.485583	CR04	Nerete	San Luis Robles	1.707867	-78.6672
22	La Sirena	Chagüi	1.732583	-78.4855	CR05	Nerete	San Luis Robles	1.707867	-78.667217
23	La Sirena	Chagüi	1.73625	-78.477583	CR06	Nerete	San Luis Robles	1.707867	-78.667217
24	La Sirena	Chagüi	1.7365	-78.477444	CR08	Nerete	San Luis Robles	1.709033	-78.666733
30	Pacora	Chagüi	1.780778	-78.484861	CR09	Nerete	San Luis Robles	1.709067	-78.66675
31	Pacora	Chagüi	1.782833	-78.481778	CR10	Nerete	San Luis Robles	1.709167	-78.666833
32	Pacora	Chagüi	1.782111	-78.489111	CR11	Nerete	San Luis Robles	1.708478	-78.666992
33	Pacora	Chagüi	1.780944	-78.48925	CR12	Nerete	San Luis Robles	1.709433	-78.666917
35	Pacora	Chagüi	1.782139	-78.468328	CR13	Nerete	San Luis Robles	1.709433	-78.666917
42	La Ceiba	Chagüi	1.717833	-78.470528	CR14	Zona Carretera	San Luis Robles	1.689089	-78.673292
47	Bella Vista	Mejicano	1.683917	-78.540917	CR15	Piñal Dulce	San Luis Robles	1.698217	-78.719167
54	Guayabo	Mejicano	1.686806	-78.573389	CR16	Piñal Dulce	San Luis Robles	1.69825	-78.719233
60	Isla Grande	Rosario	1.66871	-78.66996	CR17	Zona Carretera	San Luis Robles	1.68905	-78.673517
61	Isla Grande	Rosario	1.6664	-78.64068	CR18	Zona Carretera	San Luis Robles	1.688883	-78.673283
62	Isla Grande	Rosario	1.67286	-78.64368	CR19	Zona Carretera	San Luis Robles	1.688867	-78.673267
63	Isla Grande	Rosario	1.67286	-78.64368	CR20	Zona Carretera	San Luis Robles	1.688867	-78.6732
69	Monte Bonito	San Luis Robles	1.67817	-78.68923	CR21	Zona Carretera	San Luis Robles	1.689467	-78.673283
70	Monte Bonito	San Luis Robles	1.67287	-78.70601	CR22	Zona Carretera	San Luis Robles	1.6894	-78.6732
71	Monte Bonito	San Luis Robles	1.67256	-78.70628	CR23	Isla Grande	Rosario	1.6946	-78.643333
73	Quebrada Arriba	San Luis Robles	1.67816	-78.68919	CR24	Isla Grande	Rosario	1.6969	-78.64545
74	Nerete	San Luis Robles	1.687556	-78.667944	CR25	La Ceiba	Chagüi	1.739267	-78.470533
75	Nerete	San Luis Robles	1.687389	-78.667528	CR26	La Ceiba	Chagüi	1.739917	-78.470283
76	Nerete	San Luis Robles	1.686917	-78.666917	CR27	La Ceiba	Chagüi	1.636535	-78.407402
77	Nerete	San Luis Robles	1.686639	-78.666333	CR28	La Ceiba	Chagüi	1.758617	-78.47745
78	Nerete	San Luis Robles	1.686167	-78.665667	CR29	Palambi	Chagüi	1.780933	-78.47955
82	Quebrada Arriba	San Luis Robles	1.670472	-78.69625	CR3	Quebrada Arriba	San Luis Robles	1.692633	-78.696033
85	Isla Grande	Rosario	1.6593	-78.63534	CR30	Pacora	Chagüi	1.8025	-78.488767
89	Isla Grande	Rosario	1.666	-78.639917	CR31	Guayabo	Mejicano	1.41125	-78.34242
91	La Lomita	San Luis Robles	1.675611	-78.681972	CR32	Bella Vista	Mejicano	1.41021	-78.32273
92	La Lomita	San Luis Robles	1.674972	-78.681806	ICS 60	Corpoica	Tumaco	1.817	-78.767
93	La Lomita	San Luis Robles	1.674833	-78.681528	ICS 95	Corpoica	Tumaco	1.818	-78.768
95	Carretera Rosario	San Luis Robles	1.658417	-78.72975	IMC 67	Corpoica	Tumaco	1.819	-78.769
96	Piñal Dulce	San Luis Robles	1.67797	-78.7137	Regional	Monte Bonito	San Luis Robles	1.672727	-78.706447
97	Piñal Dulce	San Luis Robles	1.67797	-78.71387	TSH561	Quebrada Arriba	San Luis Robles	1.514877	-78.470276
98	Piñal Dulce	San Luis Robles	1.67797	-78.71129	TSH561	Corpoica	Tumaco	1.82	-78.77
99	Mascarey	San Luis Robles	1.655444	-78.720444					

Molecular characterization

The protocol of Dellaporta *et al.* (1983) was used for the extraction of the DNA. A 0.8% agarose gel was used to visualize the DNA, dyed with ethidium bromide in a Maxi-cell Primo EC-340 Electrophoresis Gel System chamber (Thermo EC, Holbrook, AZ) and visualized with UV light. The DNA concentration of each sample was determined for comparison with known DNA concentrations of Lambda bacteriophage. The quantified DNA was diluted in HPLC water at a volume of 100 μL to 10 $\text{ng } \mu\text{L}^{-1}$ and stored at -20°C .

For the RAM analysis, six polymorphic primers were used (Technologies Inc., Alameda, CA) (Espinosa *et al.*, 2004; Mahuku *et al.*, 2002; Álvarez *et al.*, 2003) (Tab. 2).

TABLE 2. RAM primers used to determine the genetic diversity of *T. cacao*.

Primer (repeat unit)	Sequence (5'a 3')
CCA	DDB(CCA) ₅
CGA	DHB(CGA) ₅
AG	HBH(AG) ₇ A
CT	DYD(CT) ₇ C
TG	HVH(TG) ₇ T
CA	DBDA(CA) ₇

The following designations were used for degenerated sites: H (A/T/C); B (G/T/C); V (G/A/C) and D (G/A/T).

The amplification reaction was prepared with a final volume of 25 μL . The reaction mixture included buffer 1X, MgCl_2 -1.5 mM, dNTPs-0.2 mM, Taq Polymerase-1U, primer-2 μM and genomic DNA-10 ng.

The amplification was carried out in a PTC 100 Programmable Thermal Controller (MJ Research, Inc., Waltham, MA). The initial denaturation was at 95°C for 5 min, with denaturation at 95°C for 30 s, hybridization at 50°C (AG and CA primers), 55°C (CCA-TG-CT primers) and 58°C (GT-CGA primers) for 45 s, an extension at 72°C for 2 min, 37 cycles from the denaturation to the extension and, finally, an extension at 72°C for 7 min.

The amplified products were separated by electrophoresis in 7% polyacrylamide gel (37:1 acrylamide-bis-acrylamide) at 159 V for 1 h in a small DNA Sequencing System FB-SEQ-3545 chamber (Fisher Biotech, Pittsburg, PA). The coloring was done with ethidium bromide with subsequent treatment with silver salts.

Statistical analysis

A binary matrix was generated which was coded as absence (zero) and presence (one). The genetic similarity between

the individuals was calculated using the similarity coefficient of Nei and Li (1979), AKA Dice (1945) (Sneath and Sokal, 1973). A cluster analysis was carried out by the UPGMA method. A dendrogram was generated using Numerical Taxonomy System (NTSYS) version 2.02 (Applied Biostatics, New York, NY). To evaluate the genetic diversity, the unbiased heterozygosity and the polymorphic loci percentage were estimated using the Tools for Population Genetic Analyses (TFPGA), version 1.3 (Northern Arizona University, Flagstaff, AZ). An F statistic test was enough for the genetic differentiation with a confidence interval of 95%.

Results and discussion

The six RAM primers used for the molecular characterization of the *T. cacao* generated 127 bands that fluctuated between 12, for the AG primer, and 33, for the TG primer. The number of polymorphic loci varied between 11 and 25 for the AG and TG primers, respectively (Tab. 3).

TABLE 3. Average estimated heterozygosity and polymorphic loci percentage for the six evaluated RAM primers in 93 genotypes of *T. cacao*.

Cebador*	No. polymorphic loci	He estimated	Porcentaje polymorphic loci (95%)	Fst	SD
CGA	19	0.30	82.6	0.11	0.02
TG	25	0.34	91.7	0.16	0.05
CCA	13	0.30	92.9	0.09	0.02
CA	20	0.23	76.9	0.12	0.03
CT	16	0.35	84.2	0.13	0.04
AG	11	0.22	75.8	0.07	0.02
Total population		0.28	84.0	0.12	0.02

*See Tab. 2 for the primer sequence.

The average heterozygosity (He) for the population was 0.28, which revealed an intermediate degree of genetic diversity considering that RAMs were used as the markers. These results suggest that, in general, a significant level of genetic diversity exists in the *T. cacao* genotypes collected in Tumaco. This diversity must be valued to characterize its potential, which in turn, will be valuable for selection and breeding programs.

The highest value, for the percentage of polymorphic loci, was seen with the CCA primer (92.9%) and the lowest with the AG primer (75.8%). The highest values for polymorphism (91.7 and 92.9%) and for He (0.34 and 0.30) were seen with the TG and CCA primers, respectively. These results indicate that these primers allowed for the evaluation of the diversity and could be appropriate for use in future

research on the evaluation of the population structure and genetic diversity of *T. cacao*.

The expected heterozygosity of 0.28 was due to, despite being a small geographic region, the fact that the origin of the materials is diverse and multiplication by botanical seed and cross-pollination of the species have contributed to maintaining the genetic diversity between the evaluated materials. Duplicates were not detected in the analyzed samples, which suggests that there has been little clonal multiplication in the propagation of the material sown by the farmers. This condition of intermediate genetic diversity potentializes the opportunity of implementing suitable selection and evaluation of local materials for characteristics of quality, disease response, and productivity, which, in the future, could identify genotypes with productive potential and possible use as parentals in breeding programs for this species.

Analogous values of genetic diversity have been reported by Llor *et al.* (2009), who analyzed 322 plants of 'Nacional' cocoa collected in different geographic zones along the Ecuadorian coast. These authors determined that the genetic structure of modern populations of the 'Nacional' cocoa is characterized by high levels of heterozygosity and genetic diversity.

Chumacero de Schawe *et al.* (2013) studied the flow of pollen and genetic diversity between cultivated and wild populations of *T. cacao*, and genotyped 143 wild plants and 86 adult cultivated plants and 374 plantlets from 19 wild trees and 25 cultivated trees with nine microsatellite loci. The principal components analysis differentiated the cultivated trees from the wild trees, supporting the idea that Bolivia holds populations of truly wild *T. cacao*. The cultivated *T. cacao* had a higher level of genetic diversity than the wild *T. cacao*, which reflected the diverse origin of the cultivated trees. The relatively high exchange of pollen from the cultivated *T. cacao* to the wild *T. cacao* compromised the genetic diversity of the wild populations, making it necessary to protect vast areas of natural forests where wild *T. cacao* thrives.

The heterozygosity values fluctuated between 0.27, for San Luis Robles and Rio Rosario, and 0.15, for the Rio Mejicano. The high heterozygosity levels ($H_e=0.27$) and percentage of polymorphic loci (83.5) found in San Luis Robles were due to the high number of genotypes (50) included in this study and their high number of sampled locations: Monte Bonito, Quebrada Arriba, Nerete, Piñal Dulce, Zona Carretera, Carretera Rosario, La Lomita and Mascarey.

The high H_e values found in the watersheds of the Rio Chagüi ($H_e=0.25$) and Rio Rosario ($H_e=0.27$) were accompanied by equally high polymorphic loci values with 72.4 and 73.2%, respectively. These results indicate that, although only a few trees were sampled (17), these two regions hold high genetic variability. As a result, it is essential to implement management, conservation and strategies for the phylogenetic resources in these regions.

The low H_e (0.15) and polymorphism (36.2%) values obtained in Rio Mejicano could be affected by the low number of genotypes and *T. cacao* farms. For this reason, more sample sites and a higher number of genotypes are needed to represent the diversity in this region.

The obtained genetic differentiation coefficient ($F_{st}=0.12\pm0.02$) suggests that an intermediate genetic differentiation exists. This result confirms that there is a moderate relationship between the geographic location and the genetic group (Tab. 3).

Studies carried out by Motamayor *et al.* (2002) on the genetic diversity of the 'Criollo', with respect to that of the 'Forastero' and 'Trinitario' have demonstrated that the genetic basis of the 'Criollo' is very narrow, despite its broad geographic distribution, because the study included a high number from our collections from the south of Mexico to the north of Colombia and Venezuela. Additionally, this study also demonstrated that the 'Forastero' presented high genetic variability, which corresponded to provenance of these genotypes. In another study, Motamayor *et al.* (2002) reported that the 'Trinitario' presented high variability for the genotypes, similar to that of 'Criollo' and 'Forastero', and a high number of genotypes that presented intermediate characteristics between both groups.

The TG primer allowed for the identification of the highest polymorphism between the samples ($F_{st}=0.16$). This result indicates that it can be used for obtaining higher differentiation between *T. cacao* genotypes (Fig. 2).

Similar results have been reported by Sereno *et al.* (2006) and Zhang *et al.* (2006), who analyzed the genetic diversity of 'Forastero' populations from Brazil and Peru with microsatellite markers. They determined that the genetic differentiation between the studied groups was very low, with values ranging from 0.018 to 0.234 for Brazil and Peru, respectively. These values confirm the hypothesis that "individuals from distinct 'Forastero' populations present high polymorphism levels".




FIGURE 2. Band pattern in *T. cacao* generated by RAM microsatellite primer (TG)_n.

The Nei-Li coefficient analysis, at a Similarity level of 0.53, differentiated the population into six principal groups (Fig. 3).

The first group consisted of 76 genotypes. Samples of the 40 genotypes were collected in San Luis Robles in the following places: Carretera Rosario (4), La Lomita (5), Mascarey (1) Monte Bonito (5), Nerete (5) Piñal Dulce (7) Carretera (7), Quebrada Arriba (6). Additionally, 15 samples were collected in Rio Chagüi, from places such as La Ceiba (4), La Sirena (3), Palambí (1) and Pácora (7). Finally, four genotypes were gathered from the Rio Mejicano: Bella Vista (2) and Guayabo (2); the Mascarey and the Rio Rosario (17): Isla Grande (15) and Iscuande (2).

This group was characterized by their Cundeamor- shaped pods, with slight presence or absence of anthocyanin on the flower bud. There were plants without notable pulvinus in the leaf petioles and absence of anthocyanin in the staminodes. Likewise, these genotypes have an intermediate architecture (90° and 135°). Some of these genotypes were 1, 71, 96, 97 and 98. It is important to highlight the genotypes numbered as: 22, 69 and 73 because they showed high on-site tolerance to *Moniliophthora perniciosa* (previously *Crinipellis perniciosa* Stahel) (Ballesteros, 2011). Additionally, this group presented the highest size and length of the seed with 1.39 and 2.98 cm, respectively; an outstanding pod size (9.97 cm), as compared to the population average (9.53 cm), and a higher average fresh seed weight (5.21 g) (Ballesteros, 2011).

The clustering of genotypes collected in Mejicano river, suggests that the genetic diversity was associated with

the rivers, where the propagation of *T. cacao* by botanical seed has possibly been favored, resulting in the presence of genotypes in a heterozygous state and manifesting that the exchange of material between farmers located in the diverse rivers of this zone has been low. These results provide useful information for the identification of areas of high priority for the *ex situ* conservation of germplasm and its potential use in genetic breeding programs.

Similarly, the identification of distribution patterns of intra-species genetic variation can provided data pertaining to the spatial-temporal dynamic of the crop, which has not been comprehensively studied at the population level. The spatial structure of the diversity manifests the need for conservation measures and collections for the natural and semi-natural populations of *T. cacao* in the Tumaco locations.

Group 2, interestingly contained materials that were recently brought to the zone by government institutions. The samples were collected in the village of Nerete, in San Luis Robles and identified with the following numbers: 74, 75, 77, CR4, CR5, CR6, CR9, and CR11. This group confirms that the new materials are genetically separated from the existing traditional varieties. If these foreign trees are massively adopted, they can displace the local cocoa diversity. This could also indicate that they have a similar origin but there is variability among them, which indicated that they are the product of clonal multiplication.

The third group was represented by the clone CCN 51 from Corpoica, Tumaco. This clone presented the lowest similarity value (0.49) with respect to the other groups. This clone




FIGURE 3. Dendrogram of the genetic structure of 93 *T. cacao* genotypes based on the Nei-Li coefficient and calculated from the combined data of the six RAM microsatellite primers.

is the result of the double hybridization of ‘Trinitario’ and ‘Forastero’ genetic material of an Amazon origin and is recognized as having a yield potential and carrying resistance to common fungal diseases (Amores *et al.*, 2009). This clone was grouped with the recently introduced materials of the Tumaco zone, suggesting that these materials could have a hybrid origin, like CCN 51 and TSH 561. At the time of sampling, these materials did not present production and the genetic origin was not known.

The fourth group contained the genotypes TSH 561 and 76. The latter is characterized by few flowers, leaves and small grains, and presents high on-site tolerance to *Moniliophthora perniciosa*.

In group five, the cocoa genotypes with the numbers 42 and 23 were found, collected in the Rio Chagüi, at the locations of La Ceiba and La Sirena, respectively. These genotypes are characterized by their Cundeamor- shaped pods and absence of anthocyanin on the flower bud. According to Ballesteros (2011), they were grouped with those that presented the highest size and length of the seed with 1.39 and 2.98 cm, respectively; an outstanding pod size (9.97 cm), when compared to the population average (9.53 cm), and a higher average fresh seed weight (5.21 g).

The sixth group showed a similarity level of 0.48. This cluster encompassed clones that came from the Corpoica Germplasm Bank in Tumaco. These trees corresponded to ‘Trinitario’ type, identified as follows: ICS 60, ICS 90, and TSH 561, and the ‘Forastero’ labeled IMC 67. These genotypes were clearly separated from the traditional cocoa varieties cultivated by the local cocoa growers and those dispersed throughout the sampled zones.

The genetic distances seen in this study were smaller than those reported by Quiroz (2002), who morphologically and molecularly characterized 63 “Nacional” cocoa genotypes from Ecuador with three reference genotypes of the ‘Trinitario’ (UF-676), ‘Forastero Amazónico’ (Matina) and ‘Criollo’ (Criollo-36), through the use of AFLPs with seven primer combinations. His results indicated a genetic distance of 0.83, a very high value for individuals of the same species. However, this distance can be explained by the hybrid nature of the materials and the ample genetic diversity introduced to the zone since 1920. Likewise, these materials have a broad relationship with the ‘Refractario’ group (Zhang *et al.*, 2008).

The results found in the present study suggest that, despite being treated as materials that are located in relatively close

areas and that are being used commercially, a significant level of genetic diversity exists in the collected *T. cacao* material.

The RAM technique allowed for the grouping of the different *T. cacao* genotypes by their collection site, suggesting that these materials have been propagated from a common plant, which could be a molecular tool for the evaluation of the genetic diversity in the *Theobroma* genus.

In Colombia, studies of genetic diversity in *T. cacao* using microsatellite molecular markers are lacking, with only two studies for this species (Londoño *et al.*, 2011). Therefore, it is important to carry out studies on genetic diversity, germplasm characterization, phylogenetics, and so on, to contribute to a better understanding of the origin and genetic variability of *T. cacao* in Tumaco.

The *T. cacao* from Tumaco evidenced genetic diversity with no uniformity among the genotypes. This is an appealing characteristic for carrying out studies that aim to generate flavor profiles, phytosanitary management, genotype conservation, and sustainability of these phyto-genetic resources.

Acknowledgements

This research was sponsored by the MIDAS A-01652 -11-C-921 contract between CIAT and ARD, carried out in the Molecular Biology Laboratory of the Universidad Nacional de Colombia, Palmira and supported by the Consejos Comunitarios of the Afrocolombians in Tumaco.

Literature cited

- Amores, F., J. Agama, F. Mite, J. Jiménez, G. Loor, and J. Quiroz. 2009. EET 544 y EET 558, nuevos clones de cacao nacional para la producción bajo riego en la Península de Santa Elena. Technical Bulletin No. 134. Estación experimental Tropical Pichilingue, Instituto Nacional Autónomo de Investigaciones Agropecuarias (INIAP), Quevedo, Ecuador.
- Álvarez, E., J.F. Mejía, and T.L. Valle. 2003. Molecular and pathogenicity characterization of *Sphaceloma manihoticola* isolates from south-central Brazil. Plant Dis. 87, 1322-1328. Doi: 10.1094/PDIS.2003.87.11.1322
- Ballesteros P., W. 2011. Caracterización morfológica de árboles élite de cacao (*Theobroma cacao* L.) en el municipio de Tumaco, Nariño, Colombia. MSc thesis. Faculty of Agricultural Sciences, Universidad de Nariño, Pasto, Colombia.
- Bartley, B.G.D. 2005. The genetic diversity of cacao and its utilization. CABI Publishing, London, UK.
- Chessman, E.E. 1944. Notes on the nomenclature, classification and possible relationships of cocoa populations. Trop. Agr. 21, 144-159. Doi: 10.1038/hdy.1994.166

- Chumacero de Schawe, C., W. Durka, T. Tschardtke, I. Hensen, and M. Kessler. 2013. Gene flow and genetic diversity in cultivated and wild cacao (*Theobroma cacao*) in Bolivia. *Am. J. Bot.* 100: 2271-2279. Doi: 10.3732/ajb.1300025
- Crouzillat, D., L. Bellanger, M. Rigoreau, P. Bucheli, and V. Pétiard. 2000. Genetic structure, characterization and selection of national cocoa compared to other genetic groups. pp. 47-64. In: *Proceedings of the International Workshop on New Technologies and Cocoa Breeding*. International Group for Genetic Improvement of Cocoa - Ingenic. Kota Kinabalu, Malaysia.
- Dellaporta, S.L., J. Wood, and J.B. Hicks. 1983. A plant DNA mini-preparation: Version II. *Plant Mol. Biol. Rep.* 1, 19-21.
- Dias, L.A.S. 2001. Origin and distribution of *Theobroma cacao* L.: a new scenario. In: Dias, L.A.S. (ed.). *Genetic improvement of cacao*. FAO, <http://ecoport.org/ep?SearchType=earticleView&earticleId=197&page=-2>; consulted: November, 2014.
- Dice, L.R. 1945. Measures of the amount of ecological association between species. *Ecology* 26, 297-302. Doi: 10.2307/1932409
- Espinosa, K., M.L. Bonilla, J.E. Muñoz, A.M. Posso, and H.D. Vásquez. 2004. Colección, caracterización fenotípica y molecular de poblaciones de uchuva *Physalis peruviana*. *Biotechnol. Sector Agropec. Agroind.* 2, 72-78.
- Hantula, J., M. Dusabenyagasani, and R.C. Hamelin. 1996. Random amplified microsatellites (RAMs) - a novel method for characterizing genetic variation within fungi. *Eur. J. For. Path.* 26, 159-166. Doi: 10.1111/j.1439-0329.1996.tb00720.x
- Ji, K., D. Zhang, L.A. Motilal, M. Boccarda, P. Lachenaud, and L.W. Meinhardt. 2012. Genetic diversity and parentage in farmer varieties of cacao (*Theobroma cacao* L.) from Honduras and Nicaragua as revealed by single nucleotide polymorphism (SNP) markers. *Genet. Resour. Crop Evol.* 60, 441-453. Doi: 10.1007/s10722-012-9847-1
- Lanaud, C., A.M. Risterucci, I. Pieretti, M. Falque, A. Bouet, and P.J.L. Lagoda. 1999. Isolation and characterization of microsatellites in *Theobroma cacao* L. *Mol. Ecol.* 8, 2141-2152. Doi: 10.1046/j.1365-294x.1999.00802.x
- Lerceteau, E., J. Quiroz, J. Soria, S. Flipo, V. Pétiard, and D. Crouzilat. 1997. Genetic differentiation among Ecuadorian *Theobroma cacao* L. accessions using DNA and morphological analyses. *Euphytica* 95, 77-87. Doi: 10.1023/A:1002993415875
- Londoño M., J., D.M. Gil V., S.B. Aguilar M., F.A. Rivera P., and G.A. López G. 2011. Caracterización molecular de clones de *Theobroma cacao* L., por medio de marcadores moleculares microsatélites. *Luna Azul* 32, 52-60. Doi: 10.17151/luaz.2011.32.6
- Loor, R.G. 2007. Contribution à l'étude de la domestication de la variété de cacaoyer Nacional d'Équateur: recherche de la variété native et de ses ancêtres sauvages. PhD thesis. Ecole Nationale Supérieure d'Agronomie (ENSAM), Montpellier, France.
- Loor, R.G., A.M. Risterucci, B. Courtois, O. Fouet, M. Jeanneau, E. Rosenquist, F. Amores, A. Vasco, M. Medina, and C. Lanaud. 2009. Tracing the native ancestors of the modern *Theobroma cacao* L. population in Ecuador. *Tree Genet. Genomes* 5, 421-433. Doi: 10.1007/s11295-008-0196-3
- Mahuku, G.S., M.A. Henríquez, J. Muñoz, and R.A. Buruchara. 2002. Molecular markers dispute the existence of the Afro-Andean group of the bean angular leaf spot pathogen, *Phaeoisariopsis griseola*. *Phytopathology* 92, 580-589. Doi: 10.1094/PHYTO.2002.92.6.580
- Marcano, M., S. Morales, M.T. Hoyer, B. Courtois, A.M. Risterucci, O. Fouet, T. Pugh, E. Cros, V. Gonzalez, M. Dagert, and C. Lanaud. 2009. A genomewide admixture mapping study for yield factors and morphological traits in a cultivated cocoa (*Theobroma cacao* L.) population. *Tree Genet. Genomes* 5, 329-337. Doi: 10.1007/s11295-008-0185-6
- Motamayor, J.C., A.M. Risterucci, P.A. Lopez, C.F. Ortiz, A. Moreno, and C. Lanaud. 2002. Cacao domestication I: the origin of the cacao cultivated by the Mayas. *Heredity* 89, 380-386. Doi: 10.1038/sj.hdy.6800156
- Motamayor, J.C., P. Lachenaud, J.W. da Silva e Mota, R. Loor, D.N. Kuhn, J.S. Brown, and R.J. Schnell. 2008. Geographic and genetic population differentiation of the Amazonian chocolate tree (*Theobroma cacao* L.). *PloS One* 3, e3311. DOI: 10.1371/journal.pone.0003311
- Motilal, L.A., D. Zhang, P. Umaharan, S. Mischke, V. Mooleedhar, and L.W. Meinhardt. 2010. The relic Criollo cacao in Belize - genetic diversity and relationship with Trinitario and other cacao clones held in the International Cocoa Genebank, Trinidad. *Plant Genet. Resour.* 8, 106-115. Doi: 10.1017/S1479262109990232
- Municipio de Tumaco. 2006. Plan de Ordenamiento Territorial (POT). Tumaco, Colombia. pp. 25-26.
- Muñoz F., J.E., A.C. Morillo C., and Y. Morillo C. 2008. Microsatélites amplificados al azar (RAM) en estudios de diversidad genética vegetal. *Acta Agron.* 57, 219-226.
- Nei, M., and W.H. Li. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA* 76, 5269-5273. Doi: 10.1073/pnas.76.10.5269
- Palacio G., D. 2007. Caracterización (fingerprinting) de clones de *Theobroma cacao* L., de interés comercial de la zona cafetera marginal baja (ZCMB) utilizando marcadores tipo fAFLP. MSc thesis. Faculty of Sciences, Universidad de Los Andes, Bogotá.
- Pautasso, M. 2009. Geographical genetics and the conservation of forest trees. *Perspect. Plant Ecol. Evol. Syst.* 11, 157-189. Doi: 10.1016/j.ppees.2009.01.003
- Pound, F.J. 1945. A note on the cocoa population of South America. In: *Rep. Proc. Cocoa Res. Conf. London*. Colonial 192, 95-97. Reprinted 1982 in *Arch. Cacao Res.* 1, 93-97.
- Quiroz V., J.G. 2002. Caracterización molecular y morfológica de genotipos superiores de cacao nacional (*Theobroma cacao* L.) de Ecuador. MSc thesis. Tropical Agricultural Research and Higher Education Center (CATIE). Turrialba, Costa Rica.
- Sereno, M.L., P.S.B. Albuquerque, R. Vencovsky, and A. Figueira. 2006. Genetic diversity and natural population structure of cacao (*Theobroma cacao* L.) from the Brazilian Amazon evaluated by microsatellite markers. *Conserv. Genet.* 7, 13-24. Doi: 10.1007/s10592-005-7568-0
- Sneath, P.H.A. and R.R. Sokal. 1973. Numerical taxonomy: the principles and practice of numerical classification. W.H. Freeman and Co., San Francisco, CA.
- Thomas, E., M. Van Zonneveld, J. Loo, T. Hodgkin, G. Galluzzi, and J. Van Etten. 2012. Present spatial diversity patterns of *Theobroma cacao* L. in the neotropics reflect genetic differentiation

- in Pleistocene refugia followed by human-influenced dispersal. PLoS ONE 7, e47676. Doi:10.1371/journal.pone.0047676
- Vallejo, G. and E. Peña. 2006. Centro Regional de Investigación el Mira. Brochure No. 47 Corpoica, Tumaco, Colombia.
- Zhang, D., E. Arevalo-Gardini, S. Mischke, L. Zúñiga-Cernades, A. Barreto-Chavez, and J. Adriazola del Aguila. 2006. Genetic diversity and structure of managed and semi-natural populations of cocoa (*Theobroma cacao* L.) in the Huallaga and Ucayali Valleys of Peru. Ann. Bot. 98, 647-655. Doi: 10.1093/aob/mcl146
- Zhang, D., M. Boccara, L.A. Motilal, D.R. Butler, P. Umaharan, S. Mischke, and L.W. Meinhardt. 2008. Microsatellite variation and population structure in the “Refractario” cacao of Ecuador. Conserv. Genet. 9, 327-337. Doi: 10.1007/s10592-007-9345-8
- Zhang, D., W.J. Martínez, E.S. Johnson, E. Somarriba, W. Phillips-Mora, C. Astorga, S. Mischke, and L.W. Meinhardt. 2012. Genetic diversity and spatial structure in a new distinct *Theobroma cacao* L. population in Bolivia. Genet. Res. Crop Evol. 59, 239 - 252. Doi: 10.1007/s10722-011-9680-y