



Agronomía Colombiana

ISSN: 0120-9965

agrocolfabog@gmail.com

Universidad Nacional de Colombia
Colombia

Narváez-Cuenca, Carlos-Eduardo; Silva-Bustos, Kelly-Johanna; Restrepo-Sánchez, Luz-Patricia

Effects of thermal processing combined with sucrose on the vitamin C content, total phenolic content, antioxidant activity, and sensory characteristics of arazá (*Eugenia stipitata* McVaugh) purée during frozen storage

Agronomía Colombiana, vol. 33, núm. 2, 2015, pp. 212-219

Universidad Nacional de Colombia
Bogotá, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=180341167010>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Effects of thermal processing combined with sucrose on the vitamin C content, total phenolic content, antioxidant activity, and sensory characteristics of arazá (*Eugenia stipitata* McVaugh) purée during frozen storage

Efecto del procesamiento térmico en combinación con sacarosa en el contenido de vitamina C, fenoles totales, actividad antioxidante y características sensoriales durante el almacenamiento de puré de arazá congelado (*Eugenia stipitata* McVaugh)

Carlos-Eduardo Narváez-Cuenca¹, Kelly-Johanna Silva-Bustos¹, and Luz-Patricia Restrepo-Sánchez¹

ABSTRACT

The effects of a 80°C, 1 minute thermal treatment (*H* treatment) and of the same treatment combined with sucrose (*SH* treatment) on the chemical and sensory qualities of arazá purée were evaluated during 4 months of storage at -20°C. For the control, an untreated sample (*C*) was included. The chemical qualities evaluated were vitamin C content, total phenolic compounds (TPC), and antioxidant activity (AoA), which were measured with the ferric reducing ability of plasma (FRAP), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), and 1,1-diphenyl-2-picrylhydrazyl (DPPH), while a trained panel was used for the sensory quality evaluation. After 4 months of frozen storage, *SH* was more effective than *H* at controlling the loss of chemical quality. *SH* provided a similar FRAP-AoA (89%) and vitamin C content (87%), as well as a higher TPC (145%), ABTS-AoA (272%), and DPPH-AoA (115%), when compared to *C* before frozen storage. The total sensory qualities in both the *SH* purée (15±1) and the *H* purée (16±1) after 4 months at -20°C were comparable to those of the *C* purée before the frozen storage (18±2). Therefore, using sucrose combined with a thermal treatment and subsequent frozen storage preserves the chemical and sensory qualities of arazá purée.

Key words: thermal treatment, tropical fruit, combined methods, Amazonian fruit.

RESUMEN

Se evaluó el efecto del tratamiento térmico a 80°C durante 1 minuto (tratamiento *H*) y este mismo tipo de tratamiento combinado con adición de sacarosa (tratamiento *SH*) sobre la calidad química y sensorial de puré de arazá durante 4 meses de almacenamiento a -20°C. Además, se incluyó un control sin tratamiento térmico ni sacarosa (control *C*). Los parámetros químicos evaluados fueron: vitamina C, compuestos fenólicos totales (TPC) y actividad antioxidante (AoA) medida por los métodos de la capacidad de reducción férrica del plasma (FRAP), ácido 2,2'-azino-bis (3-etilbenzotiazolina-6-sulfónico) (ABTS) y 1,1-difenil-2-picrilhidrazil (DPPH). La calidad sensorial se evaluó con un panel entrenado. Después de cuatro meses de almacenamiento congelado el tratamiento *SH* fue más efectivo que el *H* en el control de la calidad química. *SH* proporcionó valores de AoA-FRAP (89%) y contenido de vitamina C (87%) similares, como también mayores valores de TPC (145%), AoA-ABTS (272%) y AoA-DPPH (115%) al compararlos con el control *C* antes del almacenamiento congelado. La calidad sensorial del puré *SH* (15±1) y del puré *H* (16±1) después de 4 meses a -20°C fue comparable a la del puré *C* antes del almacenamiento congelado (18±2). El uso de sacarosa combinado con tratamiento térmico y almacenamiento congelado preserva la calidad sensorial y química de puré de arazá.

Palabras clave: tratamiento térmico, fruta tropical, métodos combinados, frutas amazónicas.

Introduction

Arazá (*Eugenia stipitata* McVaugh) is an exotic, tropical fruit that is grown in Latin American countries, such as Colombia, Peru, and Brazil. The fruit is a fleshy berry (4-7 cm in diameter) with 6 to 16 seeds (1-4 cm in diameter) per fruit. It has an attractive flavor, a vitamin C content [8.9-26.6 mg/100 g fresh weight (FW)] (García-Reyes and

Narváez-Cuenca, 2010; Contreras-Calderón *et al.*, 2011) that is comparable to those of pineapple (*Ananas comosus*) and banana (*Musa acuminata* AAA) (Hernández *et al.*, 2006), and a total phenolic content (TPC) (19.3-111.0 mg gallic acid equivalents/100 g FW) similar to those found in tropical fruits such as mango (*Mangifera indica*) and granadilla (*Passiflora ligularis*) (García-Reyes and Narváez-Cuenca, 2010; Contreras-Calderón *et al.*, 2011; Garzón *et*

al., 2012), as measured with a Folin-Ciocalteu (FC) reagent (Vasco *et al.*, 2008).

Attempts to delay the browning of the skin and excessive softening of fresh arazá fruits have been described in the literature (Narváez-Cuenca, 2003; Hernández *et al.*, 2009; Carrillo *et al.*, 2011). When stored at 20°C, green-mature fruits reach sensorial ripeness after one week with a subsequent decay of the sensory quality several days later (Hernández *et al.*, 2009; Carrillo *et al.*, 2011). The refrigeration of green-mature fruits at 12°C, with or without exposure to 1-methylcyclopropene, reduces the loss of organic acids and extends the sensory quality of arazá fruits for up to two weeks (Carrillo *et al.*, 2011). Heat shock prior to refrigeration when using green-mature fruits was also useful in extending arazá fruit sensory quality for up to two weeks (Narváez-Cuenca, 2003).

The arazá fruit is predominantly used to prepare nectars, jams, and candies, and it is rarely consumed whole. Based on frozen storage experiments that were conducted over a period of 2 months (Millán *et al.*, 2007; García-Reyes and Narváez-Cuenca 2010), the preservation of arazá fruit as a frozen purée was promising. Millán *et al.* (2007) concluded that the combination of thermal treatments of arazá purée at 80°C for 1 min, quick freezing (by immersion in liquid nitrogen), and storage at -20°C preserved the sensorial and chemical [vitamin C, TPC, and antioxidant activity (AoA)] qualities of the arazá purée. These authors also indicated that the use of slow thawing yields purées with better textures than those prepared with fast thawing. Silva *et al.* (2011) tested the effect of the addition of sucrose (in concentrations ranging from 0 to 30%, *w/w*) to arazá purées on the sensorial quality of the stored frozen product and concluded that the presence of 20% (*w/w*) sucrose controlled the degradation of the sensory characteristics for the one-month storage period that was assessed. In the mentioned studies, the use of sucrose and slow freezing was comparable to the use of thermal processing in combination with quick freezing in terms of the sensorial texture stability of the arazá purée. Until now, sucrose has not been used in combination with thermal processing and frozen storage trials have not lasted longer than two months in the evaluation of arazá purée. Because there are up to three harvests of arazá fruit per year, the present research was conducted over a period of 4 months using frozen storage and evaluating the effect of thermal processing (at 80°C for 1 min) combined with sucrose (20% *w/w*) and the same thermal processing without sucrose on the chemical and sensorial qualities of arazá purée.

Materials and methods

Materials

The 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), 2,2'-azino-bis-(3-3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), dithiothreitol, 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,4,6-tripyridyl-S-triazine (TPTZ), and Folin Ciocalteu (FC) reagent were supplied by Sigma (St. Louis, MO). Sucrose was obtained from Ingenio Manuelita (Bogota, Colombia) and all of the other chemicals came from Panreac Química (Barcelona, Spain). Ripe arazá fruits (fully-yellow skin color) (Carrillo *et al.*, 2011) were obtained from superstores in Bogota, Colombia. Vendors reported the fruit came from Cundinamarca, Colombia, but the exact location was unknown.

Preparation of samples and storage

The ripe arazá fruits were randomly split into nine lots (10-16 fruits per lot). Each lot was washed with tap water and peeled by hand. The seeds were removed and the pulp was puréed for 10 s in a domestic blender. Nine samples of purée (2 kg each) were obtained and immediately processed. Three samples of the purée were treated thermally in a kettle equipped with a stirrer. These samples were held at 80°C for 1 min once the temperature at the central part of the 2 kg-purée reached said temperature and immediately cooled to 15°C by water circulation (*H* treatment) (García-Reyes and Narváez-Cuenca, 2010). Three other samples were mixed with sucrose to obtain a final concentration of 20% (*w/w*) sucrose and then heated as described for the *H* treatment (*SH* treatment). Finally, three samples were used as the control (*C*), where neither sugar nor heating were included. All samples (*H* and *SH* treatments, and *C*) were packed in polyethylene bags (100 g of purée per bag), slowly frozen (0.2°C/min), and stored in darkness at -20°C for four months. After storage, the chemical and sensory quality parameters were evaluated. The following chemical analyses were performed on the *H* and *SH* treatments and the *C* samples: vitamin C content, TPC, and AoA (measured by ABTS, FRAP, and DPPH). The sensorial quality of the nectars prepared from the purée samples was measured by a trained panel. Measurements were performed before freezing and after 1, 2, 3, and 4 months of storage at -20°C. Except for day 0 (before freezing), the samples were thawed at room temperature and then analyzed.

Analysis of the chemical quality

Vitamin C. The vitamin C extraction and quantification were based on previous studies (Gökmen *et al.*, 2000). To extract the vitamin C, 1 g of sample was added to 4 mL of deionized water. The mixture was vortexed for 2 min and

filtered first through a black band paper filter and then through a 0.45 µm Millipore (Milford, MA) disposable filter. The filtrate was used for the vitamin C determination, which included the measurement of both ascorbic acid and dehydroascorbic acid in a single run. The dehydroascorbic acid present in the filtrate was reduced to ascorbic acid by treatment with 1 mg/mL dithiothreitol over a period of 2 h, in darkness and at room temperature. The vitamin C content was determined by reverse-phase high-performance liquid chromatography (RP-HPLC). The chromatographic system consisted of a HiChrom C18 stainless steel column (250 x 4 mm I.D.; 5 µm particle size; Reading, UK) operated at 18°C. The eluate was monitored at 254 nm using an UV-visible detector operating at 40°C. The eluent was 200 mM KH₂PO₄, adjusted to pH 2.4 with H₃PO₄. The flow rate was 0.5 mL min⁻¹. The sample volume for injection was 20 µL. A calibration curve was constructed for ascorbic acid solutions ranging from 10 to 50 mg mL⁻¹ (five data points, $r^2 = 0.999$). The results were expressed as mg of vitamin C per 100 g of FW purée.

Total phenolic content and antioxidant activity. For the TPC and AoA, an extraction was performed following previous studies (García-Reyes and Narváez-Cuenca, 2010). A 2.5 g of purée were mixed with 5 mL of 80% (v/v) aqueous methanol and shaken for 1 h at 200 rpm. The resulting mixture was centrifuged (4,000 xg, 5 min at 4°C) and the supernatant was kept at 4°C. A second extraction was performed. The two supernatants were mixed, and the volume was brought to 10 mL in a volumetric flask, topped off with distilled water, and used for the TPC and AoA analyses. The undiluted extracts were analyzed for the TPC and the extracts were diluted from 1:1 to 1:5 (extract: 80% aqueous methanol, v/v) for the AoA measurements.

The TPC was determined using the FC method described by Velioglu *et al.* (1998). A 100 µL aliquot of the aqueous methanol extract (80% v/v) was mixed with 750 µL of FC reagent (10% v/v). After 5 min, 750 µL of a NaHCO₃ (6% w/v) solution were added. After 90 min, the absorbance of the reaction mixture was measured at 725 nm. The results were expressed as gallic acid equivalents (mg GAE/100 g FW purée). The calibration curve for gallic acid was constructed in concentrations ranging from 0.01 to 0.30 mg mL⁻¹ (five data points, $r^2 = 0.998$).

The AoA of the extracts was measured with three methods: ABTS, FRAP, and DPPH. The ABTS assay was based on Re *et al.* (1999). A mixture of 7.0 mM ABTS and 2.45 mM potassium persulfate (final concentration) was prepared and stored in darkness for 16 h. The resulting ABTS^{•+}

solution was diluted with ethanol to reach an absorbance of 0.70 at 734 nm. Aliquots (10 µL) of the diluted extracts were incubated with 1,000 µL of the ABTS^{•+} solution for 60 min at 30°C and the absorbance was monitored at 734 nm every 5 s.

The FRAP assay was carried out according to the method described by Benzie and Strain (1996). The FRAP reagent was prepared by mixing 25.0 mL of sodium acetate buffer (300 mM, pH 3.6) with 2.5 mL of TPTZ (10 mM) in HCl (40 mM) and 2.5 mL of ferric chloride (20 mM). The FRAP reagent (900 µL) was mixed with 90 µL of distilled water and 30 µL of diluted extract. The mixture was incubated at 37°C and the absorbance was monitored at 593 nm every 5 s for 45 min.

The DPPH assay was performed following Sánchez-Moreno *et al.* (1998). A methanolic stock solution of 50 mM DPPH was diluted with methanol to obtain a working DPPH solution with an absorbance of 1.10 at 515 nm. The diluted extract (25 µL) was mixed with 975 µL of a working methanolic DPPH solution. The mixture was incubated at 20°C and its absorbance was measured at 515 nm every 5 s for 1 h. The scavenging effect (as a percentage) was calculated with $[(A_1 - A_2)/A_0] \times 100$, where A_0 was the absorbance of the control (without extract), A_1 was the absorbance in the presence of the extract once the plateau was reached, and A_2 was the absorbance without DPPH. In the three different assays, ABTS, FRAP, and DPPH standard curves were created with Trolox in concentrations ranging from 100 to 1,500 µM (five data points, $r^2 = 0.999, 0.989, 0.975$, respectively). In all of the cases, the absorbance reached a plateau during the given incubation time. The results were expressed as µmol of Trolox per g of FW purée.

Analysis of the sensory quality

The nectars were prepared by blending the purée, water, and sucrose to obtain a mixture with 25% (w/w) arazá purée and 12°Brix. Quantitative descriptive analysis (QDA) of the prepared nectars was carried out by a trained panel of seven members. QDA was performed for color and general aspects, as well as aroma, taste, and texture of the arazá purée. The panelists were both male and female, non-smokers, and 40 to 55 years old; they were trained twice per week over a period of two months on the evaluation of the quality parameters (Tab. 1) by preparing nectars with fresh purée, purée boiled for 1-5 min, and purée submitted to 1-3 cycles of frozen storage for 1 d with subsequent thawing. The evaluation was performed in a room designed specifically for sensory evaluation with individual booths and controlled, white lighting. In each session, each panelist

tested nine randomly codified samples. Each panelist was asked to rate the product using intermediate scores (Tab. 1). Each parameter had a different weight according to its importance, as defined in a previous study (Millán *et al.*, 2007). The total quality was calculated as the sum of the different parameters.

TABLE 1. Scores for the sensory evaluation of the nectar prepared from the arazá purée.

Score	Description
Color and general aspect	
2	Brilliant yellow color and very homogeneous material
1	Slight yellow, some suspended material
0	Atypical color, bleached yellow, sediment
Aroma	
7	Acidic, fresh, fruity
5	Slightly acidic, fresh, fruity
0	Acetic, moldy
Taste	
6	Acidic, good balance acidic/sweet
4	Slightly acidic, slightly boiled taste
0	Extremely boiled taste, metallic
Texture	
5	Viscous, homogeneous
3	Disaggregated, slightly homogeneous
0	Extremely fluid

Statistical analysis

The statistical analyses were performed using Statgraphics Plus® 5.1 for Windows (Manugistics, Rockville, MD). The mean values and their standard deviations were reported. A two-way analysis of variance was performed with factor A as the two treatments and the control, *H*, *SH*, and *C*, and factor B as the five periods of frozen storage, 0 (before freezing), 1, 2, 3, and 4 months. The comparisons among the means were performed with Tukey's test at a significance level of $P \leq 0.05$.

Results and discussion

Chemical quality of fresh arazá purée

At day 0, the chemical quality of the fresh, untreated arazá purée was assessed by quantifying the vitamin C content, TPC, and AoA (Fig. 1A-E). The vitamin C content (20.3 ± 0.8 mg/100 g FW) (Fig. 1A) was within the range previously reported for fresh arazá purée (8.9–26.6 mg/100 g FW) (García-Reyes and Narváez-Cuenca, 2010; Contreras-Calderón *et al.*, 2011).

The TPC (Fig. 1B), expressed as mg GAE/100 g FW, of the fresh, untreated purée (26.5 ± 3.6) was lower than the previously reported values (35.0 ± 8.6 and 111.0 ± 3.6) when the extraction was performed with 80% (v/v) aqueous methanol (García-Reyes and Narváez-Cuenca, 2010).

The AoA of the fresh arazá purée was higher when measured by the FRAP method (8.8 ± 1.1 μ mol Trolox/g FW of purée) (Fig. 1D) than when measured by the ABTS method (3.2 ± 0.5 μ mol Trolox/g FW of purée) (Fig. 1C) or by the DPPH method (3.4 ± 0.3 μ mol Trolox/g FW of purée) (Fig. 1E). This trend indicates that the antioxidants present in fresh arazá purée have a better reducing capacity (as measured by FRAP) than the free radical scavenging capacity (as measured by ABTS and DPPH). This trend might be related to the polyphenolic composition of the extract. Caffeic acid, reported in the skin and pulp of arazá (Cuellar *et al.*, 2013) at a concentration of 1 mg L^{-1} , yielded a FRAP-AoA value that was 1.4-times higher than the value obtained by either DPPH-AoA or ABTS-AoA (Maqsood and Benjakul, 2010). While the FRAP-AoA found in the fresh arazá purée was within the range reported for 24 fruits from Colombia (ranging from 3.2 to 175 μ mol Trolox/g FW), the ABTS-AoA was below the values found in the same 24 Colombian fruits (ranging from 6.3 to 144 μ mol Trolox/g FW) (Contreras-Calderón *et al.*, 2011). The fresh arazá purée had a DPPH-AoA value within the range of values reported in 19 tropical fruits from Ecuador (ranging from 0.3 to 76 μ mol Trolox/g FW) (Vasco *et al.*, 2008).

The higher values for the vitamin C content, TPC, and ABTS-AoA previously reported for fresh arazá purée (García-Reyes and Narváez-Cuenca, 2010), when a similar extraction method was used, might be the result of differences in the year, harvest location, and storage conditions, as shown in other fruits (Al-Turki *et al.*, 2010; Guzmán-Maldonado *et al.*, 2010). The lower TPC, FRAP-AoA, ABTS-AoA, and DPPH-AoA values found in the arazá purée when the extraction was performed with 100% (v/v) acetone (Garzón *et al.*, 2012) as compared to the values found by us when the extract was prepared in 80% (v/v) aqueous methanol suggest a more important contribution from the hydrophilic antioxidants, such as hydroxybenzoic and phenolic acids, rather than from the hydrophobic antioxidants, such as carotenoids, which are present in arazá fruits (Garzón *et al.*, 2012; Cuellar *et al.*, 2013).

Chemical quality of treated arazá purée before frozen storage

The effect of thermal treatments or sucrose with thermal treatments before storage at -20°C on the chemical quality




FIGURE 1. Effects of thermal treatments (light grey bars) and sucrose (20% w/w) combined with thermal treatments (dark grey bars) on the vitamin C content (A), total phenolic content (B), ABTS-antioxidant activity (C), FRAP-antioxidant activity (D), DPPH-antioxidant activity (E), and total sensory quality (F) of arazá purée stored at -20°C . The control samples are represented by the white bars and the data bars are the means of three replicates with their standard errors. Different lowercase letters indicate significant differences ($P \leq 0.05$) based on Tukey's comparison test within each treatment or the control during storage. Asterisks denote significant differences ($P \leq 0.05$) based on Tukey's comparison test between the treatments and the control at the same storage time.

of arazá purée (Fig. 1A-E). Compared to the C sample, there was no significant effect of the H treatment on the vitamin C content, TPC, ABTS-AoA, or FRAP-AoA. The SH treatment did not significantly change the TPC or FRAP-AoA. In contrast, at time 0, while the H treatment yielded a statistically higher DPPH-AoA ($4.0 \pm 0.2 \mu\text{mol Trolox/g FW}$ of purée) as compared to the control ($3.4 \pm 0.3 \mu\text{mol Trolox/g FW}$ of purée), the SH treatment resulted in a statistically higher vitamin C content ($23.0 \pm 0.4 \text{ mg/100}$

g FW), ABTS-AoA ($5.1 \pm 0.7 \mu\text{mol Trolox/g FW}$ of purée), and DPPH-AoA ($4.0 \pm 0.2 \mu\text{mol Trolox/g FW}$ of purée) as compared to the control ($20.3 \pm 0.8 \text{ mg vitamin C/100 g FW}$, $3.2 \pm 0.5 \mu\text{mol Trolox/g FW}$ of purée as measured by ABTS, and $3.4 \pm 0.3 \mu\text{mol Trolox/g FW}$ of purée as measured by DPPH).

The increase observed for the chemical quality of the fresh arazá purée after application of the H or SH treatments

suggests a positive role for both the thermal treatments and the sucrose additives. The stability of the vitamin C content observed in the thermal treatment of the arazá purée is in agreement with the lack of significant vitamin C loss found after thermal treatments on guava (*Psidium guajava* L. cv. Chung shan) purée (Yen and Lin, 1996) or watercress (*Nasturtium officinale* R. Br.) leaves (Gonçalves *et al.*, 2009), but contrasts the degradation of this compound after infrared thermal treatments on fresh-cut mango (Sogi *et al.*, 2012). During the thermal treatment, phenolics and other compounds contributing to the AoA that were part of the insoluble bound fraction must have been released from the purée matrix, as previously shown by Sogi *et al.* (2012) in infrared heated fresh-cut mango. These authors found an increase in the TPC and AoA, as measured by DPPH, FRAP, and ORAC assays, as a result of the infrared treatment. The stabilizing effect of sucrose on the chemical qualities of arazá purée corroborates a previous report from Kopjar *et al.* (2008). These authors reported that the addition of trehalose to strawberry cream improves the stability of the anthocyanins in the product prepared by evaporation at 80°C.

Chemical quality of treated arazá purée during frozen storage

During frozen storage of the C sample, there was a decrease ($P \leq 0.05$) in the vitamin C content, TPC, FRAP-AoA, and DPPH-AoA, and an increase ($P \leq 0.05$) in the ABTS-AoA (Fig. 1A-E). Interestingly, the H and SH treatments yielded greater values ($P \leq 0.05$) than the C sample for each chemical quality parameter evaluated during frozen storage (Fig. 1A-E).

The vitamin C content decreased in the C sample during the frozen storage, with a calculated half-life time of 93 d after applying a first order kinetic model. The degradation of the vitamin C was retarded with the H and SH treatments. The half-life of the vitamin C was extended to 166 d with the H treatment and 278 d with the SH treatment. After 4 months of frozen storage, the vitamin C content in the C, H treated, and SH treated purée samples represented 40, 64, and 87%, respectively, as compared to the vitamin C content of the fresh, untreated purée sample. The ratio values (expressed as a percentage) between the TPC after four months of frozen storage of the C, H, and SH treatments and the TPC of the fresh, untreated purée sample were calculated. The ratio values for ABTS-AoA, FRAP-AoA, and DPPH-AoA were calculated as well. The SH treated purée samples had the highest ratio values (145 for TPC, 272 for ABTS-AoA, 89 for FRAP-AoA, and 115% for DPPH-AoA), followed by the H treated purée samples

(114, 178, 91, and 88%, respectively) and then the C purée samples (66, 116, 67, and 76%).

In the present research, we found that both the H and SH treatments controlled the degradation of vitamin C, TPC, and AoA, as compared to the control. Furthermore, the SH treatment exhibited better control of chemical quality losses during frozen storage, as compared to the H treatment. The observed protective effect of sucrose on phenolic compounds and vitamin C upholds previous results for frozen fruits (Oszmiański *et al.*, 2009; Rincon and Kerr, 2010). The use of sugars, *e.g.*, glucose, fructose and sucrose, to coat whole fruits or to mix with fruit purées prior to freeze-drying, freezing, or evaporation at 80°C has positive effects on the chemical quality (Kopjar *et al.*, 2008; Galmarini *et al.*, 2009; Kopjar *et al.*, 2009; Loncaric *et al.*, 2014). These sugars effectively increase both aroma and color retention and reduce the loss of TPC, anthocyanins, and AoA during storage at either room temperature or freezing temperatures. The positive effect of sugars on the chemical quality stability has also been related to an increase in the transition glass temperature, which is associated with low shrinkage, collapse, and crystallization of freeze-dried products (Roos, 1995). The increase in the FRAP-AoA of the solutions containing sucrose has been attributed to the production of non-enzymatic browning reaction products during heating at a low pH (Tsai *et al.*, 2005). The potential susceptibility of phenolic compounds to a thermal treatment was overcome by the higher stability of these compounds during frozen storage of the arazá purée. The stabilizing effect of thermal treatments on individual phenolic compounds was previously shown in grapefruit juice during frozen storage (Igual *et al.*, 2011).

Sensory quality of treated arazá purée during frozen storage

The trained panelists indicated that the H and SH treatments generated a boiled taste (day 0); nevertheless, the effect of the treatments was not significant when the total quality of the nectars was evaluated (Fig. 1F). The total sensory quality of the C sample was negatively affected during frozen storage, moving from an initial value (before frozen storage) of 18 ± 2 to 8 ± 3 after 4 months. No statistically significant differences were obtained between the H and the SH treated samples at any storage time when evaluating the total sensory quality. The total sensory quality of the H and SH treated samples was greater than the quality of the C sample during frozen storage. After 4 months, the H (16 ± 1) and SH (15 ± 1) treatments produced total quality levels that were comparable to that of the fresh, untreated purée (18 ± 2). The use of sugars has been reported to have

a positive effect on the retention of sensory qualities, such as aroma, color, and texture in whole fruits, fruit purées or fruit juice during storage at room temperature or at -18°C (Kopjar *et al.*, 2008; Kopjar *et al.*, 2009; Kopjar *et al.*, 2012). Furthermore, thermal processing has been reported to enhance sensory quality by eliminating throat irritation, as compared to fresh purées (Ledeker *et al.*, 2014).

Conclusions

After frozen storage of the HS treated purée (20% (w/w) sucrose plus thermal treatment) for four months, the vitamin C content, TPC, AoA, and sensorial quality were comparable to or higher than those of the control purée before frozen storage. The use of 20% (w/w) sucrose with a thermal treatment in combination with frozen storage is, therefore, useful in extending both the chemical and sensory qualities of arazá purée. Further studies to describe the changes in the phenolic composition during frozen storage might be of interest when the biological activity of such compounds is taken into account.

Literature cited

- Al-Turki, S., M.A. Shahba, and C. Stushnoff. 2010. Diversity of antioxidant properties and phenolic content of date palm (*Phoenix dactylifera* L.) fruits as affected by cultivar and location. *J. Food Agric. Environ.* 8, 253-260.
- Benzie, I.F.F., and J.J. Strain. 1996. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Anal. Biochem.* 239, 70-76. Doi: 10.1006/abio.1996.0292
- Carrillo, M.P., M.S. Hernández, J. Barrera, O. Martínez, and J.P. Fernández-Trujillo. 2011. 1-Methylcyclopropene delays arazá ripening and improves postharvest fruit quality. *LWT - Food Sci. Tech.* 44, 250-255. Doi: 10.1016/j.lwt.2010.05.029
- Contreras-Calderón, J., L. Calderón-Jaimes, E. Guerra-Hernández, and B. García-Villanova. 2011. Antioxidant capacity, phenolic content and vitamin C in pulp, peel and seed from 24 exotic fruits from Colombia. *Food Res. Int.* 44, 2047-2053. Doi: 10.1016/j.foodres.2010.11.003
- Cuellar, F.A., E. Ariza, C. Anzola, and P. Restrepo. 2013. Estudio de la capacidad antioxidante del arazá (*Eugenia stipitata* Mc Vaugh) durante la maduración. *Rev. Colomb. Quím.* 42, 213-235.
- Galmarini, M.V., C. Schebor, M.C. Zamora, and J. Chirife. 2009. The effect of trehalose, sucrose and maltodextrin addition on physicochemical and sensory aspects of freeze - dried strawberry puree. *Int. J. Food Sci. Tech.* 44, 1869-1876. Doi: 10.1111/j.1365-2621.2008.01890.x
- García-Reyes, R.-H., and C.-E. Narváez-Cuenca. 2010. The effect of pasteurization on the quality of frozen arazá (*Eugenia stipitata* Mc Vaugh) pulp. *J. Food Qual.* 33, 632-645. Doi: 10.1111/j.1745-4557.2010.00348.x
- Garzón, G.A., C.-E. Narváez-Cuenca, R.E. Kopec, A.M. Barry, K.M. Riedl, and S.J. Schwartz. 2012. Determination of carotenoids, total phenolic content, and antioxidant activity of arazá (*Eugenia stipitata* Mc Vaugh), an Amanoian fruit. *J. Agric. Food Chem.* 60, 4709-4717. Doi: 10.1021/jf205347f
- Gökmen, V., N. Kahraman, N. Demir, and J. Acar. 2000. Enzymatically validated liquid chromatographic method for the determination of ascorbic and dehydroascorbic acids in fruit and vegetables. *J. Chrom. A.* 881, 309-316. Doi: 10.1016/S0021-9673(00)00080-7
- Gonçalves, E.M., R.M.S. Cruz, M. Abreu, T.R.S. Brandão, and C.L.M. Silva. 2009. Biochemical and colour changes of watercress (*Nasturtium officinale* R. Br.) during freezing and frozen storage. *J. Food Eng.* 93, 32-39. Doi: 10.1016/j.jfoodeng.2008.12.027
- Guzmán-Maldonado, S.H., G. Herrera-Hernández, D. Hernández-López, R. Reynoso-Camacho, A. Guzmán-Tovar, F. Vaillant, and P. Brat. 2010. Physicochemical, nutritional and functional characteristics of two underutilised fruit cactus species (*Myrtillocactus*) produced in central Mexico. *Food Chem.* 121, 381-386. Doi: 10.1016/j.foodchem.2009.12.039
- Hernández, Y., M.G. Lobo, and M. González. 2006. Determination of vitamin C in tropical fruits: a comparative evaluation of methods. *Food Chem.* 96, 654-664. Doi: 10.1016/j.foodchem.2005.04.012
- Hernández, M.S., J. Barrera, O. Martínez, and J.P. Fernández-Trujillo. 2009. Postharvest quality of arazá fruit during low temperature storage. *LWT - Food Sci. Tech.* 42, 879-884. Doi: 10.1016/j.lwt.2008.11.009
- Igual, M., E. García-Martínez, M.M. Camacho, and N. Martínez-Navarrete. 2011. Changes in flavonoid content of grapefruit juice caused by thermal treatment and storage. *Innov. Food Sci. Emerg. Technol.* 12, 153-162. Doi: 10.1016/j.ifset.2010.12.010
- Kopjar, M., V. Piližota, J. Hribar, M. Simčič, E. Zlatič, and N.N. Tiban. 2008. Influence of trehalose addition and storage conditions on the quality of strawberry cream filling. *J. Food Eng.* 87, 341-350. Doi: 10.1016/j.jfoodeng.2007.12.011
- Kopjar, M., N.N. Tiban, V. Piližota, and J. Babic. 2009. Stability of anthocyanins, phenols and free radical scavenging activity through sugar addition during frozen storage of blackberries. *J. Food Process. Preserv.* 33(Suppl. 1), 1-11. Doi: 10.1111/j.1745-4549.2008.00244.x
- Kopjar, M., K. Jakšić, and V. Piližota. 2012. Influence of sugars and chlorogenic acid addition on anthocyanin content, antioxidant activity and color of blackberry juice during storage. *J. Food Process. Preserv.* 36, 545-552. Doi: 10.1111/j.1745-4549.2011.00631.x
- Ledeker, C.N., S. Suwonsichon, D.H. Chambers, and K. Adhikari. 2014. Comparison of sensory attributes in fresh mangoes and heat-treated mango purées prepared from Thai cultivars. *LWT - Food Sci. Tech.* 56, 138-144. Doi: 10.1016/j.lwt.2013.11.011
- Lončarić, A., K. Dugalić, I. Mihaljević, L. Jakobek, and V. Piližota. 2014. Effects of sugar addition on total polyphenol content and antioxidant activity of frozen and freeze-dried apple purée. *J. Agric. Food Chem.* 62, 1674-1682. Doi: 10.1021/jf405003u
- Maqsood, S. and S. Benjakul. 2010. Comparative studies of four different phenolic compounds on *in vitro* antioxidative activity and the preventive effect on lipid oxidation of fish oil emulsion and fish mince. *Food Chem.* 119, 123-132. Doi: 10.1016/j.foodchem.2009.06.004

- Millán, E.R., L.P. Restrepo, and C.-E. Narváez. 2007. Efecto del escalado, de la velocidad de congelación y de descongelación sobre la calidad de la pulpa congelada de arazá (*Eugenia stipitata* Mc Vaugh). Agron. Colomb. 25, 333-338.
- Narváez-Cuenca, C.-E. 2003. Efecto del choque térmico de arazá (*Eugenia stipitata* Mc Vaugh) sobre la tolerancia al frío. Rev. Colomb. Quím. 32, 93-102.
- Oszmiański, J., A. Wojdyło, and J. Kolniak. 2009. Effect of L-ascorbic acid, sugar, pectin and freeze-thaw treatment on polyphenol content of frozen strawberries. LWT - Food Sci. Tech. 42, 581-586. Doi: 10.1016/j.lwt.2008.07.009
- Re, R., N. Pellegrini, A. Proteggente, A. Pannala, M. Yang, and C. Rice-Evans. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radical Biol. Med. 26, 1231-1237. Doi: 10.1016/S0891-5849(98)00315-3
- Rincon, A. and W.L. Kerr. 2010. Influence of osmotic dehydration, ripeness and frozen storage on physicochemical properties of mango. J. Food Process. Preserv. 34, 887-903. Doi: 10.1111/j.1745-4549.2009.00404.x
- Roos, Y. 1995. Phase transitions in foods. Academic Press, New York, NY.
- Sánchez-Moreno, C., J.A. Larrauri, and F. Saura-Calixto. 1998. A procedure to measure the antiradical efficiency of polyphenols. J. Sci. Food Agric. 76, 270-276. Doi: 10.1002/(SICI)1097-0010(199802)76:2%3c270::AID-JSFA945%3e3.0.CO;2-9; 10.1002/(SICI)1097-0010(199802)76:2<270::AID-JSFA945>3.3.CO;2-0
- Silva, B.K.J., C.-E. Narváez-Cuenca, and L.-P. Restrepo-Sánchez. 2011. Effectiveness of sucrose during the frozen storage of arazá (*Eugenia stipitata* McVaugh) pulp. Agron. Colomb. 29, 441-445.
- Sogi, D.S., M. Siddiq, S. Roidoung, and K.D. Dolan. 2012. Total phenolics, carotenoids, ascorbic acid, and antioxidant properties of fresh-cut mango (*Mangifera indica* L., cv. Tommy Atkin) as affected by infrared heat treatment. J. Food Sci. 77, C1197-C1202. Doi: 10.1111/j.1750-3841.2012.02933.x
- Tsai, P.J., L. Delva, T.Y. Yu, Y.T. Huang, and L. Defossé. 2005. Effect of sucrose on the anthocyanin and antioxidant capacity of mulberry extract during high temperature heating. Food Res. Int. 38, 1059-1065. Doi: 10.1016/j.foodres.2005.03.017
- Vasco, C., J. Ruales, and A. Kamal-Eldin. 2008. Total phenolic compounds and antioxidant capacities of major fruits from Ecuador. Food Chem. 111, 816-823. Doi: 10.1016/j.foodchem.2008.04.054
- Velioglu, Y.S., G. Mazza, L. Gao, and B.D. Oomah. 1998. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. J. Agric. Food Chem. 46, 4113-4117. Doi: 10.1021/jf9801973
- Yen, G.-C. and H.-T. Lin. 1996. Comparison of high pressure treatment and thermal pasteurization effects on the quality and shelf life of guava puree. Int. J. Food Sci. Tech. 31, 205-213. Doi: 10.1111/j.1365-2621.1996.331-32.x