



Agronomía Colombiana

ISSN: 0120-9965

agrocolfabog@gmail.com

Universidad Nacional de Colombia
Colombia

Vergara, Manuel Fernando; Vargas, Jessica; Acuña, John Fabio
Physicochemical characteristics of blackberry (*Rubus glaucus* Benth.) fruits from four
production zones of Cundinamarca, Colombia
Agronomía Colombiana, vol. 34, núm. 3, septiembre-diciembre, 2016, pp. 336-345
Universidad Nacional de Colombia
Bogotá, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=180350123005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Physicochemical characteristics of blackberry (*Rubus glaucus* Benth.) fruits from four production zones of Cundinamarca, Colombia

Características físicoquímicas de frutos de mora de Castilla (*Rubus glaucus* Benth.) provenientes de cuatro zonas productoras de Cundinamarca, Colombia

Manuel Fernando Vergara^{1,2}, Jessica Vargas¹, and John Fabio Acuña¹

ABSTRACT

Blackberries play an important role in human nutrition, due to the elevated content of certain bioactive compounds including ascorbate, anthocyanins, phenolic acids, carbohydrates, and proteins. Blackberry *Rubus glaucus* from Colombian Andes has a high demand on regional and international markets and there is much interest in its cultivation to increase the commercial offer. In four blackberry producing municipalities of Cundinamarca province, fruits were harvested for different quality tests based on the availability of hydric resources in the zone and their influence on fruit characteristics according to the precipitation of each zone, since the crops did not have an irrigation system. The variables, such as size, fresh weight, texture, pH, titratable total acids (TTA), total soluble solids (TSS), and color were evaluated to compare the quality of the fruits between the municipalities, finding that fruits from Silvania had the greater size but, in turn, were the least consistent. El Colegio zone presented the sweetest fruits and together with Pasca had the highest acidity percentage. The color of Pasca and Silvania fruits was the most attractive for commercialization parameters. These parameters, even when management and agro-climatic conditions between municipalities were very similar, allows knowing the positive attributes and determine the factors that need to be improved and to project on a short-term future productions acceptable on the international market.

Key words: drought, water deficiency, growth, development, Rosaceae, post-harvest.

RESUMEN

La mora desempeña un papel importante en la nutrición humana, debido a su elevado contenido en bioactivos que incluyen ascorbato, antocianinas, ácidos fenólicos, carbohidratos y proteínas. La especie *Rubus glaucus* proveniente de los Andes colombianos, tiene alta demanda en el mercado regional e internacional y existe elevado interés en su cultivo para aumentar la oferta comercial. En cuatro diferentes municipios, altamente productores del departamento de Cundinamarca, se cosecharon frutos para realizar diferentes pruebas de calidad con base en la disponibilidad del recurso hídrico en la zona y como este afecta las cualidades de los mismos según la precipitación presentada en cada zona, debido a que los cultivos no cuentan con un sistema de riego disponible. Variables como tamaño, peso fresco, textura, pH, acidéz total titulable (ATT), sólidos solubles totales (SST) y color fueron evaluadas para comparar la calidad de los frutos entre municipios, encontrando que en “Silvania” se produjeron los frutos de mayor tamaño pero a su vez son los menos consistentes. La zona de “El Colegio” presentó los frutos mas dulces y junto con “Pasca” tuvieron el mayor porcentaje de acidez, finalmente el color de frutos de Pasca y Silvania es el mas atractivo para su comercialización, esto, aún cuando el manejo y las condiciones agro-climáticas son muy similares, permite conocer las cualidades positivas y determinar los factores que hacen falta para mejorar y proyectar en un futuro a corto plazo con producciones que tengan aceptación en el mercado internacional.

Palabras clave: sequía, deficit hídrico, crecimiento, desarrollo, Rosaceae, poscosecha.

Introduction

In Colombia, the Andean blackberry (*Rubus glaucus* Benth.), also known as the Castilla blackberry, is a traditional rural crop grown by small producers but with a high production potential (Moreno-Medina *et al.*, 2016), having an area of 12,281 ha, a production of 105,218 t year⁻¹ and an average productivity of 8.6 t ha⁻¹ (Carvalho and Betancur,

2015). Cundinamarca (3,275 ha, 24,239 t year⁻¹ and 7.4 t ha⁻¹), Santander (1,793 ha, 19,190 t year⁻¹ and 10.7 t ha⁻¹) and Antioquia (1,430 ha, 13,728 t year⁻¹, 9.6 t ha⁻¹) are the provinces best positioned in terms of area and production, but Risaralda (14.0 t ha⁻¹) and Santander provinces presented the best performance (Carvalho and Betancur, 2015). The plants are sown in relatively cold areas at an altitude between 1,800 and 2,400 m a.s.l., with 70-80% relative air

Received for publication: 17 October, 2016. Accepted for publication: 30 November, 2016.

Doi: 10.15446/agron.colomb.v34n3.62755

¹ Grupo de Gestión en Tecnología, Innovación y Diseño Agroindustrial (GTI), Departamento de Ingeniería Civil y Agrícola, Facultad de Ingeniería, Universidad Nacional de Colombia. Bogotá (Colombia).

² Corresponding author. mfvergaras@unal.edu.co



humidity, 11-18°C air temperature, 1,500-2,500 mm year precipitation, direct solar radiation between 1,200 and 1,600 h year⁻¹, and an average soil pH of 5.5 (Franco, 2001).

The genus *Rubus* is one of the most diverse in morphological and genetic terms and presents a wide range of wild and cultivated species that are appreciated for the edible fruits (Jennings, 1988). This crop enjoys high consumption in Colombia (1.5 kg year⁻¹ per capita in 2013), with wide consumption acceptance as fresh produce (60%), and is highly regarded in the processed industry (40%) due to its exotic flavor and easy agroindustrial processing (Escobar, 2014).

Berries have been recognized to play an important role in human nutrition providing health-benefits against a wide range of diseases, mainly due to their elevated content in certain bioactive compounds, including ascorbate, anthocyanins, phenolic acids, and carotenoids (Manganaris *et al.*, 2013). Therefore, most bioactive compounds within the plants are secondary metabolites, whose synthesis can be triggered in response to biotic and abiotic stresses, such as UV radiation, drought, wounding as well as infections (Terry *et al.*, 2007; Jahangir *et al.*, 2009).

The blackberry crop is highly sensitive to different issues, for example droughts (Morales and Villegas, 2012) and global climate projections suggest that the frequency of drought events in many regions would increase affecting drought-sensitive plant species. In non-tropical areas, for example, it is expected that water restriction will limit perennial species production in the arid and semi-arid production regions, unless enough water will be stored for irrigation (Walthall *et al.*, 2012). Short-term droughts are projected to occur as frequently as once per year, while occasional long-term droughts (> 6 months) are being projected, where perennial horticulture crops will be highly affected (Walthall *et al.*, 2012). Otherwise, in the United Kingdom, the climate change effects can have a negative impact on berry crops, affecting fruit quality and productivity, and among adaptation strategies to deal with water shortage of these crops are: irrigation scheduling, deficit irrigation techniques and other techniques to reduce water inputs maintaining yield and high quality of fruits (Else and Atkinson, 2010).

High-quality supported by irrigation is an important feature to ensure postharvest life to cope commercialization and transformation processes, because the blackberry is a non-climacteric fruit which has a fragile skin, susceptible to contamination by microorganisms and a very short postharvest life between 3 to 5 d (Ramirez *et al.*, 2013; Sora *et al.*, 2016). During the ripening process a fruit

undergoes continuous physicochemical changes that affect acceptability, quality, and storage time (Ayala *et al.*, 2013) reducing the content of organic acids, allowing loss of firmness and increases the concentration of sugars, among others (Moreno and Deaquiz, 2016). In addition, physical factors, such as color and its uniformity, are parameters that directly define the quality of the fruits, because it is considered to interfere with features, such as flavor and intensity (Moreno and Deaquiz, 2016).

Postharvest studies with fruits of this species seek appropriate handling to ensure the quality and marketing requirements, considering the physical and chemical changes that occur during the maturation, depending on various factors such as light, temperature, moisture, and soil fertility among others (Gómez-Romero *et al.*, 2010).

Hence, under the framework of the “Corredor Tecnológico Agroindustrial II - Fresa y Mora” project, it was considered important to characterize the physical and chemical properties of fruits based on the hydric status of plants and harvested fruits in four different high blackberry producing municipalities in Cundinamarca, Colombia. This, in order to know its performance and to generate short-term improvements in such products that can be projected as a producing region and improve both the quality of life and the recognition of farmers based on an international market approach.

Materials and methods

Location

The study was conducted in four representative producing areas of the province of Sumapaz, Cundinamarca, that refer to different municipalities (Tab. 1). These zones were strategically chosen because of their agroclimatic potential that promotes better plant growth and fruit development and have a long tradition of blackberry growing.

To carry out the study, a representative sample of fruits was collected per municipality, and classified by three maturity stages (4, 5 and 6) after the Colombian Technical Standard (NTC 4106; Icontec, 1997), where fruits from each state were taken for variables evaluation. The tests were performed 1 d after harvest in the postharvest laboratory of the Agricultural Engineering Department of the Universidad Nacional de Colombia, Bogotá.

Environmental data

Recorded rainfall data (in mm), accumulated over a period of 1 year (April 2015 to April 2016), were obtained from

TABLE 1. Climatic conditions of each production site.

Municipality	Crop location	Altitude (m a.s.l.)	Average day temperature (°C)	Relative air humidity (%)
El Colegio	4°32'29.9"N 74°24'28.01"W	1,882	17	75
Pasca	4°14'10"N 74°19'31.5"W	2,647	13	80
San Bernardo	4°6'4.57"N 74°24'52.85"W	2,410	15	84
Silvania	4°27'33.4"N 74°19'25.6"W	2,200	16	80

the database of the “Instituto de Hidrología, Meteorología y Estudios Ambientales (IDEAM)” from the Ministry of “Ambiente y Desarrollo Sustentable” of Colombia, whereas the relative air humidity and average day temperature were recorded by dataloggers (RHT10, Extech Instruments®, Waltham, MA) located in each location.

Fresh and dry mass of fruits

In order to obtain dry mass of fruits, three samples were taken from three fruits at each maturity stage, the samples were weighed and later dried in oven at 70 °C with measurements every 24 h until obtain a constant weight, having its percentages using the following formula:

$$\text{Dry mass (\%)} = \frac{\text{initial weight}}{\text{final weight}} \times 100 \quad (1)$$

Texture

To determine texture from the side to the center of the fruit, a CT3 texturometer (Brookfield Engineering Laboratories, Middleborough, MA) was used with a punch accessory diameter of 0.2 cm, with reference the maximum load and final charge applied in grams (g), whereas deformation was obtained in millimeters (mm).

Total titratable acidity (TTA)

TTA was determined with potentiometer TitroLine® WA 20 (SI Analytics, Weilheim, Germany), taking 10 g of pure fruit extract plus 10 g of distilled water to obtain a 1:1 solution. The computer showed the pH and the volume of reagent required for titration in millimeters (mm), so that the percentage of acidity (malic acid for blackberry fruits) was calculated by the following formula:

$$\text{Acidity (\%)} = \frac{V \times N \times C}{W} \times 100 \quad (2)$$

Where

V: NaOH used volume (mL)

N: NaOH normality (0.1 N)

C: malic acid constant (0.067 g meq⁻¹)

W: sample weight (g)

Total soluble solids (TSS)

For TSS determination, a J357 refractometer (Rudolph Research Analytical, Hackettstown, NJ) was used with one pure extract juice of blackberry as sample and the results were expressed in percentage units. Additionally, the reading was corrected for acidity using the formula obtained by Herrera (2010):

$$\text{Corrected TSS} = \text{TSS (\%)} + \text{acid (\%)} \times 0.0118 \quad (3)$$

Color

For color measurement the system of Hunter *L a b* and a chroma meter CR 400 (Konica Minolta Sensing Europe, Bremen, Germany) was used. After its calibration, the readings were taken at three different sites of each fruit.

Statistical analysis

All statistical analysis was performed with R software (v 2.15.2) (Development Core Team 2012). Data were reviewed using a mixed lineal effect model, with Tukey test for variance and homogeneity with a confidence level of $\alpha = 0.05$ to identify significant differences among the variables evaluated.

Results and discussion


Water availability

In order to record the average rainwater per month, from april 2015 to april 2016, the rain water amounts in each evaluated zone were registered from Ideam wheater stations (Tab. 2).

Figure 1 shows the amount of rainfeed water in each municipality and, as an average, the dotted line of the base requirement for a blackberry crop to fulfill its normal physiological functions in the conditions of the province of Cundinamarca. Although the crops are located in different sites, the characteristics such as soil, management, and its structures are very similar among them. For optimal growth and development of blackberry crops, it is necessary to supply about 30 mm of water per week (4.28 mm d⁻¹) (Casaca, 2005). A drip irrigation system is recommended

TABLE 2. Locations and the weather stations close to each experimental farm (Ideam, 2016).

Municipality	Location	Station name	Location
El Colegio	4°32'29.9"N 74°24'28.01"W	Hidroparaiso	4°57'31.6"N 74°40'48.33"W
Pasca	4°14'10"N 74°19'31.5"W	Pasca	4°31'01"N 73°31'17.5"W
San Bernardo	4°6'4.57"N 74°24'52.85"W	El Tulcan	4°1'5.24"N 74°35'97.55"W
Silvania	4°27'33.4"N 74°19'25.6"W	Alto San Miguel	4°44'96.6"N 74°29'97.2"W

**FIGURE 1.** Daily rainfall average per month in each experimental blackberry farm on a one year period (april 2015 to april 2016), after Ideam (2016).

by Casaca (2005) because it adapts to any topographical condition and there is no contact between the water and the aerial part of plants and, thus, takes full advantage of water resource. Flowering and fruit growth are the main periods to irrigate the crop, because constant irrigation increases performance, increasing fruit size and quantity of fruits per plant (Fischer and Orduz-Rodríguez, 2012).

Regardless of climatic change alterations presented throughout the year as El Niño and La Niña phenomena, it is evident that rain water supply in nearly all months (Fig. 1) was not enough to cover the minimum need amount for all fruit development. After Jaleel *et al.* (2009) during a prolonged water deficit, the overall plant development is delayed and leaf size is reduced; inducing anatomical changes due to modifications in cell size and a premature senescence and, ultimately, plant death observed in several species.

For adaptation strategies, it is important to increase the irrigation efficiency implementing different irrigation methods, such as drip irrigation or the partial root-zone drying technique. In other species, partial “root-zone drying” can allow a 29-50% of water saving with a similar yield and a concomitant water use efficiency increase (Monneveux *et*

al., 2013). Water restriction during full flowering and fruit production is particularly harmful. A study showed that blackberry plants had decreased stomatal conductance and transpiration after 24 d under dry-out conditions (drought stress), while the leaf turgor did not show variation (Parra *et al.*, 1999). This result suggests the occurrence of osmotic adjustment activity, in fact, blackberry plants without irrigation presented double proline accumulation compared with plants under irrigation (Parra *et al.*, 1999).

Fruit size

Fruits size, expressed in volume, among municipalities and by maturity stage (with 6,32 cm³; 6,92 cm³ and 6,47 cm³ for maturity stages 4, 5 and 6, respectively) showed slightly significant differences due to environmental conditions of the four producing areas (Fig. 2). Probably, because the agronomic management applied to each of them (foliar and soil fertilization, pesticide application, spalier, weed management, pruning and harvesting) were very similar, so the homogeneity can be attributed to these factors. Studies done by Ayala *et al.* (2013) in blackberry in the province of Tolima showed an average fruit volume for maturity stages 4, 5 and 6 of 6,81 cm³; 7,55 cm³ and 7,49 cm³, respectively, with values higher than in our study, where the agronomic management was based on a traditional knowledge.

TABLE 3. Fresh weight, dry biomass, and water content of blackberry fruits per maturity stage (4, 5 and 6) by each site and 6) by each site.

Municipality	Fresh weight (g)			Biomass (g)			Water content (g)		
	4	5	6	4	5	6	4	5	6
El Colegio	16.87 ab	19.64 ab	15.83 ab	2.18 a	2.45 a	2.12 a	14.69 a	17.19 a	13.7 a
Pasca	18.06 a	19.96 a	21.86 a	2.23 a	2.37 a	2.36 a	15.83 a	17.59 a	19.5 a
San Bernardo	19.54 ab	19.24 ab	16.62 ab	2.46 a	2.55 a	2.03 a	17.08 a	16.69 a	14.59 a
Silvania	15.3 b	15.13 b	15.3 b	1.94 b	1.94 b	1.59 b	13.36 b	13.19 b	13.71 b




FIGURE 2. Average volume of blackberry fruits harvested in four municipalities and classified at three different maturity stages. Means with different letters indicate significant differences according to the Tukey test ($P \leq 0.05$). Error bars indicate standard error.

Previous studies by García (2012) in Silvania municipality reported a harvested fruit average fruit length of 31.2 mm during the rainy season, while in this study an average length of 26.5 mm was obtained (data not shown), including also the municipality of Silvania. The low weight and size of blackberry fruits in our study, among other agronomic management factors, could be affected most by the lack of water supply. Fischer and Orduz-Rodríguez (2012) indicated that water deficiency can reduce the growth and quality of fruits, which normally contains in its maturity stage up to 90% of water and, especially, during the fruit enlargement phase a constant water supply is the basis for its normal growth. Hsiao *et al.* (1976) stated that one of the first responses to a water deficit is reduced plant growth due to cell expansion inhibition.

Fresh and dry matter of fruits

The fresh weight, dry biomass, and water content by stage of maturity is shown in Tab. 3. No significant differences were registered between municipalities demonstrating very consistent results. Weight variation was not determined directly by the biomass but more by the amount of water contained in cells and rainfall of the four studied

zones were more or less similar (IDEAM, 2016). Vicente *et al.* (2009) corroborate that water is the most abundant component in fruits and vegetables comprising about 90% of their mass. The water content values vary according to the product structure, what is the reason that fruits and vegetables are highly perishable, especially when having a high water content that can easily allow growing pathogen microorganisms (Fischer, 2009).

With the formation of fruits an increase in their density occurs, being, in many cases, directly proportional to its weight, as a result of internal transformations through the expansion of cells and tissues (Quimbaya, 2009). However, because of the low water availability in soil, which decreases photosynthesis, carbohydrate accumulation and, finally, overall plant growth are limited (Chaves and Oliveira, 2004). In addition, drought stress causes stomata closure, decreases gas exchange by reducing transpiration and, thus, the photosynthetic rate (Glass *et al.*, 2003; Kiziloglu *et al.*, 2006), all of them affecting density of cells.

Texture of fruits

Figure 3 shows that consistency and firmness of fruits were reduced as a consequence of maturation by the decrease of osmotic cellular pressure through chemical reactions in the processes of degradation of pectin, starch, and cellulose (Quimbaya, 2009), as well as by reducing its fiber content and from degradative processes of cell walls through which fruits become softer and more susceptible to physical damages during postharvest handling (Arias and Toledo, 2000).

According to data reported by Farinango (2010) for blackberry *Castilla* fruits, resistance to penetration varies from 0.23 kgf to 0.48 kgf for maturity stages 2 to 5 but, in our study, in El Colegio, Pasca and San Bernardo municipalities, the values between 0.21 and 0.45 kgf were obtained (data not shown). Although there were similar results, the municipality of Silvania presented a significant drop in the consistency and strength (Fig. 3), which can be attributed, primarily, to the employment of a blackberry hybrid between Castilla and San Antonio variety, and this intercrossing presented an improved performance, but their




FIGURE 3. A. Consistency and B. Firmness of blackberry fruits per maturity stage and municipality.

physical characteristics werenot the most suitable to obtain favorable postharvest results.

Secondly, the municipality of Silvania received low amounts of rainwater throughout the year (Fig. 1), this could have affected the process of overall production of plants by weakening the cell walls and the cohesive forces that hold together the cells between them, generating a disintegration of cell structure, besides the internal movement of water related to osmosis processes (Angón-Galván *et al.*, 2006). Also, in soil analysis from Silvania. it was found that Ca levels (data not shown) were very low (2.64 cmol kg⁻¹ exchangeable Ca) compared with the other municipalities, which have acceptable levels (Pasca 5.04 cmol kg⁻¹ and San Bernardo 4.01 cmol kg⁻¹). According to Castaño *et al.* (2010), calcium reinforces the cell wall and tissues of plants, calcium pectate gives resistance to infection by fungi or bacteria, increases the hardness of the fruits and delays ripening, consequently giving stability to the cell membrane, and favors the cation and anion balance within the cell and osmoregulation.

Titrateable total acids (TTA) and total soluble solids (TSS) of fruits

The behavior of TTA showed that, as ripening occurred, acidity decreased (Fig. 4A); similar results were obtained by Díaz *et al.* (2011), who claimed that the acids contained in fruits are consumed in respiration across the time, giving rise to new compounds (Kays, 2004), which explains its decline during the ripening process. However, concentrations of TTA, as shown in figure 4A, were considerably higher compared with those in NTC 4106 (Icontec, 1997), where the contents at maturity stages 4, 5, and 6 corresponded to 3.0, 2.8, and 2.5%, respectively.

The municipality of San Bernardo had the lowest levels differing significantly ($P \leq 0.05$) from the others based on their maturation process. The studies by Moreno and Deaquiz (2016) in *Rubus alpinus* showed concentrations of 2.28 and 1.85% at stages 5 and 6 TTA, respectively, indicating these high levels can be viewed positively for accessing a world-class market. Clark and Finn (2008) indicated that the United States industry prefers an original blackberry flavor, good color, low pH, high content of soluble solids and acidity. and low number of seeds, while, for consumers, the qualities, such as sweetness, acidity, astringency, color, firmness, and absence of seeds are important for both processed and fresh fruits.

It is worthwhile to mention that TSS measurement is associated with dissolved sugars in the cell sap (Osterloh *et al.*, 1996), and these authors claimed that the amount of sugars in fruit would be determined by the variety, the assimilatory performance of leaves, the relationship between leaf and fruit, and the climatic conditions during fruit development, among others.

However, the results found in our study (Fig. 4B) can be compared with other blackberry studies; for example, Grijalba *et al.* (2010) recorded an average of 5.69 °Brix at maturity stage 4, values lower than those obtained by other authors, such as Vásquez *et al.* (2006), who recorded values between 7.7 and 9.1 °Brix for *Castilla* blackberry, whereas Tosun *et al.* (2008) reported values of 7-11 °Brix of blackberries in Brazil. Ali *et al.* (2011) explained that variations may occur in the chemical content in response to the location of the crop, especially due to differences in temperature ranges, which influence the organoleptic characteristics of fruits (Osterloh *et al.*, 1996). Therefore, lower TSS results obtained from the municipality of Pasca




FIGURE 4. A. Total titratable acidity and B. Total soluble solids (°Brix), obtained by three different maturity stages compared by each municipality. Means with different letters indicate significant differences according to the Tukey test ($P \leq 0.05$).

were normal because of its high elevation site and, consequently, a lower average temperature than from the other municipalities (Fischer and Orduz-Rodríguez, 2012).

Color

The color is an important parameter to assess as harvest index during the ripening of many fruits. The NTC 4106

standard (Icontec, 1997) provides a visual color scale with six stages of maturity for blackberry fruits along with images and descriptions for each maturity stage, which were established using subjective evaluations.

Figure 5 shows the behavior of the parameters L , a , and b , based on the Hunter Lab scale in the four evaluated




FIGURE 5. Hunter Lab graphics obtained by harvested and evaluated fruits selected by maturity stage (4: dashed, 5: grey, and 6: black) in the four municipalities. A. El Colegio, B. Pasca, C. San Bernardo, and D. Silvania.

municipalities. The *L* component represents the variation between black and white, where black = 100 and white = 0. *a* parameter does not have a numerical limit defined and its positive value indicates a trend towards the red hue, while its negative value indicates a trend toward green. The parameter *b* also has a definite numerical limit and its positive value indicates a trend toward the yellow hue, while its negative value indicates a tendency towards blue tone (Hunterlab, 2012).

A clear decrease of the *L* as maturity progresses was observed (Fig. 5) indicated the trend that fruit acquire to become darker, just as the value *b* is abruptly reduced showing the trend that takes the fruit to the blue color. Finally, figure 5 shows that there were no negative values eliminating the green color of the fruit structure, however, it indicated a positive decrease placing the fruit in an increasingly clear red hue. This reduction is explained by the change to purple red hue of the fruits during ripening due to the appearance of pigments, such as anthocyanins (García, 2012).

The trend of color change of the harvested fruit as the same, however significant differences in the intensity of the colors were evident (Fig. 5). Blackberries from Sylvania were the most attractive ones because of their dark color, which may be through the *Castilla* and *San Antonio* combination or can also be a reaction of hydric stress in these zones unlike the other municipalities. Terry *et al.* (2007) affirmed, for strawberry plants, that irrigation shortage resulted in berries with higher concentrations of anthocyanins and antioxidant capacity.

Conclusions

Currently it is not accurate to recommend an optimal development of crop production based just on rainwater supply, even if Colombia presents a bimodal rainfall system. It was evident that the one year period of rainwater received failed to meet the basic needs, not even for the projected yield spectations.

It is highly recommended to implement a drip irrigation system, which is friendly to water use without generating unexpected cost increases and ensures the plant health.

For blackberry crops in Cundinamarca, simple adjustments in fertilization plans can contribute to a significantly improved both physical and biochemical quality of fruits.

A high percentage of titratable acidity and low concentrations of total soluble solids are observed, this is positive

but still is difficult to project these blackberries to an international market, basically due to the size of the fruit. Therefore, it is necessary to have scientific assistance focused on fruit quality along the time which allows to do the correct improve at the right time.

It was determined that the fruits from Sylvania had the largest size, those from El Colegio were the sweetest ones and those from Pasca had the highest acidity and, finally, the most attractive color were found in fruits of Pasca and Sylvania. So, each municipality has interesting characteristics for marketing. Now, the next step is to homogenize and improve with research and constant accompaniment the qualities of blackberry fruits for the farmers of Colombia.

Acknowledgments

The authors thank the Corredor Tecnológico Agroindustrial II – Fresa y Mora project, for allowing the realization of this study with the help and willingness of farmers in each area, under the direction of Dr. Martha Bolaños from blackberry sub-project, also thank to Mrs. Elizabeth Amado, Mr. Julian Carrasco and Mr. Brian Albarracin for their support in obtaining data in the laboratory and to Mrs. Giovanna Quintero for her contributions.

Literature cited

- Ali, L., B. Svensson, B.W. Alsanius, and M.E. Olsson. 2011. Late season harvest and storage of *Rubus* berries - Major antioxidant and sugar levels. *Sci. Hortic.* 129, 376-381. Doi: 10.1016/j.scienta.2011.03.047
- Angón-Galván, P., N.F. Santos-Sanchez, and C.G. Hernández. 2006. Índices para la determinación de las condiciones óptimas para la maduración de un fruto. *Temas Cienc. Tecnol.* 10(30), 3-8.
- Arias, C.J. and J. Toledo. 2000. Manual de manejo poscosecha de frutas tropicales (papaya, piña, plátano, cítricos). FAO, Rome, Italy.
- Ayala, L.C., C.P. Valenzuela, and Y. Bohorquez. 2013. Caracterización fisicoquímica de mora de castilla (*Rubus glaucus* Benth.) en seis estados de madurez. *Biotechnol. Sector Agropecu. Agroind.* 11(2), 10-18.
- Carvalho, C.P. and J.A. Betancur. 2015. Quality characterization of Andean blackberry fruits (*Rubus glaucus* Benth.) in different maturity stages in Antioquia, Colombia. *Agron. Colomb.* 33(1), 74-83. Doi: 10.15446/agron.colomb.v33n1.47132
- Casaca, Á.D. 2005. Guía técnica de frutas y vegetales - El cultivo de la mora. Proyecto de Modernización de los Servicios de Tecnología Agrícola - PROMOSTA. San José de Costa Rica.
- Castaño, C.A., C.L. Morales, and F.H. Obando. 2010. Evaluación de las deficiencias nutricionales en el cultivo de mora (*Rubus glaucus* Benth.) en condiciones controladas para bosque montano bajo. *Agron* 16(1), 75-88.
- Chaves, M. and M. Oliveira. 2004. Mechanisms underlying plant resilience to water deficits: prospects for water-saving agriculture. *J. Exp. Bot.* 55, 2365-2384. Doi: 10.1093/jxb/erh269

- Clark, J.R. and C.E. Finn. 2008. New trends in blackberry breeding. *Acta Hortic.* 777, 41-47. Doi: 10.17660/ActaHortic.2008.777.2
- Díaz, R.O., L. Moreno, R. Pinilla, W. Carillo, L.M. Melgarejo, O. Martínez, J.P. Fernández-Trujillo, and M.S. Hernández (2011). Postharvest behavior of purple passion fruit in Xtend® bags during low temperature storage. *Acta Hortic.* 934, 727-731. Doi: 10.17660/ActaHortic.2012.934.95
- Else, M. and C. Atkinson. 2010. Climate change impacts on UK top and soft fruit production, *Outlook Agric.* 39, 257-262. Doi: 10.5367/oa.2010.0014
- Escobar, C. 2014. Cadena productiva nacional de la mora. Indicadores de apoyo. Ministerio de Agricultura y Desarrollo Rural, Bogota, Colombia
- Farinango, M.E. 2010. Estudio la fisiología poscosecha de la mora de castilla (*Rubus glaucus* Benth.) y de la mora variedad brazos (*Rubus* sp.). Facultad de Ingeniería Química y Agroindustria, Escuela Politécnica Nacional, Quito, Ecuador.
- Fischer, G. 2009. Efecto del agua en la calidad del fruto. pp. 27-32. In: Robayo, G. (ed.). Agua: Conservación, educación, agua potable. Riego y nutrición en semillas y cultivos. Galrobayov Editores Científicos-Técnicos, Bogota, Colombia.
- Fischer, G. and J.O. Orduz-Rodríguez. 2012. Ecofisiología en frutales. pp. 54-72. In: Fischer, G. (ed.). Manual para el cultivo de frutales en el trópico Produmedios, Bogotá, Colombia.
- Franco, G. 2001. Fertilización de la mora de Castilla (*Rubus glaucus* Benth.) en zonas del departamento de Caldas. In: Proc. Tercer seminario de clima frío moderado. Manizales, Colombia.
- García, G. 2012. Elaboración de un paquete tecnológico para productores, en manejo, cosecha y poscosecha de mora (*Rubus glaucus* Benth.) aplicando ingeniería de calidad y determinación de las características nutraceuticas de la fruta en precosecha, en el municipio de Silvania Cundinamarca. MSc. thesis. Universidad Nacional de Colombia, Bogota, Colombia.
- Glass, V. D. Percival, and J. Proctor. 2003. Influence of decreasing soil moisture on stem water potential, transpiration rate and carbon exchange rate of the lowbush blueberry (*Vaccinium angustifolium* Ait) in a controlled environment. *J. Hortic. Sci. Biotechnol.* 78(3), 359-364. Doi: 10.1080/14620316.2003.11511632
- Gómez-Romero, M., A. Segura-Carretero, and A. Fernández-Gutiérrez. 2010. Metabolite profiling and quantification of phenolic compounds in methanol extracts of tomato fruit. *Phytochem.* 71(16), 1848-1864. Doi: 10.1016/j.phytochem.2010.08.002
- Grijalba, C.M., L.A. Acalderon, and M.M. Pérez. 2010. Rendimiento y calidad de la fruta en mora de castilla (*Rubus glaucus* Benth.) con y sin espinas, cultivado a campo abierto en Cajicá - Cundinamarca, Colombia. *Rev. Fac. Cienc. Básicas* 6(1), 24-41.
- Hsiao, T., E. Acevedo, E. Fereres, and D. Henderson. 1976. Water stress, growth, and osmotic adjustment. *Phil. Trans. Roy. Soc. London* 273, 479-500. Doi: 10.1098/rstb.1976.0026
- Hunterlab. 2012. Hunter L, a, b color scale. Applications Note 8(9). In: <http://www.hunterlab.se/wp-content/uploads/2012/11/Hunter-L-a-b.pdf>; consulted: july, 2016.
- Ideam. 2016. Base de datos del Instituto de Hidrología, Meteorología y Estudios Ambientales, Bogota, Colombia.
- Icontec. 1997. NTC 4106. Frutas frescas. Mora de castilla. Especificaciones. Instituto Colombiano de Normas Técnicas y Certificación, Bogota, Colombia.
- Jahangir, M., I.B. Abdel-Farid, H.K. Kim, Y.H. Choi, and R. Verpoorte. 2009. Healthy and unhealthy plants: the effect of stress on the metabolism of brassicaceae. *Environ. Exp. Bot.* 67, 23-33. Doi: 10.1016/j.envexpbot.2009.06.007
- Jaleel, C.A., P. Manivannan, A. Wahid, M. Farooq, R. Somasundaram, and R. Panneerselvam. 2009. Drought stress in plants: a review on morphological characteristics and pigments composition. *Int. J. Agric. Biol.* 11, 100-105.
- Jennings, D. 1988. Raspberries and blackberries: their breeding, diseases and growth. Academic Press, London, U.K.
- Kays, S. 2004. Postharvest biology. Exon Press, Athens, GA.
- Kiziloglu, F.M. U. Sahin, T. Tunc, and S. Diler. 2006. The effect of deficit irrigation on potato evapotranspiration and tuber yield under cool season and semiarid climatic condition. *J. Agron.* 5, 284-288. Doi: 10.3923/ja.2006.284.288
- Manganaris, G.A., V. Goulas, A.R. Vicente, and L.A. Terry. 2013. Berry antioxidants: small fruits providing large benefits. *J. Sci. Food Agric.* 94, 825-833. Doi: 10.1002/jsfa.6432
- Monneveux, P., D.A. Ramírez, and M.T. Pino. 2013. Drought tolerance in potato (*S. tuberosum* L.). Can we learn from drought tolerance research in cereals? *Plant Sci.* 205-206, 76-86. Doi: 10.1016/j.plantsci.2013.01.011
- Morales, C.S. and B. Villegas. 2012. Mora (*Rubus glaucus* Benth.). pp. In: Fischer, G. (ed.). Manual para el cultivo de frutales en el trópico. Produmedios, Bogota, Colombia.
- Moreno, B.L. and Y.A. Deaquiz. 2016. Caracterización de parametros fisicoquímicos en frutos de mora (*Rubus alpinus* Macfad.). *Acta Agron.* 65(2), 130-136. Doi: 10.15446/acag.v65n2.45587
- Moreno-Medina, B.L., F. Casierra-Posada, and Michael Blanke. 2016. Índices de crecimiento en plantas de mora (*Rubus alpinus* Macfad) bajo diferentes sistemas de poda. *Rev. Colomb. Cienc. Hortic.* 10(1), 28-39. Doi: 10.17584/rcch.2016v10i1.4457
- Osterloh, A., G. Ebert, W.H. Held, H. Schulz, and E. Urban. 1996. Lagerung von Obst und Südfrüchten. Verlag Ulmer, Stuttgart, Germany.
- Parra, R.A., J.L. Rodríguez, and V.A. González. 1999. Transpiración, potencial hídrico y prolina en zarzamora bajo déficit hídrico. *Terra* 17, 125-130.
- Quimbaya, A.Y. 2009. Desarrollo de una propuesta para mejorar el manejo poscosecha de los frutos de gulupa (*Passiflora edulis* Sims.) y curuba (*Passiflora mollissima* Bailey). ICTA, Universidad Nacional de Colombia, Bogota, Colombia.
- Development Core Team. 2012. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ramírez, J., I. Aristizabal, and J. Restrepo. 2013. Conservación de mora de castilla mediante la aplicación de un recubrimiento comestible de gel de mucílago de penca de sábila. *Vitae* 20(3), 172-183.
- Sora, A.D., G. Fischer, and R. Flórez. 2006. Almacenamiento refrigerado de frutos de mora de Castilla (*Rubus glaucus* Benth.) en empaques con atmosfera modificada. *Agron. Colomb.* 24(2), 306-316.
- Terry, L.A., G.A. Chope, and J.G. Bordonaba. 2007. Effect of water deficit irrigation and inoculation with *Botrytis cinerea* on

- strawberry (*Fragaria xananassa*) fruit quality. J. Agric. Food Chem., 55, 10812-10819. Doi: 10.1021/jf072101n
- Tosun, I., N.S. Unstun, and B. Tekguler. 2008. Physical and chemical changes during ripening of blackberry fruits. Sci. Agric. 65, 87-90. Doi: 10.1590/S0103-90162008000100012
- Vásquez R., H.H. Ballesteros, C.A. Muñoz, and M.E. Cuellar. 2006. Utilización de la abeja *Apis mellifera* como agente polinizador en cultivos comerciales de fresa (*Fragaria chiloensis*) y mora (*Rubus glaucus*) y su efecto en la producción. Corpoica, Tibaitatá, Mosquera, Colombia.
- Vicente, A.R., G.A. Manganaris, G.O. Sozzi, and C.H. Crisosto. 2009. Postharvest handling: a system approach. Chapter 5. Nutritional quality of fruits and vegetables. Macmillan Publishing, London, U.K.
- Walthall, C.L., J. Hatfield, P. Backlund, L. Lengnick, E. Marshall, M. Walsh, S. Adkins, M. Aillery, E.A. Ainsworth, C. Ammann, C.J. Anderson, I. Bartomeus, L.H. Baumgard, F. Booker, B. Bradley, D.M. Blumenthal, J. Bunce, K. Burkey, S.M. Dabney, J.A. Delgado, J. Dukes, A. Funk, K. Garrett, M. Glenn, D.A. Grantz, D. Goodrich, S. Hu, R.C. Izaurralde, R.A.C. Jones, S-H. Kim, A.D.B. Leaky, K. Lewers, T.L. Mader, A. McClung, J. Morgan, D.J. Muth, M. Nearing, D.M. Oosterhuis, D. Ort, C. Parmesan, W.T. Pettigrew, W. Polley, R. Rader, C. Rice, M. Rivington, E. Rosskopf, W.A. Salas, L.E. Sollenberger, R. Srygley, C. Stöckle, E.S. Takle, D. Timlin, J.W. White, R. Winfree, L. Wright-Morton, and L.H. Ziska. 2012. Climate change and agriculture in the United States: Effects and adaptation. USDA Tech. Bull. 1935. Washington, DC.