



Iatreia

ISSN: 0121-0793

revistaiatreia@udea.edu.co

Universidad de Antioquia

Colombia

Bonilla-Ramírez, Leonardo; Jiménez-Del-Río, Marlene; Veléz-Pardo, Carlos  
Glucose promotes resistance in lymphocytes against oxidative stress-induced apoptosis  
through signaling and metabolic pathways. Implications for Parkinson's disease

Iatreia, vol. 30, núm. 2, abril-junio, 2017, pp. 117-130

Universidad de Antioquia

Medellín, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=180550477002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Glucose promotes resistance in lymphocytes against oxidative stress-induced apoptosis through signaling and metabolic pathways. Implications for Parkinson's disease

Leonardo Bonilla-Ramírez<sup>1</sup>, Marlene Jiménez-Del-Río<sup>1</sup>, Carlos Veléz-Pardo<sup>1</sup>

## SUMMARY

**Introduction:** Parkinson's disease (PD) is a neurological disorder associated with the selective loss of dopaminergic (DAergic) neurons. Clinical data suggest that oxidative stress (OS) and dysregulation of glucose (G) metabolism are early events in PD. However, no data are available to explain the molecular connection between glucose metabolism, OS, and neuronal demise in PD. Human lymphocytes share a similar dopaminergic signaling mechanism with DAergic neurons. Further, rotenone (ROT) is a mitochondrial complex I inhibitor that selectively induces apoptosis through OS in dopaminergic neurons and lymphocytes. Thus, to test the hypothesis that G metabolism and OS are linked in dopaminergic system toxicity and PD, human lymphocytes were cultured with ROT in the presence or absence of various concentrations of glucose.

**Objective:** This study examines the response of human lymphocytes to glucose (11, 55, 166, 277, 555 mM G) in the absence or presence of ROT (250 microM).

**Methods:** Light and fluorescence microscopy and immunocytochemistry techniques were used to evaluate morphological and biochemical changes in human lymphocytes.

**Results:** 55 mM G was effective in suppressing ROT-induced apoptosis in lymphocytes via 5 pathways: (i) pentose phosphate pathway (PPP), (ii) glutathione (GSH) pathway, (iii) superoxide dismutase (SOD) and catalase (CAT) antioxidant systems, and (iv) Phosphoinositide 3-kinase (PI3-K) signaling. Additionally, we report for the first time that G rescued lymphocytes from

---

<sup>1</sup> Neuroscience Research Group, Medical Research Institute, Faculty of Medicine, University of Antioquia (UdeA), calle 70# 52-21, and Calle 62 # 52-59, Building 1, Room 412; SIU, P.O. Box 1226, Medellín, Colombia  
Correspondencia: Carlos Vélez-Pardo; calberto.velez@udea.edu.co

Recibido: mayo 04 de 2016

Aceptado: agosto 12 de 2016

Cómo citar: Bonilla-Ramírez L, Jiménez-del-Río M, Vélez-Pardo C. Glucose promotes resistance in lymphocytes against oxidative stress-induced apoptosis through signaling and metabolic pathways. Implications for Parkinson's disease. *Iatreia*. 2017 Abr-Jun;30(2):117-130. DOI 10.17533/udea.iatreia.v30n2a02

ROT-induced apoptosis by (v) activating NF- $\kappa$ B and down-regulating p53 and caspase-3. Signaling (e.g., LY294002) and metabolic inhibitors of these pathways (e.g., dehydroepiandrosterone (DHEA), L-buthionine-sulfoximine (BSO), 1,3-Bis (2-chloroethyl)-1-nitrosourea (BCNU), mercaptosuccinic acid (MS), 3-amino-1,2,4-triazole (AT), sodium diethyldithiocarbamate (DDC)) only partially reversed the protective effect of 55 mM G on lymphocytes exposed to ROT.

**Conclusion:** These data suggest that high G simultaneously triggers cellular signaling and antioxidant systems to ensure global cell protection against stressful conditions in DAergic cells.

## KEY WORDS

*Apoptosis; Glucose; Lymphocytes; Oxidative stress; Rotenone; Signaling*

## RESUMEN

**La glucosa promueve la resistencia en linfocitos contra el estrés oxidativo que induce apoptosis a través de rutas de señalización y metabólica. Impacto en la enfermedad de Parkinson**

**Introducción:** la enfermedad de Parkinson (EP) es un trastorno neurológico asociado con la pérdida selectiva de neuronas dopaminérgicas (DAérgicas). Datos clínicos sugieren que el estrés oxidativo (EO) y la desregulación del metabolismo de la glucosa (G) son eventos tempranos en la EP. Sin embargo, no existe información que explique la posible asociación molecular entre el metabolismo de la glucosa, el EO y la muerte neuronal. Los linfocitos humanos comparten mecanismos de señalización DAérgicos comunes. Más aún, la rotenona (ROT) es un inhibidor que selectivamente induce apoptosis vía EO en neuronas DAérgicas y linfocitos. Para evaluar la hipótesis que el metabolismo de la G y el EO están asociados con la toxicidad del sistema DAérgico y EP, se cultivaron linfocitos humanos con ROT en presencia o ausencia de varias concentraciones de G.

**Objetivo:** este estudio examina la respuesta de los linfocitos a G (11, 55, 166, 277, 555 mM) en ausencia o presencia de ROT (250  $\mu$ M).

**Métodos:** se utilizaron técnicas de microscopía de luz y fluorescencia e inmunocitoquímica para evaluar los cambios morfológicos y bioquímicos de los linfocitos.

**Resultados:** la G 55 mM fue eficaz en suprimir la apoptosis en linfocitos inducida por ROT vía activación de 5 rutas metabólicas: (i) la vía pentosa fosfato, (ii) la vía glutatión; (iii) los sistemas antioxidantes superóxido dismutasa (SOD) y catalasa (CAT); (iv) fosfoinositol 3 cinasa (PI3-K). Además, se observó por primera vez que la G rescata linfocitos de la apoptosis inducida por ROT vía (v) activación del factor nuclear  $\kappa$ B (NF- $\kappa$ B) y por regulación a la baja de p53 y de la caspasa-3. Se demostró que los inhibidores de señalización (v.gr. LY294002) e inhibidores metabólicos (v.gr. DHEA, BSO, BCNU, MS, DCC) revierten parcialmente los efectos citoprotectores de la G 55 mM en linfocitos expuestos a ROT.

**Conclusión:** estos hallazgos sugieren que la alta concentración de G induce simultáneamente sistemas de señalización y antioxidantes para asegurar la protección global de la célula contra condiciones estresantes en células DAérgicas.

## PALABRAS CLAVE

*Apoptosis; Estrés Oxidativo; Glucosa; Linfocitos; Rotenona; Señalización*

## RESUMO

**A glucose promove a resistência em linfócitos contra o estresse oxidativo que induz apoptose a través de rotas de sinalização e metabólica. Impacto na doença de Parkinson**

**Introdução:** a doença de Parkinson (PD) é uma desordem neurológica associada a perda seletiva de neurônios dopaminérgicos (Daérgicos). Dados clínicos sugerem que o estresse oxidativo (EO) e a desregulação do metabolismo da glicose (G) são eventos iniciais no PD. No entanto, não há nenhuma informação que explica a possível associação molecular entre o metabolismo da glicose, EO e morte neuronal. Linfócitos humanos compartilham comum Daérgicos mecanismos de sinalização. Além disso, a rotenona (ROT) é um inibidor que seletivamente induz apoptose através de EO nos neurônios

Daérgicas e linfócitos. Para avaliar a hipótese de que o metabolismo do G e EO estão associadas com a toxicidade do Daérgicas e sistema de EP, linfócitos humanos com ROT foram cultivados na presença ou ausência de várias concentrações de G.

**Objetivo:** este estudo analisa a resposta dos linfócitos a G (55 11, 166, 277, 555 mM) na ausência ou presença de podridão (250 microM).

**Métodos:** luz e imunocitoquímica, microscopia de fluorescência foram utilizadas técnicas para avaliar as alterações morfológicas e bioquímicas de linfócitos.

**Resultados:** a 55mM G foi eficaz em suprimir a apoptose em linfócitos induzidos por (ROT) a través da ativação de 5 vias metabólicas: (i) a través do fosfato de pentose, (ii) a través de glutational; (iii) a superóxido dismutase de antioxidante (SOD) e catalase (CAT) sistemas; (iv) Phosphoinositide 3 cinase (PI3-K). Além disso, foi observada pela primeira vez que o G resgata a apoptose de linfócitos induzida por (ROT) via ativação (v) por fator nuclear kappa-B, (NF-kB) e por regulação por baixo de p53 e Caspase-3. Mostrou que inibidores (por exemplo, LY294002), inibidores de sinalização e metabólicos (por exemplo, DHEA, BSO, BCNU, MS, DCC) reverteu parcialmente os efeitos dos tilacoides do G 55 mM de linfócitos expostos à (ROT).

**Conclusão:** estes resultados sugerem que a alta concentração de G induz simultaneamente antioxidantes e sistemas de sinalização para garantir a proteção global da célula de condições estressantes nas células Daérgicas.

## PALAVRAS-CHAVE

*Apotopse; Estresse oxidativo; Glicose; Linfócitos; Rotenona; Sinalização*

## INTRODUCTION

Parkinson's disease (PD) is a progressive neurodegenerative movement disorder associated with the selective loss of dopamine (DAergic) neurons in the *substantia nigra* pars compacta (SNpc). Although the specific molecular mechanisms leading to neuronal death in PD are not yet fully understood (1), increasing evidence from human and animal studies has suggested that oxidative stress (OS) plays a major role in the cell death process (2). Circumstantial and

epidemiological studies suggest that PD can be induced with environmental toxins that inhibit mitochondrial complex I (e.g., rotenone, ROT) (3). ROT is often used to model PD *in vitro* and *in vivo* (4,5). Interestingly, Xu and co-workers (6) have shown strong evidence for an association between low cerebral glucose (G) metabolism and the severity of PD. Furthermore, it has been demonstrated that dysregulation of G metabolism is an early event in sporadic PD (7). Despite these advances, no information is available to explain the relationship between OS, abnormal G metabolism, and neuronal demise in PD. Most importantly, the understanding of the molecular mechanisms that control neuronal survival during OS and bioenergetics crisis is essential for therapeutic intervention in PD.

Lymphocytes represent a dopaminergic system to analyze the phenomenon of OS in this neurological disorder (8,9). Specifically, we (10,11) and others (12-16) have used lymphocytes to understand the relationship between OS phenomena and PD. Previous studies have shown that ROT induced apoptosis in lymphocytes in a concentration- and time-dependent manner by a molecular signaling mechanism under standard G condition (i.e., 11 mM G, hereafter 11 G) (17). ROT toxicity thus appears to result primarily from OS signaling. Interestingly, the authors also reported that high G concentration (55 mM) abates ROT-induced apoptosis in lymphocytes (17). However, the mechanism(s) by which high G protects lymphocytes against OS is not yet established.

The aim of the present work was to evaluate the survival response of lymphocyte cells cultured with different G concentrations in the absence or presence of ROT. Our findings might contribute to understanding the mechanism of cell resistance/death signaling under stressful conditions. Taken together, our data suggest G metabolism should be considered as an additional dietary factor to be properly monitored and adjusted to confront OS and combat neuronal demise in PD patients.

## MATERIALS AND METHODS

Reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA) if not otherwise specified and were of

analytical grade or better. The 3,3'-dihexyloxacarbocyanine iodide ( $\text{DiOC}_6(3)$ , Cat. #D-273) and dihydrorhodamine (DHR, cat. #D-633) were obtained from Invitrogen Molecular Probes (Eugene, OR, USA).

## Isolation of lymphocytes

Peripheral blood lymphocytes (PBL) from venous blood of healthy adult males (ages ranging from 30 to 40 years old) were obtained by gradient centrifugation (Lymphocyte separation medium, density: 1.007 G/M; Bio-Whittaker), according to ref. (17).

## Experiments with peripheral blood lymphocytes (PBL)

**Assessment of cellular viability (CV) in PBL by light and fluorescence microscopy analysis using trypan blue (TB) and acridine orange/ethidium bromide (AO/EB) double staining.**

The PBL cells, at a cell density of  $1 \times 10^6$  cells/mL, were exposed to ROT (250  $\mu\text{M}$ ) in RPMI-1640 medium (Gibco/Invitrogen reference # 11879-020) in the absence (0 mM corresponding to 285 mOsmol/L) or presence of glucose (11, 55, 166, 277, 555 (mM) G), corresponding to 296, 340, 452, 563, 3 400 mOsmol/L, respectively) and different products of interest for 24 h at 37 °C. The PBL were then used for light and fluorescent microscopy analysis, according to ref. (17). The apoptotic index was assessed three times in independent experiments blind to researchers.

## Determination of intracellular reactive oxygen species (ROS)

**Assessment of superoxide anion radical generation**

Lymphocytes ( $1 \times 10^6$  cells/mL) were exposed to ROT (250  $\mu\text{M}$ ) in the absence or increasing concentrations of glucose (G 11, 55, 166, 277, 555) under similar experimental conditions as mentioned above. Superoxide anion radicals ( $\text{O}_2^-$ ) were evaluated

with nitro blue tetrazolium (NBT) reagent, as described in ref. (17). The assessment was repeated three times in independent experiments.

## Assessment of hydrogen peroxide ( $\text{H}_2\text{O}_2$ )

PBL cells ( $1 \times 10^6$  cells/mL) were incubated with ROT (250  $\mu\text{M}$ ), and increasing concentrations of G.  $\text{H}_2\text{O}_2$  was detected in PBL cells by using the sensitive, uncharged, non-fluorescent dihydrorhodamine 123 (DHR), as described in ref. (17). The experiments were performed in three independent settings. Parallel to  $\text{O}_2^-$  and  $\text{H}_2\text{O}_2$  evaluation, apoptotic cell percentage was established according to AO/EB staining.

## Assessment of mitochondrial transmembrane potential ( $\Delta\psi$ mt)

The PBL were treated as described above. Then, cells were incubated for 30 min at 37 °C with cationic lipophilic  $\text{DiOC}_6(3)$  (1  $\mu\text{M}$ , final concentration) to evaluate ( $\Delta\psi$  mt), according to ref. (17). The experiments were performed in three independent settings.

**Assessment of LY-294002, dehydroepiandrosterone (DHEA), L-buthionine-sulfoximine (BSO), 1,3-Bis (2-chloroethyl)-1-nitrosourea (BCNU or carmustine), 3-amino-1,2,4-triazole (AT), sodium diethyldithiocarbamate (DDC) and mercaptosuccinic acid (MS) inhibitors on lymphocytes exposed to rotenone in high glucose**

The PBL cell suspension ( $1 \times 10^6$  cells/mL) in 55 G was pre-incubated with LY294002 (5  $\mu\text{M}$ , specific PI3K inhibitor), dehydroepiandrosterone (DHEA) (5, 50, 100  $\mu\text{M}$ , specific glucose-6 phosphate dehydrogenase (G6PD) inhibitor (18,19), 1,3-Bis (2-chloroethyl)-1-nitrosourea (BCNU) (10, 100  $\mu\text{M}$ , specific glutathione reductase (GR) inhibitor (20), mercaptosuccinic acid (MS) (10 mM, specific GPx inhibitor), 3-amino-1,2,4-triazole (AT) (25, 50 mM, specific catalase [CAT] inhibitor (21), and sodium diethyldithiocarbamate (DDC) (0.5, 1, 5 mM, specific SOD inhibitor (21)), for three hours, or pre-incubated with L-buthionine-sulfoximine (BSO) (1, 5, 10 mM, specific  $\gamma$ -glutamyl-cysteine synthetase (g-GCS) inhibitor (21)) for 24 h at 37 °C before exposure to ROT (250  $\mu\text{M}$ ). PBL were then evaluated for apoptotic morphology and ( $\Delta\psi$  mt), as described above. The optimal concentration of inhibitors reported in

table 1 was based on our previous experience with inhibitors (e.g., 5 microM LY294002 and 10 nM PDTC (17), based on cited literature (e.g., 50 mM DHEA) (19), or based on the minimal concentration at which maximal inhibition and/or non-toxic effects were reported.

#### **Immunocytochemistry detection of NF- $\kappa$ B, p53, and caspase-3 transcription factor proteins**

The supplier's protocol (Santa Cruz Biotechnology, goat ABC Staining System: cat # sc-2023) was followed for immunocytochemistry using primary goat polyclonal antibodies NF- $\kappa$ B p65 (C-20)-G (Santa Cruz Biotechnology cat#sc-372-G), p53 (FL-393) (Santa Cruz Biotechnology cat #sc-6243-G), and cleaved caspase-3 p11 (h176) (Santa Cruz Biotechnology cat #sc-22171-R).

#### **Statistical analysis**

Data are presented as the means  $\pm$  S.D. of three independent experiments. One-way ANOVA analyses with Bonferroni or Games-Howell *post-hoc* comparison were calculated using SPSS 18 software. A *p*-value of \* $< 0.05$  and \*\* $< 0.001$  was considered significant.

#### **Photomicrography**

The light microscopy photomicrographs were taken using a Zeiss (AxioStart 50) microscope equipped with a Canon PowerShot G5 digital camera.

### **RESULTS**

#### **Glucose protects lymphocytes against rotenone (ROT) irrespective of ROS and in a concentration and time-dependent fashion: Effect on the (delta psi (mt))**


To establish the PBL response to G and ROT, we initially exposed cells to increasing concentrations of G (0-555) alone or in combination with ROT (hereafter 250 mM). As shown in figure 1A, lymphocytes cultured in glucose-free medium (GF) or G alone up to 166

G induced almost no apoptotic morphology (i.e.,  $0.2 \pm 1$  % AO/EB staining). Higher G concentrations provoked apoptotic morphology (i.e.,  $30 \pm 3$ ,  $68 \pm 3$ , and  $96 \pm 2$  % AO/EB at 277, 388, and 555 G, respectively). As expected, ROT induced apoptosis in lymphocytes, but the percentage of apoptotic cell death was heterogeneous. Cell death was reduced by 70 % in the presence of 11 G and ROT (i.e.,  $32 \pm 2$  % AO/EB staining positive cells), while ROT in GF medium induced 100 % AO/EB apoptosis. Interestingly, ROT toxicity was further decreased (i.e.,  $2 \pm 1$  % AO/EB) when co-incubated with 55 G (figure 1B), but ROT-induced apoptosis slightly increased (i.e.,  $8 \pm 2$  % AO/EB) when co-cultured with 166 G (figure 1B). Cell viability (CV) evaluated with trypan blue staining inversely matched the percentage of nuclear morphological changes assessed with AO/EB staining (figures 1A and B). AO/EB staining was selected for further apoptotic morphology evaluation, because the evaluation of morphological cell changes by this method was validated as the most reliable method for the detection and quantification of cell death in terms of percentage of apoptosis, when compared to other methods (22).

It is known that either G or ROT generate ROS (23,24). As shown in figure 1A, 55 G and 166 G moderately produced superoxide anion ( $O_2^{\cdot-}$ ) and  $H_2O_2$ , as determined by nitroblue tetrazolium (NBT) reduction into formazan (FORZ), and dihydrorhodamine (DHR) into rhodamine-123 (R-123) staining, respectively. PBL cells cultured in GF medium and 11 G showed either non detectable or low  $O_2^{\cdot-}/H_2O_2$  production, respectively. By contrast,  $H_2O_2$  was dramatically increased in cells treated with ROT in GF medium (i.e., almost 100 % R-123), but no  $O_2^{\cdot-}$  was detected (figure 1B). Interestingly, when ROT was co-incubated with 55 G or 166 G, the percentages of  $O_2^{\cdot-}/H_2O_2$  positive cells were almost constant (i.e., 15-22 % FORZ and 25-33 % R-123), but ROS were higher in 11 G and ROT (40 % FORZ and 42 % R-123). A similar effect was detected in cells exposed to higher G concentration alone such as 277 G and 388 G (55-60 % FORZ and 33-26 % R-123, respectively), but extremely low percentages of  $O_2^{\cdot-}$  ( $5 \pm 1$  % FORZ) and  $H_2O_2$  positive cells ( $4 \pm 1$  % R-123) were identified in the cells treated with the highest (555 G) concentration of G (data not shown).

ROT inhibits the mitochondrial NADH-quinone oxidoreductase complex; we therefore evaluated the effect of ROT on the mitochondrial membrane





**Figure 1. Glucose (G) protects lymphocytes against rotenone (ROT)-induced apoptosis independently of reactive oxygen species.** (A) Lymphocytes ( $1 \times 10^6$  cells/mL) were incubated with increasing concentrations of G (0, 11, 55, 166 (mM)) either alone or (B) in combination with ROT (250 microM) for 24 h at 37 °C. The evaluation of  $O_2^-$ ,  $H_2O_2$ , apoptosis, and ( $\Delta\psi$  mt) were performed as described in the *Methods* section. The percentage of positive formazan (FORZ), R123,  $DiO_6(3)$  high-polarized mitochondria, cell viability (CV) and AO/EB stained cells are expressed as the mean of percentages  $\pm$  SD from three independent experiments

potential ( $\Delta\psi_{mt}$ ) on cells cultured in G at different concentrations. PBL exposed to ROT in GF medium induced a complete mitochondrial depolarization (i.e., 0 % DiOC<sub>6</sub>(3)<sup>high</sup> positive cells). By contrast, no effect on mitochondrial potential depolarization was observed when PBL were incubated with 55 G - 166 G and ROT (figure 1B) or G alone (figure 1A). Noticeably, lymphocytes exposed to ROT with 11 G induced moderate mitochondrial potential depolarization (figure 1B). Since 55 G was found to be a minimal concentration that almost completely protected lymphocytes against ROT-induced mitochondrial membrane depolarization and apoptosis, this concentration of G was thus used for further experiments.

We then evaluated the time course of H<sub>2</sub>O<sub>2</sub> generation, mitochondrial depolarization, and nuclear morphological changes in PBL treated with ROT in GF and in 55 G. PBL exposed to ROT in GF medium steadily produced H<sub>2</sub>O<sub>2</sub> over time with a maximum percentage value at 24 h compared to cells cultured in GF medium alone. Noticeably, the apoptotic morphology and mitochondrial depolarization were detected after 3 h of ROT incubation. Although H<sub>2</sub>O<sub>2</sub> generation was produced at low percentage levels over time when cells were exposed to ROT and 55 G, no mitochondrial depolarization or apoptotic morphology were detected under this treatment condition (data not shown).

### Glucose protects lymphocytes against rotenone through the pentose phosphate pathway

Once G enters the cell, it can participate in at least two metabolic pathways, namely the glycolysis and the pentose phosphate pathways (PPP) (25). The former pathway turns glucose to pyruvate, which either turns into Acetyl-CoA, an essential molecule in the generation of ATP through the tricarboxylic acid (TCA) cycle and mitochondrial potential, or it may react as an antioxidant on mitochondria (26, 27). Therefore, we investigated the role of pyruvate in ROT-induced apoptosis in lymphocytes. PBL cells were therefore incubated with increasing concentrations of sodium pyruvate (1, 5, 10, 25 mM SP) in the presence or absence of ROT (250 microM) in GF medium. SP alone was innocuous to cells.

Given that lymphocytes produced no detectable H<sub>2</sub>O<sub>2</sub> in GF medium, no antioxidant function of pyruvate was evaluated. However, almost 100 % apoptotic morphology, mitochondrial depolarization, and H<sub>2</sub>O<sub>2</sub> production were detected when cells were cultured with ROT together with SP (data not shown).

These observations prompted us to evaluate the role of the PPP in this paradigm (28). PPP converts glucose to NADPH. This last compound reduces glutathione (GSH) via glutathione reductase (GR), which converts reactive H<sub>2</sub>O<sub>2</sub> into H<sub>2</sub>O and O<sub>2</sub> by glutathione peroxidase (GPx). We also evaluated the importance of superoxide dismutase (SOD) and catalase (CAT) as antioxidant enzymes in ROT-induced death in this PBL model. As described in table 1, except BSO, all inhibitors provoked moderate increments in the percentage of apoptosis (25-30 %) and mitochondrial depolarization (20-35 %) in cells treated with ROT compared to cells exposed to the inhibitor alone or 55 G (control).

### Glucose protects lymphocytes against rotenone induced apoptosis through NF- $\kappa$ B activation and p53 and caspase-3 inhibition


To further characterize the molecular response of PBL to ROT intoxication, cells were left in GF or 55 G in the absence or presence of ROT. As depicted in figure 2, PBL in GF medium alone showed normal nuclei morphology with no detectable death markers, i.e. 0 % DAB<sup>+</sup> nuclei of NF- $\kappa$ B, p53, and caspase-3 (figures 2A-C), whereas ROT in GF medium induced cell shrinkage and 100 % DAB<sup>+</sup> (apoptotic) nuclei (figures 2D-F). Strikingly, PBL cells exposed to ROT and 55 G or 55 G alone showed NF- $\kappa$ B activation (figure 2G: ~52 % DAB<sup>+</sup> (non-apoptotic) nuclei; figure 2J ~48 % DAB<sup>+</sup> (non-apoptotic) nuclei), but p53 and caspase-3 expression were negative, i.e., ~1 % DAB<sup>+</sup> nuclei (figures 2H, I, K, L).

To confirm the involvement of signaling molecules in lymphocyte cell death induced by ROT, PBL were treated with PDTC (10 nM, specific NF- $\kappa$ B inhibitor), PFT (50 nM, specific p53 inhibitor), and NSCI (10 microM, specific caspase-3 inhibitor) in either GF or 11 G alone or in combination with ROT for 24 h at



37 °C. All inhibitors reduced ROT-induced apoptosis (e.g., 9-11 % AO/EB) and increased mitochondrial membrane potential (e.g., 85-87 % DiOC<sub>6</sub>(3)<sup>high</sup> positive cells) when cultured in the presence of 11 G compared to cells incubated in 11 G and ROT (e.g., 34 % AO/EB,

and 64 % DiOC<sub>6</sub>(3)<sup>high</sup>, respectively). It is noteworthy to mention that inhibitors were ineffective at protecting lymphocytes against ROT-induced apoptosis and mitochondrial depolarization when cultured in GF medium (data not shown).



**Figure 2. Glucose induces NF-kB activation and p53 and caspase-3 inhibition.** Lymphocyte cells ( $1 \times 10^6$  cells/mL) were incubated in glucose-free (GF) medium (A-F) or glucose (55 mM) G (G-L) or treated with ROT (250 microM) in GF (D, E, F) or 55 G (J, K, L) for 24 h at 37 °C. Cells were stained with anti-NF-kB-p65 (A, D, G, J), anti-p53 (B, E, H, K), and anti-caspase-3 (C, F, I, L) antibodies according to the procedure described in the *Methods* section. Notice that NF-kB, p53, and caspase-3 positive-nuclei (numbers represent the percentage of positive (%) dark brown colored nuclei) reflecting nuclear translocation/activation. Magnification 400x (A-G, J), 600x (H, I, K, L), inset magnification 800x

## Glucose protects lymphocytes against rotenone through PI3K activation

We evaluated whether the activation of PI3K was involved in lymphocyte resistance to ROT-induced oxidative stress, because NF- $\kappa$ B was activated and mitochondrial potential was not impaired in lymphocytes even in the presence or absence of

ROT in 55 G. Thus, PBL cells were incubated either with LY294002 (5  $\mu$ M) alone, or in combination with ROT in 55 G. Table 1 shows that the presence of LY294002 blocked the protective effect of 55 G in lymphocytes against ROT-induced apoptosis and mitochondrial depolarization compared to cells exposed to ROT and 55 G.

**Table 1. Effect of glutathione, metabolic and signaling inhibitors on lymphocytes cultured in 55 mM glucose (55G) medium with or without rotenone (ROT, 250 mM) exposure**

| Treatment                                | AO/EB (%)  | DiOC <sub>6</sub> (3) <sup>high</sup> (%) |
|--|------------|---|
| 55G                                      | 0          | 100                                       |
| ROT (250 $\mu$ M)                        | 2 $\pm$ 1  | 97 $\pm$ 2                                |
| PDTC (10nM)                              | <1 $\pm$ 0 | 99 $\pm$ 1                                |
| PDTC (10nM) + ROT (250 $\mu$ M)          | 3 $\pm$ 1  | 97 $\pm$ 2                                |
| LY294002 (5 $\mu$ M)                     | 1 $\pm$ 0  | 98 $\pm$ 1                                |
| LY294002 (5 $\mu$ M) + ROT (250 $\mu$ M) | 30 $\pm$ 2 | 65 $\pm$ 3                                |
| DHEA (50 $\mu$ M)                        | 3 $\pm$ 1  | 96 $\pm$ 1                                |
| DHEA (50 $\mu$ M) + ROT (250 $\mu$ M)    | 25 $\pm$ 2 | 65 $\pm$ 2                                |
| AT (25mM)                                | <1 $\pm$ 0 | 98 $\pm$ 1                                |
| AT (25mM) + ROT (250 $\mu$ M)            | 34 $\pm$ 1 | 63 $\pm$ 2                                |
| DDC (500 $\mu$ M)                        | <1 $\pm$ 0 | 99 $\pm$ 1                                |
| DDC (500 $\mu$ M) + ROT (250 $\mu$ M)    | 25 $\pm$ 2 | 82 $\pm$ 2                                |
| BCNU (100 $\mu$ M)                       | <1 $\pm$ 0 | 99 $\pm$ 1                                |
| BCNU (100 $\mu$ M) + ROT (250 $\mu$ M)   | 26 $\pm$ 2 | 65 $\pm$ 2                                |
| MS (10 mM)                               | 4 $\pm$ 1  | 97 $\pm$ 2                                |
| MS (10 mM) + ROT (250 $\mu$ M)           | 20 $\pm$ 2 | 65 $\pm$ 3                                |
| BSO (10mM)*                              | 2 $\pm$ 1  | 96 $\pm$ 2                                |
| BSO (10mM) + ROT (250 $\mu$ M)           | 72 $\pm$ 2 | 40 $\pm$ 3                                |

PBL cell suspension (1 x 10<sup>6</sup> cells/mL) was cultured in 55 (mM) G without or with specific PI3K, G6PD, GR, SOD, CAT and GPx inhibitors in the absence or presence of ROT (250  $\mu$ M) for 24 h at 37 °C. Notice that BSO (10 mM), inhibitor of g-GCS, was pre-incubated for 48 h before ROT exposure. Nuclear morphologic changes and loss of the mitochondrial membrane potential ( $\Delta\psi$  mt) indicative of apoptosis were evaluated using AO/EB and DiOC<sub>6</sub>(3) staining, as described in the *Methods* section. Apoptosis percentage and high-polarized mitochondria (green fluorescent DiOC<sub>6</sub>(3) high positive cells) are expressed as the mean percentage (%)  $\pm$  SD from two independent experiments (with two replicas for each experiment)

## DISCUSSION

The present study shows for the first time that high glucose (55 G) was effective in suppressing ROT-induced apoptosis in lymphocytes via five pathways, involving (i) the pentose phosphate pathway (PPP), (ii) the glutathione (GSH) pathway, (iii) the SOD and CAT antioxidant systems, and (iv) PI3-K signaling. Moreover, it is also shown for the first time that G induced lymphocyte survival through (v) NF- $\kappa$ B activation and down-regulation of p53 and caspase-3. These experiments revealed the complex nature of G protection in cells. This conclusion is based on the following observations. We confirm that ROT generates  $O_2^{\cdot-}/H_2O_2$ , which was associated with decreased ( $\Delta\psi$  mt) and morphological apoptotic nuclei under standard culture conditions (i.e., 11 mM glucose, 11 G), as described in ref. (17). In accordance with those observations, lymphocytes cultured in GF and exposed to ROT displayed the typical morphological characteristics of apoptosis such as drastic cellular volume diminution, chromatin condensation and/or nuclei fragmentation concomitantly with a complete mitochondrial membrane depolarization. Interestingly, 55 G was able to maintain ( $\Delta\psi$  mt) potential and nuclear morphology against ROT toxicity to control values. These observations comply with the notion that G provides the cell's energy by serving as a primary substrate for the generation of ATP, thereby providing ( $\Delta\psi$  mt) maintenance and cell survival. Indeed, pyruvate which is a natural metabolic intermediate *en-route* to the mitochondrial tricarboxylic acid cycle, may exert antioxidant activity, thus inhibiting ( $\Delta\psi$  mt) collapse and apoptosis. However, we found that sodium pyruvate (SP, 1-25 mM range), as sole energy source, did not protect lymphocytes against ROT-induced apoptosis. Moreover, the percentages of apoptosis and mitochondrial depolarization were similar to those observed when lymphocytes were cultured in GF supplemented with ROT. Although our data clearly suggest that SP did not protect lymphocytes against ROT-induced mitochondrial dysfunction and apoptosis, we do not discard the possibility that SP may serve as an  $H_2O_2$  scavenger under other specific experimental conditions (e.g., lymphocytes exposed to ROT in the presence of G plus SP). Our data thus suggest that G metabolism in non-dividing cells might be preferentially routed through the PPP

(28, 29) rather than the glycolysis pathway, which is the prevalent G metabolic route in active dividing cells when challenged with oxidant stressors (30, 31). Interestingly, the inactivation of the glycolytic enzyme glyceraldehyde-3-phosphate dehydrogenase (GAPDH) by  $H_2O_2$  may serve as a cellular switch to reroute the metabolic carbohydrate flux from glycolysis to the PPP under OS conditions (32).

We found that except BSO, all other metabolic and signaling inhibitors moderately reversed 55 G-induced survival in lymphocytes under rotenone exposure. These data suggest that G produces functional redundancy (i.e., one part in a system can completely or partially compensate the loss of another) to ensure global cell protection against stress stimuli. Several considerations support this view. First, as a consequence of the pharmacological blockage of G6PD, the supply of NADPH in lymphocyte cells can be fulfilled through several one-step enzymatic reactions catalyzed by NADP<sup>+</sup>-dependent dehydrogenases from the Krebs' cycle, by pyridine nucleotide transhydrogenase powered by the proton motive force, and/or by NADH kinase which directly phosphorylates NADH to form NADPH (33). In case of PPP breakdown, glycolysis might take over NADPH production, but G is absolutely required to run the defense mechanism. Second, blockage of glutathione system might activate other NADPH-dependent oxidative defense systems such as thioredoxin reductase and peroxiredoxin protein (34). One may conclude therefore that the continuous supply of NADPH depends on G availability and on the cyclic metabolic PPP/glycolysis network.

Taken together, our data comply with the notion that lymphocyte cells have alternative metabolic pathways to combat OS, even in circumstances in which some of the key enzymes fail to accomplish their function by either accident (i.e., pharmacologic inhibition) or by genetic defects (e.g., G6PD mutations). These data further suggest that ROT kills cells predominantly through OS (17, 35), and that a proper source of NADPH, GSH, and antioxidant functioning enzymes must be warranted for lymphocytes to undergo complete resistance to ROT-induced OS. Collectively, these data and ours suggest that pharmacological inhibition, deficiency, or genetically altered expression of the cellular antioxidant proteins are critical components of response to stressful conditions and that these

antioxidant systems act in a cooperative way to ensure cell protection. It is worth mentioning that lymphocytes (and neurons) exposed to extremely high glucose concentrations induce OS, mitochondrial dysfunction, and apoptosis (e.g., 277 and 388 G) (36). We concluded that optimal cellular protection against stress stimuli is achieved only when there is an appropriate balance between G and GSH metabolism via the PPP pathway, and the activities of SOD, CAT, and GPx are maintained.

The NF- $\kappa$ B plays a critical role in cellular metabolism (37). We have demonstrated that 55 G alone or in combination with ROT was able to induce the activation of NF- $\kappa$ B. Moreover, NF- $\kappa$ B was activated by ROT in lymphocytes cultured in GF medium. Since both G and ROT generated ROS, it is reasonable to think that a common molecule (e.g.,  $H_2O_2$ ) might be responsible for NF- $\kappa$ B activation in both experimental settings. In accordance with this view,  $H_2O_2$ -induced NF- $\kappa$ B through the indirect activation of kinases (38), which in turn phosphorylate the I $\kappa$ B $\alpha$  inhibitor as well as p65—two molecular components of NF- $\kappa$ B (39-41). Remarkably, we also demonstrated for the first time that the pro-apoptotic transcription factor p53 was undetectable when cells were co-incubated in 55 G and ROT, but it was evidently activated by ROT in lymphocytes cultured in GF medium. Consequently, a connection between NF- $\kappa$ B and p53 is clearly established in lymphocytes under different conditions. These data thus comply with the notion that the activation of NF- $\kappa$ B is necessary but not sufficient to promote survival or induce apoptosis in lymphocytes (42). Accordingly, we found mitochondrial depolarization and caspase-3 activation concomitantly with apoptotic morphology in lymphocytes exposed to ROT in GF medium. Taken together, our data suggest that activated NF- $\kappa$ B might induce pro-apoptotic proteins which balance the cell fate towards death response in cells under limited or nil amounts of G combined with ROT (or  $H_2O_2$ ). On the contrary, activated NF- $\kappa$ B induces anti-apoptotic proteins which balance the cell fate towards survival in a cell under optimal G concentration (e.g., 55 G) and ROT. Interestingly, NF- $\kappa$ B also up regulated g-GCS expression and enzyme activity in human bronchial epithelial NCI-H292 cells (43), thereby keeping GSH available in response to OS. These data suggest that G might promote gene transcription of survival and G metabolic genes via NF- $\kappa$ B activation and suppress gene transcription of pro-apoptotic proteins through p53 inactivation.

It is known that the PI3K-protein kinase B (PKB, also known as Akt) pathway serves as a central pivotal component of cell survival and G transport pathways. We found that pharmacological inhibition of PI3K by LY294002 reversed the survival effect of G against ROT in lymphocytes. This observation suggests that PI3K might be implicated in G-induced survival signaling in lymphocytes. Indeed, it has been shown that Akt enhances both the ubiquitination-promoting function of Mdm2 (murine double minute) and Mdm2 protein stabilization by phosphorylation of Ser<sup>186</sup>, Ser<sup>166</sup>, and Ser<sup>188</sup>, which results in the reduction of p53 protein (44, 45). Based on our present data, it is reasonable to assume that p53 is modulated by G through the PI3K-Akt pathway (46). Our findings therefore reveal that p53, but not NF- $\kappa$ B, is the critical transcription factor that may possibly balance the expression of pro-death proteins towards intracellular death decision under noxious stimuli.

Recently, Yoon and Oh (47) have shown that high G levels (> 17.5-35 mM) protected MN9D cells (i.e., dopaminergic neuronal cells) against 1-methyl-4 phenylpyridinium (MPP+)-induced apoptosis via downregulation of caspase-3 activation, DNA fragmentation, and ROS inhibition. Taken together, these data suggest that high G levels modulate DAergic neuronal and lymphocyte cells response against stressful stimuli by suppressing OS and apoptosis-induced pathways. This information may contribute to a better understanding of the intracellular molecular mechanism by which G promotes cell survival against OS. Indeed, in the brain *in vivo*, about half of the G leaving the capillaries is transferred down its concentration gradient via facilitated diffusion on the GLUT-1 transporter from the lumen (approx. 100 mM G) to the brain extracellular space (approx. 60 mM G) to neuron cells (approx. 30 mM G) via the GLUT-3 transporter (48). Recent data suggest that dysregulation of G metabolism might represent an early metabolic biomarker of sporadic PD (6). Because lymphocytes express GLU-1 and GLU-3 transporter proteins (49), display molecular death machinery (50), and G metabolism (25) similar to neurons (8,47,51), we consider that lymphocyte cells represent a remarkable non-neural cell model to understanding the signaling and metabolic regulation of apoptosis in response to stressful stimuli. Therefore, the metabolism-OS-cell death axis is a fundamental issue in the therapeutic design to prevent, delay, or



ameliorate the treatment of individuals at risk of PD as encountered in Antioquia, Colombia (52).

## Acknowledgements

This work was supported by "Comité for Development and Research [Comite para el Desarrollo y la Investigacion]-CODI", Universidad de Antioquia (UdeA), grants #2545 to CV-P & MJ-Del-Rio.

## Conflict of interest

None to declare

## BIBLIOGRAPHIC REFERENCES

- Venderova K, Park DS. Programmed cell death in Parkinson's disease. *Cold Spring Harb Perspect Med*. 2012 Aug;2(8). pii: a009365. DOI 10.1101/cshperspect.a009365.
- Blesa J, Trigo-Damas I, Quiroga-Varela A, Jackson-Lewis VR. Oxidative stress and Parkinson's disease. *Front Neuroanat*. 2015 Jul;9:91. DOI 10.3389/fnana.2015.00091.
- Goldman SM. Environmental toxins and Parkinson's disease. *Annu Rev Pharmacol Toxicol*. 2014;54:141-64. DOI 10.1146/annurev-pharmtox-011613-135937.
- Xiong N, Long X, Xiong J, Jia M, Chen C, Huang J, et al. Mitochondrial complex I inhibitor rotenone-induced toxicity and its potential mechanisms in Parkinson's disease models. *Crit Rev Toxicol*. 2012 Aug;42(7):613-32. DOI 10.3109/10408444.2012.680431.
- Johnson ME, Bobrovskaya L. An update on the rotenone models of Parkinson's disease: their ability to reproduce the features of clinical disease and model gene-environment interactions. *Neurotoxicology*. 2015 Jan;46:101-16. DOI 10.1016/j.neuro.2014.12.002.
- Xu Y, Wei X, Liu X, Liao J, Lin J, Zhu C, et al. Low Cerebral Glucose Metabolism: A Potential Predictor for the Severity of Vascular Parkinsonism and Parkinson's Disease. *Aging Dis*. 2015 Nov;6(6):426-36. DOI 10.14336/AD.2015.0204.
- Dunn L, Allen GF, Mamais A, Ling H, Li A, Duberley KE, et al. Dysregulation of glucose metabolism is an early event in sporadic Parkinson's disease. *Neurobiol Aging*. 2014 May;35(5):1111-5. DOI 10.1016/j.neurobiolaging.2013.11.001.
- Alberio T, Pippione AC, Zibetti M, Olgiati S, Cecconi D, Comi C, et al. Discovery and verification of panels of T-lymphocyte proteins as biomarkers of Parkinson's disease. *Sci Rep*. 2012;2:953. DOI 10.1038/srep00953.
- Allen Reish HE, Standaert DG. Role of  $\alpha$ -synuclein in inducing innate and adaptive immunity in Parkinson disease. *J Parkinsons Dis*. 2015;5(1):1-19. DOI 10.3233/JPD-140491.
- Jimenez del Rio M, Velez Pardo C. The hydrogen peroxide and its importance in the Alzheimer's and Parkinson's disease. *Cent Nerv Syst Agents Med Chem*. 2004 Dec;4(4):279-85.
- Jimenez del Rio M, Velez Pardo C. The bad, the good, and the ugly about oxidative stress *Oxid Med Cell Longev*. 2012;2012:163913. DOI 10.1155/2012/163913.
- Younes-Mhenni S, Frih-Ayed M, Kerkeni A, Bost M, Chazot G. Peripheral blood markers of oxidative stress in Parkinson's disease. *Eur Neurol*. 2007;58(2):78-83.
- Battisti C, Formichi P, Radi E, Federico A. Oxidative-stress-induced apoptosis in PBLs of two patients with Parkinson disease secondary to alpha-synuclein mutation. *J Neurol Sci*. 2008 Apr;267(1-2):120-4.
- Prigione A, Isaias IU, Galbussera A, Brighina L, Begni B, Andreoni S, et al. Increased oxidative stress in lymphocytes from untreated Parkinson's disease patients. *Parkinsonism Relat Disord*. 2009 May;15(4):327-8. DOI 10.1016/j.parkreldis.2008.05.013.
- Colamartino M, Santoro M, Duranti G, Sabatini S, Ceci R, Testa A, et al. Evaluation of levodopa and carbidopa antioxidant activity in normal human lymphocytes in vitro: implication for oxidative stress in Parkinson's disease. *Neurotox Res*. 2015 Feb;27(2):106-17. DOI 10.1007/s12640-014-9495-7.
- Ide K, Yamada H, Umegaki K, Mizuno K, Kawakami N, Hagiwara Y, et al. Lymphocyte vitamin C levels as potential biomarker for progression of Parkinson's disease. *Nutrition*. 2015 Feb;31(2):406-8. DOI 10.1016/j.nut.2014.08.001.
- Avila-Gomez IC, Velez-Pardo C, Jimenez-Del-Rio M. Effects of insulin-like growth factor-1 on rotenone-induced apoptosis in human lymphocyte cells. *Basic Clin Pharmacol Toxicol*. 2010 Jan;106(1):53-61. DOI 10.1111/j.1742-7843.2009.00472.x.

18. Schwartz AG, Whitcomb JM, Nyce JW, Lewbart ML, Pashko LL. Dehydroepiandrosterone and structural analogs: a new class of cancer chemopreventive agents. *Adv Cancer Res.* 1988;51:391-424.
19. Tian WN, Braunstein LD, Pang J, Stuhlmeier KM, Xi QC, Tian X, et al. Importance of glucose-6-phosphate dehydrogenase activity for cell growth. *J Biol Chem.* 1998 Apr;273(17):10609-17.
20. Weydert CJ, Zhang Y, Sun W, Waugh TA, Teoh ML, Andringa KK, et al. Increased oxidative stress created by adenoviral MnSOD or CuZnSOD plus BCNU (1,3-bis(2-chloroethyl)-1-nitrosourea) inhibits breast cancer cell growth. *Free Radic Biol Med.* 2008 Mar;44(5):856-67.
21. Han YH, Moon HJ, You BR, Kim SZ, Kim SH, Park WH. The effects of buthionine sulfoximine, diethyldithiocarbamate or 3-amino-1,2,4-triazole on propyl gallate-treated HeLa cells in relation to cell growth, reactive oxygen species and glutathione. *Int J Mol Med.* 2009 Aug;24(2):261-8.
22. Leite M, Quinta-Costa M, Leite PS, Guimarães JE. Critical evaluation of techniques to detect and measure cell death--study in a model of UV radiation of the leukaemic cell line HL60. *Anal Cell Pathol.* 1999;19(3-4):139-51.
23. Tada-Oikawa S, Hiraku Y, Kawanishi M, Kawanishi S. Mechanism for generation of hydrogen peroxide and change of mitochondrial membrane potential during rotenone-induced apoptosis. *Life Sci.* 2003 Nov;73(25):3277-88.
24. Li N, Ragheb K, Lawler G, Sturgis J, Rajwa B, Melendez JA, et al. Mitochondrial complex I inhibitor rotenone induces apoptosis through enhancing mitochondrial reactive oxygen species production. *J Biol Chem.* 2003 Mar;278(10):8516-25.
25. Maciver NJ, Jacobs SR, Wieman HL, Wofford JA, Colloff JL, Rathmell JC. Glucose metabolism in lymphocytes is a regulated process with significant effects on immune cell function and survival. *J Leukoc Biol.* 2008 Oct;84(4):949-57. DOI 10.1189/jlb.0108024.
26. Jagtap JC, Chandele A, Chopde BA, Shastry P. Sodium pyruvate protects against H<sub>2</sub>O<sub>2</sub> mediated apoptosis in human neuroblastoma cell line-SK-N-MC. *J Chem Neuroanat.* 2003 Oct;26(2):109-18.
27. Wang LZ, Sun WC, Zhu XZ. Ethyl pyruvate protects PC12 cells from dopamine-induced apoptosis. *Eur J Pharmacol.* 2005 Jan;508(1-3):57-68.
28. Riganti C, Gazzano E, Polimeni M, Aldieri E, Ghigo D. The pentose phosphate pathway: an antioxidant defense and a crossroad in tumor cell fate. *Free Radic Biol Med.* 2012 Aug;53(3):421-36. DOI 10.1016/j.freeradbiomed.2012.05.006.
29. Ralser M, Wamelink MM, Kowald A, Gerisch B, Heeren G, Struys EA, et al. Dynamic rerouting of the carbohydrate flux is key to counteracting oxidative stress. *J Biol.* 2007 Dec;6(4):10.
30. Kondoh H, Leonart ME, Bernard D, Gil J. Protection from oxidative stress by enhanced glycolysis; a possible mechanism of cellular immortalization. *Histol Histopathol.* 2007 Jan;22(1):85-90.
31. Shi DY, Xie FZ, Zhai C, Stern JS, Liu Y, Liu SL. The role of cellular oxidative stress in regulating glycolysis energy metabolism in hepatoma cells. *Mol Cancer.* 2009 Jun 5;8:32. DOI 10.1186/1476-4598-8-32.
32. Grant CM, Quinn KA, Dawes IW. Differential protein S-thiolation of glyceraldehyde-3-phosphate dehydrogenase isoenzymes influences sensitivity to oxidative stress. *Mol Cell Biol.* 1999 Apr;19(4):2650-6.
33. Shi F, Li Y, Li Y, Wang X. Molecular properties, functions, and potential applications of NAD kinases. *Acta Biochim Biophys Sin (Shanghai).* 2009 May;41(5):352-61.
34. Holmgren A, Lu J. Thioredoxin and thioredoxin reductase: current research with special reference to human disease. *Biochem Biophys Res Commun.* 2010 May;396(1):120-4. DOI 10.1016/j.bbrc.2010.03.083.
35. Watabe M, Nakaki T. ATP depletion does not account for apoptosis induced by inhibition of mitochondrial electron transport chain in human dopaminergic cells. *Neuropharmacology.* 2007 Feb;52(2):536-41.
36. Russell JW, Golovoy D, Vincent AM, Mahendru P, Olzmann JA, Mentzer A, et al. High glucose-induced oxidative stress and mitochondrial dysfunction in neurons. *FASEB J.* 2002 Nov;16(13):1738-48.
37. Tornatore L, Thotakura AK, Bennett J, Moretti M, Franzoso G. The nuclear factor kappa B signaling pathway: integrating metabolism with inflammation. *Trends Cell Biol.* 2012 Nov;22(11):557-66. DOI 10.1016/j.tcb.2012.08.001.
38. Marinho HS, Real C, Cyrne L, Soares H, Antunes F. Hydrogen peroxide sensing, signaling and regulation of transcription factors. *Redox Biol.* 2014 Feb;2:535-62. DOI 10.1016/j.redox.2014.02.006.



39. Takada Y, Mukhopadhyay A, Kundu GC, Mahabeleshwar GH, Singh S, Aggarwal BB. Hydrogen peroxide activates NF-kappa B through tyrosine phosphorylation of I kappa B alpha and serine phosphorylation of p65: evidence for the involvement of I kappa B alpha kinase and Syk protein-tyrosine kinase. *J Biol Chem*. 2003 Jun;278(26):24233-41.
40. Yang WS, Seo JW, Han NJ, Choi J, Lee KU, Ahn H, et al. High glucose-induced NF-kappaB activation occurs via tyrosine phosphorylation of IkappaBalpha in human glomerular endothelial cells: involvement of Syk tyrosine kinase. *Am J Physiol Renal Physiol*. 2008 May;294(5):F1065-75. DOI 10.1152/ajprenal.00381.2007.
41. Kamata H, Manabe T, Oka Si, Kamata K, Hirata H. Hydrogen peroxide activates IkappaB kinases through phosphorylation of serine residues in the activation loops. *FEBS Lett*. 2002 May;519(1-3):231-7.
42. Czabotar PE, Lessene G, Strasser A, Adams JM. Control of apoptosis by the BCL-2 protein family: implications for physiology and therapy. *Nat Rev Mol Cell Biol*. 2014 Jan;15(1):49-63. DOI 10.1038/nrm3722.
43. Ray S, Watkins DN, Misso NL, Thompson PJ. Oxidant stress induces gamma-glutamylcysteine synthetase and glutathione synthesis in human bronchial epithelial NCI-H292 cells. *Clin Exp Allergy*. 2002 Apr;32(4):571-7.
44. Ogawara Y, Kishishita S, Obata T, Isazawa Y, Suzuki T, Tanaka K, et al. Akt enhances Mdm2-mediated ubiquitination and degradation of p53. *J Biol Chem*. 2002 Jun;277(24):21843-50.
45. Feng J, Tamaskovic R, Yang Z, Brazil DP, Merlo A, Hess D, et al. Stabilization of Mdm2 via decreased ubiquitination is mediated by protein kinase B/Akt-dependent phosphorylation. *J Biol Chem*. 2004 Aug;279(34):35510-7.
46. Xing CG, Zhu BS, Liu HH, Lin F, Yao HH, Liang ZQ, et al. LY294002 induces p53-dependent apoptosis of SGC7901 gastric cancer cells. *Acta Pharmacol Sin*. 2008 Apr;29(4):489-98. DOI 10.1111/j.1745-7254.2008.00770.x.
47. Yoon SY, Oh YJ. Glucose Levels in Culture Medium Determine Cell Death Mode in MPP(+)-treated Dopaminergic Neuronal Cells. *Exp Neurobiol*. 2015 Sep;24(3):197-205. DOI 10.5607/en.2015.24.3.197.
48. Spector R. Nutrient transport systems in brain: 40 years of progress. *J Neurochem*. 2009 Oct;111(2):315-20. DOI 10.1111/j.1471-4159.2009.06326.x.
49. Piatkiewicz P, Czech A, Tatoń J. Glucose transport in human peripheral blood lymphocytes influenced by type 2 diabetes mellitus. *Arch Immunol Ther Exp (Warsz)*. 2007 Mar-Apr;55(2):119-26.
50. Kerr JFR, Gobe GC, Winterford CM, Harmon BV. Anatomical methods in cell death. In: *Methods in Cell Biology*. Vol. 46. San Diego: Academic Press; 1995. p. 1-39.
51. Gebril HM, Avula B, Wang YH, Khan IA, Jekabsons MB. (13)C metabolic flux analysis in neurons utilizing a model that accounts for hexose phosphate recycling within the pentose phosphate pathway. *Neurochem Int*. 2016 Feb;93:26-39. DOI 10.1016/j.neuint.2015.12.008.
52. Pineda-Trujillo N, Carvajal-Carmona LG, Buriticá O, Moreno S, Uribe C, Pineda D, et al. A novel Cys212Tyr founder mutation in parkin and allelic heterogeneity of juvenile Parkinsonism in a population from North West Colombia. *Neurosci Lett*. 2001 Feb;298(2):87-90.

