



Acta Scientiarum. Biological Sciences

ISSN: 1679-9283

eduem@uem.br

Universidade Estadual de Maringá
Brasil

Lopes Ceschini, Tiago; Massato Takemoto, Ricardo; Hideki Yamada, Fábio; Aquino Moreira, Luis
Henrique de; Cezar Pavanelli, Gilberto

Endoparasites of *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889), collected in the
tributaries Corvo and Guairacá of Paranapanema river, Paraná State, Brazil

Acta Scientiarum. Biological Sciences, vol. 32, núm. 2, 2010, pp. 125-130

Universidade Estadual de Maringá

.png, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=187114387004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Endoparasites of *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889), collected in the tributaries Corvo and Guairacá of Paranapanema river, Paraná State, Brazil

Tiago Lopes Ceschini*, Ricardo Massato Takemoto, Fábio Hideki Yamada, Luis Henrique de Aquino Moreira and Gilberto Cezar Pavanelli

Universidade Estadual de Maringá, Av. Colombo, 5790, 87020-900, Maringá, Paraná, Brazil. *Author for correspondence.
E-mail: tiagobiology@yahoo.com.br

ABSTRACT. *Steindachnerina brevipinna* is a freshwater fish from the Curimatidae family, and its occurrence is restricted to some South American countries (Argentina, Brazil, Paraguay and Uruguay), distributed in the Paraguay, lower and upper Paraná and lower Uruguay rivers. There are no parasitological studies related to this fish. The present study records and illustrates a species of digenean, *Sphincterodiplostomum musculosum* from subfamily Diplostominae, present on the ovary and three species of nematodes, *Cosmoxyinema vianai* Travassos, 1949, *Travnema travnema* Pereira, 1938, *Spinoxyuris* sp., of superfamily Oxyuridea, parasitizing the intestine of *S. brevipinna* collected in the tributaries Guairacá and Corvo of the Paranapanema River, Parana State, Brazil.

Key words: oxyuridae, diplostominae, *Steindachnerina brevipinna*, Corvo, Guairacá.

RESUMO. Endoparasitos de *Steindachnerina brevipinna* (Eigenmann e Eigenmann, 1889), coletados nos tributários Corvo e Guairacá do rio Paranapanema, Paraná, Brasil. *Steindachnerina brevipinna* é um peixe dulcícola da família Curimatidae e sua ocorrência é restrita a alguns países da América do Sul (Argentina, Brasil, Paraguai e Uruguai) e estão distribuídos nos rios Paraguai, baixo e alto Paraná e baixo Uruguai. Ainda não existem estudos parasitológicos relacionados a esse peixe. No presente trabalho foram registrados e ilustrados uma espécie de digêneo, *Sphincterodiplostomum musculosum* Dubois, 1936 da subfamília Diplostominae presente no ovário e três espécies de nematodos, *Cosmoxyinema vianai* Travassos, 1949, *Travnema travnema* Pereira, 1938 e *Spinoxyuris* sp., pertencentes à superfamília Oxyuridea localizados no intestino, para os espécimes de *S. brevipinna* coletados nos tributários Corvo e Guairacá do rio Paranapanema, Estado do Paraná, Brasil.

Palavras-chave: oxyuridae, diplostominae, *Steindachnerina brevipinna*, Corvo, Guairacá.

Introduction

The parasites of family Diplostomidae Poirer, 1886 when adults are found in birds and mammals, totaling 41 genera distributed in different subfamilies according to their host. The subfamily Alariinae includes 11 genera parasites of mammals, with genera parasitizing birds and divided into three subfamilies, Diplostominae Poirier, 1886; Crassiphialinae Sudarikov, 1960; and Codonocephalinae Sudarikov, 1959 (LUNASCHI; DRAGO, 2006).

Most nematodes of superfamily Oxyuridea Cobbold, 1864 are parasites of neotropical fishes of South America and only a few genera of family Pharyngodonidae occur in marine and freshwater fish of tropical and subtropical regions. There are seven genera known to the superfamily Oxyuridea in South America, with 11 species described (MORAVEC, 1998). The species belonging to the genus *Brasilnema*

Moravec, Kohn and Fernandes, 1992, *Cosmoxyinema* Travassos, 1949, *Cosmoxyinemoides* Travassos, 1949, *Ichthyouris* Inglis, 1969, *Parasynodontidia* Moravec, Kohn and Fernandes, 1992, *Travnema*, Pereira, 1938, are registered to fishes from Brazil (PEREIRA, 1938; TRAVASSOS, 1949; FERNANDES et al., 1983; MORAVEC et al., 1992); two species of *Parasynodontidia* and *Spinoxyuris* Petter, 1994, of fishes from Paraguay (PETTER, 1994); and only one species of *Ichthyouris* from Guyana (INGLIS, 1962). One species of Diplostomidae and three species of Pharyngodonidae were found in *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889), a freshwater fish of Curimatidae family, which is found in lakes, rivers and channels (GRAÇA; PAVANELLI, 2007). Therefore, due to their feeding habit as detritivores (FINK; FINK, 1978), *S. brevipinna* probably has a greater contact with intermediate hosts and free-living larvae that facilitates parasitism.

There are no publications of parasitological studies related to this fish, but only records of endoparasites of the genus *Diplostomum* sp. for *Steindachnerina insculpta* (PAVANELLI et al., 2004), an endemic species of the upper Paraná river (PAVANELLI; BRITSKI, 1999).

This manuscript reports and illustrates a digenean species of subfamily Diplostominae and two species and one genus of nematodes of superfamily Oxyuridea found in *S. brevipinna* collected in tributaries Guairacá and Corvo of the Paranapanema river.

Material and methods

Sixty-three specimens of *Steindachnerina brevipinna* were collected between April and September 2006 in the tributaries Guairacá and Corvo, in the lower Paranapanema river, a region flooded by the reservoir of the Rosana hydroelectric power plant (Figure 1) (22°36'S; 52°52'W), state of Paraná, Brazil (CESP, 1998).




Figure 1. Location of the rivers Corvo and Guairacá, (tributaries of the low Paranapanema river) and of the reservoir of the Rosana hydroelectric power plant.

Gills nets with different mesh sizes were used for capturing the fish, disposed for a period of 24 hours, with collections every eight hours. Parasites were collected and processed according to Eiras et al. (2006)

The measurements were standardized in millimeters and the illustrations were made with the aid of a drawing tube attached to a Nikon YS 2 microscope. Representative specimens were deposited in the Helminthological Collection of the Oswaldo Cruz Institute, Rio de Janeiro, Rio de Janeiro State, Brazil.

Results and discussion

Class Trematoda

Subclass Digenea

Family Diplostomidae Poirier, 1886

Subfamily Diplostominae Poirier, 1886

Sphincterodiplostomum musculosum Dubois, 1936 (Figure 2)




Figure 2. *Sphincterodiplostomum musculosum* (metacercariae) Dubois, 1936. PS – pseudosuckers; OV – oral sucker; PH – pharynx; OE – oesophagus; VT – vitellaria; IC – intestinal caeca; VS – ventral sucker; TO – tribocytic organ; OV – ovary; TE – testes; SP – sphincter.

Measurements based on 15 specimens.

Body, 1.775 – 3.075 (2.553) long and 1.075 – 1.675 (1.396) wide, *forebody*, 1.150 – 1.900 (1.530) long and 1.075 – 1.675 (1.508) wide; *hindbody*, 0.600 – 1.375 (1.033) long and 0.500 – 0.800 (0.665) wide, with a dense vitellaria concentration in the anterior

region of *forebody*. Pharynx 0.084 – 0.117 (0.104) long and 0.045 – 0.084 (0.054) wide, located after an subterminal oral sucker, 0.100 – 0.200 (0.145) long and 0.130 – 0.200 (0.213) wide, with two laterals pseudosuckers. Ventral sucker, 0.130 – 0.210 (0.162) long and 0.180 – 0.250 (0.213) wide, above of tribocytic organ, 0.290 – 0.440 (0.384) long 0.350 – 0.650 (0.490) wide, located in medio-posterior region of *forebody*. Subterminal sphincter, one ovary and a pair of testes in tandem, posterior teste with 0.210 – 0.400 (0.290) long and 0.200 – 0.510 (0.410) wide and anterior teste with 0.130 – 0.390 (0.217) and 0.250 – 0.350 (0.307), located in medio-posterior region of *hindbody*.

Taxonomic Summary

Host: *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889).

Site: Ovary.

Locality: Tributaries Guairacá and Corvo of the low Paranapanema river, Paraná State, Brazil.

Material: CHIOC: no. 37263a-c

Mean Intensity of infection: 52.63, being found up to 320 parasites in a single host.

Prevalence: 90.47 %

Others hosts: *Cyphocharax gilbert* (ABDALLAH et al., 2005) and *Hemisorubim platyrhynchos* to *Sphincterodiplostomum* sp. (GUIDELLI et al., 2003).

Distribution

Sphincterodiplostomum musculosum first described by Dubois (1936) in fishes of Neotropical region (NIEWIADOMSKA, 2002). Was found also in Guandu river, Rio de Janeiro State, Brazil (ABDALLAH et al., 2005), in Paraná river, Brazil, as *Sphincterodiplostomum* sp. (GUIDELLI et al., 2003) and their adult form in birds of Formosa Province, Argentina (LUNASCHI; DRAGO, 2006).

Comments

Quoy and Gaimard in 1824 described this species parasitizing the eyes of *Cyphocharax gilbert*. In *S. brevipinna* was found in large quantities in the ovaries, being the most prevalent endoparasite. The adult form was described by Lunaschi and Drago (2006), parasitizing the intestine of *Ardea alba*, Linnaeus 1758.

The illustration shows clearly the body bipartite in *forebody* and *hindbody*. The anterior region has vitellaria concentrated, oral sucker, pseudosuckers, ventral suckers and tribocytic organ. In posterior region has one ovary, one pair of testes and one subterminal sphincter.

Filo Nematoda

Family Pharyngodonidae Travassos, 1919

Superfamily Oxyuridea Cobbold, 1864

Cosmoxyinema vianai Travassos, 1949 (Figure 3)




Figure 3. *Cosmoxyinema vianai* Travassos, 1949. a) (anterior region): BC – buccal capsule; PH – pharynx; NR – nerve ring; IT – isthmus; BU – Bulb; OV – ovary; VU – vulva; UT – uterus with eggs. b) (posterior region): CA – caudal region. c) Egg.

Measures based on 15 specimens.

Female: Elongated body with a abrupt decrease of diameter after anus, forming a thin and long tail, 2.325 – 4.225 (3.368) long and 0.096 – 0.270 (0.158) wide; buccal capsule absent of teeth and well developed; oesophagus, 0.280 – 0.390 (0.320) long and 0.048 – 0.091 (0.069) wide, elongated and divided in: anterior region of oesophagus, 0.288 – 0.390 (0.327) long and 0.038 – 0.069 (0.047) wide; Isthmus, 0.012 – 0.021 (0.015) long and 0.090 – 0.024 (0.016) wide; and bulb, 0.062 – 0.088 (0.074) long and 0.048 – 0.091 (0.069) wide; vulva situated at the end of first quarter of body, 0.600 – 1.020 (0.739) of anterior end until the vulva; egg asymmetric, elongated with a small operculum, 0.079 – 0.187 (0.146) and 0.024 – 0.048 (0.037).

Male: Unknown.

Taxonomic Summary

Host: *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889).

Site: Intestine.

Locality: Tributaries Guairacá and Corvo of the low Paranapanema river, Paraná State, Brazil.

Material: CHIOC: no. 37260, no. 37261.

Mean Intensity of infection: 1.33, being found maximum three parasites in a single host.

Prevalence: 28.75%.

Others hosts: *Pseudocurimata gilberti* (= *Cyphocarax gilberti*) and *Pseudocurimata* sp. (MORAVEC, 1998).

Distribution

Species originally described by Travassos (1949), in Barra Seca river and Juparana Lake, Espírito Santo State, Brazil. Later recorded in Paraná and Mogi Guaçu river, São Paulo State, Brazil by Vicente et al. (1985) and Moravec et al. (1992) respectively.

Comments

The genus *Cosmoxyinema* was described by Travassos in 1949, and has only one species (MORAVEC, 1998). Of all nematodes found in *S. brevipinna*, *Cosmoxyinema vianai* showed most long. In most cases only two parasites were found in a single host, always female. There was no record for males.

According Moravec (1998), Travassos not find the male of the species. The same is true for genus *Cosmoxyinemoides*, very similar to the *Cosmoxyinema*.

The illustration shows in the anterior region, cuticle striated, oesophagus and nerve ring, eggs distributed only in one region of the body, ovary and muscular vagina; in posterior region and an abrupt decrease of diameter after anus, forming a thin and long tail.

Travnema travnema Pereira, 1938 (Figure 4)




Figure 4. a-c) Female of *T. travnema* Pereira, 1938. a) (anterior region): BC – buccal capsule; PH – pharynx; NR – nerve ring; IT – isthmus; BU – bulb; b) (vulva region): VU – Vulva. c) egg development, initial fase. d-e) Male of *T. travnema* Pereira, 1938. d) (anterior region): BC – buccal capsule; PH – pharynx; NR – nerve ring; IT – isthmus; BU – bulb. e) (posterior region): SP – Spicule.

Measures based on 15 female and one male specimens.

Female: small compared to other nematodes, 1.600 – 2.425 (1.989) long and 0.084 – 0.264 (0.163) wide; Buccal capsule lack teeth and well developed; oesophagus, 0.144 – 0.187 (0.167) long and 0.031 – 0.069 (0.048) wide, divided in: anterior region of oesophagus (developed and muscular) 0.072 – 0.086 (0.079) long and 0.031 – 0.048 (0.041) wide; Isthmus, 0.024 – 0.040 (0.032) long and 0.016 – 0.024 (0.019) wide; and bulb, 0.036 – 0.079 (0.056) long and 0.031 – 0.069 (0.049) wide; vulva situated at the posterior region of second third of body, with a distance of 0.700 – 1.250 (0.947) anterior region to reproductive organ; egg asymmetric, elongated with at small operculum, 0.122 – 0.179 (0.227) long and 0.040 – 0.072 (0.050) wide.

Male: body size two times smaller than the female, 1.130 long and 0.055 wide. Oesophagus, 0.125 long and 0.025 wide; divided in: anterior region of oesophagus, 0.055 long and 0.020 wide; Isthmus, 0.037 long and 0.013 wide; and Bulb, 0.009 long and 0.014 wide; Conical tail with sharp curvature, 0.118 long and 0.080 wide in anterior region and 0.010 near to end; One spicule, 0.047 long and wide base 0.007, gubernaculum absent.

Taxonomic Summary

Host: *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889).

Site: Intestine.

Locality: Tributaries Guairacá and Corvo of the low Paranapanema river, Paraná State, Brazil.

Material: CHIOC: no. 37258, no. 37259.

Mean Intensity of infection: 1.26, maximum of two parasites found in a single host.

Prevalence: 30.15%.

Others hosts: *Pseudocurimata elegans*, *P. gilbert* (= *Cyphocarax gilbert*), *P. plumbea* and *Astyanax bimaculatus lacustris* (MORAVEC, 1998).

Distribution

Described by Pereira (1938), *Travnema travnema* was found in northeastern Brazil, in Porangaba, Soure and Tauape lakes, Fortaleza, Ceará State, Brazil. Later found in Mogi Guaçu, Pirassununga and Paraná rivers, Brazil by Kohn et al. (1985), Kohn and Fernandes (1987) and Moravec et al. (1992).

Comments

The genus *Travnema* was described by Pereira (1938) and has two species, *T. araujo* and *T. travnema*. The main difference between them is the presence of denticles in the buccal capsule of the first species. Besides being one of the oldest genera of Superfamily Oxyuridea, with *Laurotravassoxyuris*

described in same year by Vigueiras (MORAVEC, 1998), *T. travnema* had the highest distribution and prevalence of all parasites found in *S. brevipinna*.

This parasite is small and therefore difficult to visualize on stereoscopic microscope, being best viewed after viscera washing with sieve (154 μ m).

The illustration shows in the anterior region of the parasite, transverse striations, buccal capsule, oesophagus and nerve ring, vulva located in posterior region of the second third of body, in posterior region of parasite can be observed, the final part of intestine and anus; egg in initial and final stage.

Spinioxyuris sp. (Figure 5)




Figure 5. a) *Spinioxyuris* sp. Petter, 1994. b) (anterior region): BC – buccal capsule; PH – pharynx; BU – bulb. c) (posterior region): SP – spicule; GC – genital cone; CA – caudal alae; TA – tail.

Measures based on one male.

Male: small body, 1.130 long by 0.055 wide. Pharynx without isthmus, 0.192 long and 0.045 wide; bulb with irregular surface, 0.012 long and 0.045 wide. Terminal tail presence, 0.118 long, 0.080 anterior and 0.010 posterior wide; caudal alae presence. Genital cone well developed with only one simple and grooved spicule, 0.066 long and 0.009 wide base, gubernaculum absent.

Female: Unknown.

Taxonomic Summary

Host: *Steindachnerina brevipinna* (Eigenmann and Eigenmann, 1889).

Site: Intestine.

Locality: Tributaries Guairacá and Corvo of the low Paranapanema river, Paraná State, Brazil.

Material: CHIOC: no. 37262.

Mean Intensity of infection: 1, only one parasite found in a single host.

Prevalence: 1.58%.

Others hosts: *Oxydoras kneri* to *Spinioxyuris oxydoras* (MORAVEC, 1998).

Distribution

Genus described by Petter (1994), in Paraná river, Alto Paraná Province, Paraguay.

Comments

The *Spinioxyuris* genus was described by Petter (1994) with records of both sexes for a single species, *Spinioxyuris oxydoras* in host *Oxydoras kneri*, collected in Paraná river, Alto Paraná Province, Paraguay (MORAVEC, 1998). This parasite species possibly belongs to a new species, but this single specimen limits the identification only to genus level.

The illustration shows in the anterior region of the parasite, one irregular bulb which probably obtained such features due to methods of fixation. In posterior region can be observed a developed genital cone, caudal alae and only a single spicule with grooves.

References

- ABDALLAH, V. D.; AZEVEDO, R. K.; LUQUE, J. L. Ecologia da comunidade de metazoários parasitos do sairú *Cyphocharax gilbert* (Quoy e Gaimard, 1824) (Characiformes: Curimatidae) do rio Guandu, estado do Rio de Janeiro, Brasil. **Revista Brasileira de Parasitologia Veterinária**, v. 14, n. 4, p. 154-159, 2005. CESP-Companhia Energética de São Paulo.
- Conservação e manejo nos reservatórios:** limnologia, ictiologia e pesca. São Paulo: CESP, 1998.
- EIRAS, J. C.; TAKEMOTO, R. M.; PAVANELLI, G. C. **Métodos de estudo e técnicas laboratoriais em parasitologia de peixes**. 2. ed. Maringá: Eduem, 2006.
- FERNANDES, M. T.; CAMPOS, M. S.; ARTIGAS, P. T. *Travnema araujoi* n. sp. (Nematoda, Pharyngodonidae), parasito de *Curimatus gilberti* Quoy & Gaimard (Pisces, Characidae). **Revista da Faculdade de Medicina Veterinária e Zootecnia da Universidade de São Paulo**, v. 20, n. 1, p. 35-41, 1983.
- FINK, W. I.; FINK, S. A Amazônia Central e seus peixes. **Acta Amazonica**, v. 8, n. 4, p. 19-42, 1978.
- GRAÇA, W. J.; PAVANELLI, C. S. **Peixes da planície de inundação do alto rio Paraná e áreas adjacentes**. Maringá: Eduem, 2007.
- GUIDELLI, G. M.; ISAAC, A.; TAKEMOTO, R. M.; PAVANELLI, G. C. Endoparasite Infracommunities of *Hemisorubim platyrhynchos* (Valenciennes, 1840) (Pisces: Pimelodidae) of the Baía river, upper Paraná river floodplain, Brazil: specific composition and ecological aspects. **Brazilian Journal of Biology**, v. 63, n. 2, p. 261-268, 2003.
- INGLIS W. G. *Ichthyouris* ro gen. et sp. nov. (Nematoda): an oxyurid from a freshwater fish. **Journal of Helminthology**, v. 36, n. 1, p. 45-50, 1962.
- KOHN, A.; FERNANDES, B. M. M. Comparative study of helminth parasites of fishes from Mogi Guassu River, collected during expeditions between 1927 and 1985. **Memórias do Instituto Oswaldo Cruz**, v. 82, n. 1, p. 483-500, 1987.
- KOHN, A.; FERNANDES, B. M. M.; MACEDO, B.; ABRAMSON, B. Helminth parasites of freshwater fishes from Pirassununga, SP, Brazil. **Memórias do Instituto Oswaldo Cruz**, v. 80, n. 3, p. 327-336, 1985.
- LUNASCHI, L. I.; DRAGO, F. B. First report of adult specimens of *Sphincterodiplostomum musculosum* (Digenea, Diplostomidae). **Parasitology International**, v. 55, n. 1, p. 7-10, 2006.
- MORAVEC F. **Nematodes of freshwater fishes of the neotropical region**. Praha: Academia, 1998.
- MORAVEC, F.; KOHN, A.; FERNANDES, B. M. M. Three new species of oxyuroid nematodes, including two new genera, from freshwater catfishes in Brazil. **Systematic Parasitology**, v. 21, n. 3, p. 189-201, 1992.
- NIEWIADOMSKA, K. Family Diplostomidae Poirier, 1886. In: GIBSON, D. I.; JONES, A.; BRAY, R. A. (Ed.). **Keys to the Trematoda**. London: CAB International and The Natural History Museum, 2002. p. 167-196.
- PAVANELLI, C. S.; BRITSKI, H. A. Description of a new species of *Steindachnerina* (Teleostei: Characiformes: Curimatidae) from the upper Paraná river basin, Brazil. **Ichthyological Exploration of Freshwaters**, v. 10, n. 3, p. 211-216, 1999.
- PAVANELLI, G. C.; MACHADO, M. H.; TAKEMOTO, R. M.; GUIDELLI, G. M.; LIZAMA, M. A. P. Helminth fauna of fishes: diversity and ecology aspects. In: THOMAZ, S. M.; AGOSTINHO, A. A.; HAHN, N. S. (Ed.). **The upper Paraná river and its floodplain: physical aspects, ecology and conservation**. Leiden: Backhuys Publishers, 2004. p. 309-329.
- PEREIRA, C. *Travnema travnema* n. gen. n. sp. (Nematoda: Oxyuridae) parasito de *Curimatus elegans* (Pisces: Characidae) no Nordeste Brasileiro. In: **Livro Jubilar do Dr. Lauro Travassos**. Rio de Janeiro: Brasil, 1938. p. 381-385.
- PETTER, A. J. Nematodes de Poissons de Paraguay. VII. Oxyuroidea: *Spinioxyuris oxydoras* n. g., n. sp. **Revue Suisse de Zoologie**, v. 101, n. 3, p. 761-769, 1994.
- TRAVASSOS, L. Contribuição ao conhecimento da fauna helmintológica dos peixes de água doce do Brasil. IV. Dois novos generos de Cosmocercidae (Nematoda) e uma nota helmintológica. **Memórias do Instituto Oswaldo Cruz**, v. 46, n. 3, p. 633-637, 1949.
- VICENTE, J. J.; RODRIGUES, H. O.; GOMES, D. C. Nematóides do Brasil. 1. Parte: nematóide de peixes. **Atas da Sociedade de Biologia do Rio de Janeiro**, v. 25, n. 1, p. 1-79, 1985.

Received on June 26, 2008.

Accepted on February 19, 2009.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.