



Revista Ciência Agronômica

ISSN: 0045-6888

ccarev@ufc.br

Universidade Federal do Ceará
Brasil

Vieira, Rafael Augusto; Scapim, Carlos Alberto; Tessmann, Dauri José; Teruhiko Hata, Fernando
Diallel analysis of yield, popping expansion, and southern rust resistance in popcorn lines
Revista Ciência Agronômica, vol. 42, núm. 3, julio-septiembre, 2011, pp. 774-780
Universidade Federal do Ceará
Ceará, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=195318899025>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Diallel analysis of yield, popping expansion, and southern rust resistance in popcorn lines¹

Análise dialélica da produtividade, capacidade de expansão e resistência à ferrugem-polissora em linhagens de milho-pipoca

Rafael Augusto Vieira^{2*}, Carlos Alberto Scapim³, Dauri José Tessmann³ e Fernando Teruhiko Hata³

Abstract - Ten popcorn inbred lines were crossed in a circulant partial diallel matching scheme (group of lines derived from IAC 112 × group of lines from 'Zaeli'). The hybrids were evaluated in respect to grain yield (GY), popping expansion (PE), and resistance to southern rust (*Puccinia polysora*, Pp), in randomized block design, with three replications. Data was examined by analyses of variance and Kempthorne and Curnow's diallel analysis. General combining ability (GCA) was significant ($p < 0.10$) in the set 2-Zaeli for GY, PE, and partial resistance to Pp. This indicated that additive gene effects are predominant, and therefore breeding methods of recurrent selection are recommended. However, none of lines had outstanding GCA for GY and PE. The line P_{8,5} was recommended to form breeding populations for resistance to Pp. Specific combining ability (SCA) was detected for GY. The P_{9,24} × P_{8,6} might be promising for GY, if exploited using interpopulational breeding methods.

Key words - *Zea mays*. General and specific combining ability. *Puccinia polysora*. Plant breeding.

Resumo - Dez linhagens de milho-pipoca (grupo de linhagens IAC 112 × grupo de linhagens 'Zaeli') foram cruzadas em esquema dialélico parcial circulante. Os híbridos foram avaliados quanto ao rendimento de grãos (RG), capacidade de expansão (CE) e resistência parcial à ferrugem-polissora (*Puccinia polysora* Pp), em delineamento de blocos completos com tratamentos ao acaso, com três repetições. As características foram utilizadas para análises de variância e dialélicas de Kempthorne e Curnow. A capacidade geral de combinação (CGC) foi significativa ($p < 0,10$) no grupo Zaeli para RG, CE e resistência à Pp. Isto indicou a predominância de genes aditivos, e que métodos de seleção recorrente são recomendados para estas características. Não foram encontradas linhagens com CGC favoráveis para RG e CE. A linhagem P_{8,5} foi recomendada para integrar populações para melhoramento visando resistência à Pp. A capacidade específica de combinação somente foi detectada para RG. O híbrido P_{9,24} × P_{8,6} pode ser promissor para RG se explorando usando métodos intrapopulacionais de melhoramento.

Palavras-chave - *Zea mays*. Capacidades geral e específica de combinação. *Puccinia polysora*. Melhoramento vegetal.

* Autor para correspondência

¹Recebido para publicação em 09/06/2010; aprovado em 25/04/2011

Financiamento via CNPq - National Counsel of Technological and Scientific Development, Brazil and Araucaria Foundation

²Universidade Estadual de Maringá, Maringá-PR, Brasil, rfavieira@msn.com

³Universidade Estadual de Maringá, Maringá-PR, Brasil, cscapim@uem.br, djtessmann@uem.br, prox_fdinhu@hotmail.com

Introduction

Popcorn and other specialty maize are producing value-added properties to the grains in Brazil. Despite the important growth in the consumption of popcorn, diverse studies of the Ministry of Agriculture have determined that production is still limited with regards to the potential market (ARNHOLD et al., 2009; VIEIRA et al., 2009a). However, this situation is being reverted by important advances achieved in the breeding programs (ARNHOLD et al., 2009; SANTOS et al., 2008; SCAPIM et al., 2010; SILVA et al., 2009; TRINDADE et al., 2010; VIEIRA et al., 2009a).

To obtain superior lines and hybrids is one of the major strategies in popcorn breeding programmes for development of high-yielding and popping expansion cultivars. Both additive and nonadditive gene effects are exploited during populations improvement and development of inbred lines. The relative importance of additive and nonadditive effects on total genetic variance is fundamental. This defines which intrapopulation or interpopulation methods are most advantageous.

The genetic potential of maize lines and hybrids has been studied since Jones (1918) suggested the commercial use of double-crosses hybrids. In order to evaluate genotypes, the diallel crosses and top-crosses are useful. Diallel provides information on combining ability and gene effects, and constitute a valuable tool for maize breeders. Nevertheless, the evaluation of diallel crosses among inbred lines are recent and incipient in popcorn germplasm (LARISH; BREWBAKER, 1999; PINTO et al., 2007a, 2007b; VIEIRA et al., 2009b).

The southern rust is caused by *Puccinia polysora* Underw., and well-adapted to high and low lands in Brazil, and other tropical and sub-tropical environments where maize grown (SHURTLEFF, 1992). This rust has affected leaves causing yield losses up to 50% (CASTELLANOS et al. 1998; RODRIGUES-ARNON et al., 1980). The partial resistance to southern rust, or slow-rusting, consists of the ability to decrease the disease rates of development, usually by small lesions and less sporulation (ZUMMO, 1988). Since the area under the disease progress curve (AUDPC) measures the disease development over a period, this criteria is commonly to assess partial resistance levels under field conditions.

To prevent yield losses, the development of genotypes with genetic resistance to diseases is a major strategy in maize breeding (VIEIRA et al., 2009b). In fact, the flint germplasm available to popcorn breeding is narrow and does not performed as well as common maize. Consequently, it provides poorer agronomic traits including diseases susceptibility (ZIEGLER; ASHMAN, 1994).

In the present study, the combining ability of popcorn inbred lines was evaluated in circulant partial diallel matching scheme for grain yield, popping expansion, and partial resistance to southern rust (*Puccinia polysora*, Pp). This study intended to measure the breeding possibilities using this germplasm, and firstly reports the inheritance pattern for the resistance to Pp in popcorn germplasm.

Material and methods

Two sets of popcorn lines were crossed in circulant partial diallel matching scheme, with three hybrid combinations per line. The sets were: i) set 1-IAC112: five S_8 inbred lines, derived from self-pollinations of the hybrid IAC 112, a cultivar with temperate and tropical genealogy; and, ii) set 2-Zaeli: five S_8 inbred lines, derived from self-pollinations of the 'Zaeli', a unknown genetic material but with temperate genealogy. The fifteen hybrids were obtained and evaluated in randomized block design, with three replications, in Maringá, PR, Brazil (23°21' South latitude, 52°04' West longitude and 510 m altitude), during the 2007/08 cropping season. One check hybrid (single hybrid IAC 112, moderately susceptible to Pp) was included, and its mean was compare to diallel crosses. In order to study the diallel crosses, the check hybrid was not included in statistical analyses.

The experimental plots consisted of a 5 m long row with 25 plants and spaced 0.9 m apart. Plots were fertilized with 20 kg ha⁻¹ N, 50 kg ha⁻¹ P₂O₅, and 50 kg ha⁻¹ K₂O at sowing, and 100 kg ha⁻¹ N sidedressing, when the plants reached development stage V₄. Cultural treatments were based on the crop requirements; however, fungicide applications were not performed. The Pp severity was evaluated under natural infestations on hybrids. The ear leaf and the first leaves below and above of five plants were sampled and evaluated per plot. Three evaluations were carried out approximately every 12 days, after 15 days of flowering. Severity data was used to calculate the area under the disease progress curve (AUDPC), as proposed by Campbell and Madden (1990).

The ears were hand-harvested and threshed each plot, and grain yield data was obtained and corrected to 15% moisture. Next, data was corrected by number of plants per plot, as according to Schmildt et al. (2001), and transformed to kg ha⁻¹. Popping expansion was evaluated in a sample of 30.0 g of grains that were popped in a automatic machine developed by Embrapa-CNPDIA, at 237 °C during 120 s. Grains popped were collected from central region of ears, of which had storage until 14% moisture was reached. The popped volume was measured using a beaker of 2,000 mL capacity.

Grain yield, popping expansion, AUDPC values for southern rust, and the southern rust severity in the last evaluation (at 92 days after sowing) were used in the statistical analyses. Bartlett's test was applied to verify the homogeneity of variances. The method of Kolmogorov-Smirnov was used to verify the normal distribution of errors. Since the parametric assumptions were satisfied data transformations were not necessary.

Two-way analyses of variance were carried out. Next, the means of traits were used to perform the partial diallel analysis, as Kempthorne and Curnow (1961). The statistic model was:

$$Y_{ij} = \mu + g_i + g_j + s_{ij} + \varepsilon_{ij} \quad (1)$$

in which,

μ : overall mean;

g_i and g_j : effects of general combining ability (GCA) associated to the lines i (set 1-IAC 112) and j (set 2-Zaeli);

s_{ij} : effect of specific combining ability (SCA) to the combination of the lines i and j ;

and ε_{ij} : mean experimental error to the combination ij . The mean square errors and degrees of freedom from analysis of variance also were included in diallel analyses.

The percentages of sum of squares from GCA and SCA were calculated as follows: i) Percentage of sum of squares from GCA: $\%SSG = SSG / (SSG + SSS) \times 100$, in which SSG: sum of squares from GCA and SSS: sum of squares from SCA; ii) Percentage of sum of squares from SCA: $\%SSS = 100 - \%SSG$, in which $\%SSG$: percentage of sum of squares from GCA. Random effects of genotypes were considered for the analyses. Data analyses were carried out using the program Genes (Universidade Federal de Viçosa, Viçosa, MG).

Results and discussion

Mean grain yield (GY) was from 3,200.8 ($P_{9.27} \times P_{8.4}$) to 1,745.7 kg ha⁻¹ among diallel crosses (TAB. 1). For popping expansion (PE), the mean values were from 33.4 ($P_{9.27} \times P_{8.4}$) to 15.4 mL g⁻¹. Considering the means for the check hybrid IAC 112 (3,175.9 Kg ha⁻¹ for GY and 32.4 mL g⁻¹ to PE (TAB. 1), the hybrid combination $P_{9.27} \times P_{8.4}$ was considered outstanding. It suggested that lines $P_{9.27}$ (set 1-IAC 112) and $P_{8.4}$ (set 2-Zaeli) may be promising per se, or their combination resulted in positive heterosis and favorable allelic complementation.

Regarding to the partial resistance to southern rust, the hybrids ranged from 10.4 ($P_{9.24} \times P_{8.5}$) to 24.4% ($P_{9.20} \times P_{8.6}$) in severity at 92 days, and from 57.9 ($P_{9.22} \times P_{8.2}$) to 191.4 ($P_{9.20} \times P_{8.6}$) in AUDPC (TAB. 1). When compared to check IAC 112, the means of hybrids corresponds to moderately susceptible to susceptible popcorn cultivars. This revealed some possibilities with regards to the improvement of partial resistance to southern rust using this germplasm, especially if additive genes would be acting. Mean values for hybrid combinations per line are presented in the Table 1. In this sense, it was possible to emphasize the lines $P_{8.5}$ and $P_{8.2}$ from set 2-Zaeli to Pp resistance.

Hybrid effects were significant ($p < 0.10$) for GY and AUDPC (TAB. 2). It indicated that some hybrids were superior for GY, and it is possible to identify lines which were combined in outstanding hybrids. Concerning to AUDPC, which integrates the Pp evaluations, some hybrids were able to reduce the disease development over the season, and therefore they had higher levels of partial resistance. This is a favorable condition for plant breeding. Hybrid effects were not detected ($p > 0.10$) for PE and severity at 92 d (TAB. 2).

General combining ability was significant ($p < 0.10$) for GY, PE, and partial resistance to Pp in the inbred lines from set 2-Zaeli (TAB. 2). Therefore, it is possible to assume that additive gene effects are important for these traits, and suggests that alleles from some lines are outstanding compared to the other ones. Specific combining ability - from nonadditive gene effects from dominance or epistasis - was detected for GY ($p < 0.10$). This indicated that some hybrids are heterotic and had good allelic complementation.

For PE and partial resistance to Pp, the relative importance of general combining ability was higher than the importance of specific combining ability (TAB. 2). This indicates that additive genetic variability had larger breeding possibilities than nonadditive variability, and therefore the intrapopulation breeding based on recurrent selection are recommendable.

The estimates of general combining ability effects (\hat{g}_j) for the lines of set 2-Zaeli are presented at Table 3. For GY and PE, positive \hat{g}_j 's larger than its standard deviation (SD) was considered relevant. For partial resistance to Pp, negative \hat{g}_j 's larger than its SD was relevant. None of lines had favorable \hat{g}_j in respect to GY and PE. The lines $P_{8.1}$ (for GY) and $P_{8.2}$ (for PE) had negative relevant \hat{g}_j , and might reduce GY in their hybrids. In fact, since none of other lines had relevant \hat{g}_j , it denoted that lines from set 2-Zaeli are quite similar concerning its genetic frequencies. This also occurred for lines from set 1-IAC 112 ($p > 0.10$ for general combining ability). Therefore, it ratified that self-pollination of commercial hybrids are not appropriate

Table 1 - Means for grain yield^a (kg ha⁻¹), popping expansion^b (mL g⁻¹), southern rust severity at 92 days after sowing^c (%), and area under the disease progress curve^d (AUDPC) in diallel crosses obtained by circulant partial scheme

Inbred lines	Set 2-Zaeli					Mean ¹
	1' (P _{8.5})	2' (P _{8.2})	3' (P _{8.4})	4' (P _{8.1})	5' (P _{8.6})	
Set 1-IAC112						
	2,697.5 a	2,453.7	2,688.5			2,613.2
1 (P _{9.22})	23.4 b	19.4	25.9			22.9
	10.7 c	15.4	12.2			12.8
	83.5 d	57.9	93.8			78.4
2 (P _{9.27})		3,112.0	3,200.8	1,894.4		2,735.7
		24.4	33.4	26.3		28.1
		10.8	19.3	19.2		16.5
		83.1	139.0	150.0		124.0
3 (P _{9.1})			2,299.4	2,457.4	1,936.4	2,231.1
			26.9	27.8	25.4	26.7
			13.1	22.2	19.0	18.1
			99.1	151.9	151.2	134.1
4 (P _{9.24})	2,721.8			1,745.7	2,925.5	2,464.3
	23.9			21.1	28.2	24.4
	10.4			21.0	19.9	17.1
	85.5			144.7	145.1	125.1
5 (P _{9.20})	2,931.6	2,724.5			2,095.1	2,583.7
	24.8	15.7			27.6	22.7
	11.9	17.4			24.2	17.8
	88.7	86.0			191.4	122.0
Mean ²	2,783.6	2,763.4	2,729.6	2,032.5	2,319.0	
	24.0	19.8	28.7	25.1	27.1	
	11.0	14.5	14.9	20.8	21.0	
	85.9	75.6	110.7	148.9	162.5	
Check hybrid (IAC 112)			3,175.9			
			32.4			
			6.2			
			43.03			

¹Means of hybrids combinations per line from set 1-IAC 112. ²Means of hybrids combinations per line from set 2-Zaeli

to obtain lines with relative broad genetic base. In our understanding, this meet with expectation since two to four inbred lines are employed to obtain the commercial hybrids. In this instance, three lines are combined in IAC 112, the base material for the lines of set 1.

The lines P_{8.5} and P_{8.2} were the ones which most contributed to partial resistance to Pp. Due to the relevant negative \hat{g}_j , these lines decreased the overall mean of diallel, reducing final severity level and Pp progress on their hybrids. Since P_{8.2} had negative \hat{g}_j for PE, only P_{8.5} was recommended to form breeding populations for partial resistance to Pp.

Regarding to the specific combining ability for GY, two hybrid combinations with outstanding estimates of specific combining ability effects were identified (TAB. 4). These two combinations were P_{9.1} × P_{8.1} (s_{ij} = 616.1 kg ha⁻¹) and P_{9.24} × P_{8.6} (s_{ij} = 490.5 kg ha⁻¹). In conclusion, these combinations overcome its expected performance if considering the general combining ability of their lines. The hybrid combination P_{9.1} × P_{8.1} included P_{8.1} (with negative \hat{g}_j for GY), and therefore was discarded to be exploited by breeding. The hybrid P_{9.24} × P_{8.6}, however, is promising for interpopulational breeding for GY.

Table 2 - Analyses of variance and circulant partial diallel analyses of Kempthorne and Curnow (1961) for grain yield (GY, kg ha⁻¹), popping expansion (PE, mL g⁻¹), southern rust severity at 92 days after sowing (Sev92), and area under the disease progress curve (AUDPC)

Sources of variation	DF	Mean squares			
		GY	PE	sev92	AUDPC
Hybrids	14	612580.6*	51.4 ^{ns}	64.3 ^{ns}	4374.3*
General combining ability (GCA) set 1-IAC 112	4	146521.8 ^{ns}	46.1 ^{ns}	8.7 ^{ns}	950.4 ^{ns}
General combining ability (GCA) set 2-Zaeli	4	1104627.7*	101.3*	180.6*	13536.9*
Specific combining ability (SCA)	6	595295.0*	21.1 ^{ns}	24.6 ^{ns}	556.2 ^{ns}
Blocks	2	202948.0 ^{ns}	15.3 ^{ns}	784.7*	22343.1*
Error	28	204218.9	38.2	38.7	2249.3
Overall mean		2525.6	24.9	16.4	116.7
Coefficient of variation		17.9	24.8	37.8	40.6
%SS GCA set 1-IAC112		7	26	4	6
%SS GCA set 2-Zaeli		52	57	80	88
%SS SCA		42	18	16	5

SS: Sum of squares. * p < 0.10 by F-test. ^{ns} p > 0.10 by F-test**Table 3** - Estimates of general combining ability effects in the lines of set 2-Zaeli (\hat{g}_j) for grain yield (GY, kg ha⁻¹), popping expansion (PE, mL g⁻¹), southern rust severity at 92 days after sowing (Sev92), and area under the disease progress curve (AUDPC)

Inbred lines	\hat{g}_j			
	GY	PE	Sev92	AUDPC
1' (P _{8.5})	302.8	0.5	-6.1	-29.1
2' (P _{8.2})	226.8	-5.6	-2.4	-46.5
3' (P _{8.4})	212.4	2.8	-0.8	3.9
4' (P _{8.1})	-566.3	-0.8	5.1	33.2
5' (P _{8.6})	-175.7	3.1	4.2	46.2
SD $\hat{g}_j - \hat{g}_i$	367.4	3.5	4.7	39.6

SD: Standard deviation. Relevant estimates are highlighted in bold

Table 4 - Estimates of specific combining ability effects (s_{ij}) between lines of the two sets for grain yield (GY, kg ha⁻¹)

Inbred lines	Set 2-Zaeli				
	1' (P _{8.5})	2' (P _{8.2})	3' (P _{8.4})	4' (P _{8.1})	5' (P _{8.6})
Set 1-IAC112					
1 (P _{9.22})	28.8	-139.0	110.2		
2 (P _{9.27})		107.1	210.33	-317.4	
3 (P _{9.1})			-320.64	616.1	-295.5
4 (P _{9.24})	-191.8			-298.7	490.5
5 (P _{9.20})	163.0	31.9			-195.0

Relevant estimates are highlighted in bold

Both additive and nonadditive gene effects were important for GY. Their relative importance was similar on genetic variability, indicating that the exploitation of both gene effects is appropriate using recurrent reciprocal selection. In this context, to improve many breeding populations simultaneously are recommended before obtaining lines. In this manner, high-yielding, vigorous, and heterotic lines are supposed to be obtained. This strategy was supported by results from Scapim et al. (2002) and Simon et al. (2004), which found smaller inbreeding depression in improved populations. In fact, this strategy also seems to be appropriate to obtain popcorn genotypes with resistance to northern leaf blight hybrids (VIEIRA et al., 2009c). Since inbreeding depression levels are low for PE (SIMON et al., 2004), these improved populations should have genetic variability and high PE levels (superior than 28 g mL⁻¹).

When a trait is major conditioned by additive genes and heterotic effects are not strongly important, either phenotypic or genotypic means among parents are a good predictor of hybrids' performance, and screening or evaluating lines per se is crucial for maize breeding programmes. In this sense, both PE and Pp are included.

Additive gene effects have been more important than nonadditive effects for PE in landraces and open-pollinated popcorn varieties from South American germplasm, in South and Central Brazil (ANDRADE et al., 2002; FREITAS JÚNIOR et al., 2006; MIRANDA et al., 2008; PINTO et al., 2007; SCAPIM et al., 2002; VIEIRA et al., 2009a). For popcorn inbred lines, these predominance was reported by Larish and Brewbaker (1999) in U.S. germplasm, and Pinto et al. (2007a, 2007b), in Brazilian adapted germplasm. Our study corroborated with other studies and reinforces findings previously obtained.

Concerning to the partial resistance to Pp, no studies were reported previously in popcorn germplasm worldwide. The predominance of additive genes conditioning this trait was found in common corn (SILVA et al. 2001). Slow-rusting resistance was also found in common maize (BRUNELLI et al., 2004; CHÁVEZ-MEDINA et al., 2007). Nonadditive gene effects, however, were also reported (JINES et al., 2006; SILVA et al., 2001). In popcorn germplasm, Vieira et al. (2009b) reported the major importance of additive genes for partial resistance to northern leaf blight (*Exserohilum turcicum*) and gray leaf spot (*Cercospora* spp.) in popcorn inbred lines. For resistance to Pp in popcorn, this study firstly reported the major action of additive gene effects. Concerning this study, it was possible to affirm that the inheritance of slow-rusting resistance to Pp (partial resistance) was from additive genes. Suitable level of partial resistance provides a durable and no race-specific resistance, and since many

racess occur simultaneously the partial resistance is a important tool for breeding programmes.

The importance of promising inbred lines with resistance to southern rust resistance is emphasized when considering the significant correlations between yield losses and southern rust severity (CASTELLANOS et al., 1998). In this respect it is believed that popcorn breeding including P_{8.5} are useful.

Conclusions

1. General combining ability were more important than specific combining ability for grain yield, popping expansion, and partial resistance to southern rust, indicating that additive gene are predominant, and therefore recurrent selection is recommendable;
2. The inbred line P_{8.5} is recommended to form breeding populations for partial resistance to southern rust;
3. The hybrid combination P_{9.24} × P_{8.6} was promising for interpopulational breeding for GY.

Acknowledgements

The authors thank CNPq - National Counsel of Technological and Scientific Development, Brazil; and Araucaria Foundation, from the Parana State Government, Brazil, for partially fund this work.

References

- ANDRADE, R. A. *et al.* Análise dialélica da capacidade combinatória de variedades de milho-pipoca. **Acta Scientiarum. Agronomy**, v. 24, n. 05, p. 1197-1204, 2002.
- ARNHOLD, E. *et al.* Evaluation of top-cross popcorn hybrids using mixed linear model methodology. **Chilean Journal of Agricultura Research**, v. 69, n. 01, p. 46-53, 2009.
- BRUNELLI, K. R.; SILVA, H. P.; ARANHA, L. E. C. Mapeamento de genes de resistência quantitativa a *Puccinia polysora* em milho. **Fitopatologia Brasileira**, v. 27, n. 02, p. 134-140, 2004.
- CAMPBELL, C. L.; MADDEN, L. V. **Introduction to plant disease epidemiology**. New York : John Wiley, 1990. 532 p.
- CASTELLANOS, J. S.; HALLAUER, A. R. CORDOVA, H. S. Relative performance of testers to identify elite lines of corn (*Zea mays* L.) **Maydica**, v. 43, n. 03, p. 217-226, 1998.
- CHÁVES-MEDINA, J. A.; LEYVA-LÓPEZ, N. E.; PATAKY, J. K. Resistance to *Puccinia polysora* in maize accessions. **Plant Disease**, v. 91, n. 11, p. 1489-1495, 2007.

- FREITAS JÚNIOR, S. P. *et al.* Capacidade combinatória em milho-pipoca por meio de dialelo circulante. **Pesquisa Agropecuária Brasileira**, v. 41, n. 11, p. 1599-1607, 2006.
- JINES, M. P. Mapping resistance to southern rust in tropical × temperate maize recombinant inbred topcross population. **Theoretical and Applied Genetics**, v. 114, n. 04, p. 659-667, 2006.
- JONES, D. F. **The effects of inbreeding and crossbreeding upon development**. Connecticut: Connecticut Agricultural Experimental Station Bulletin, 1918. 100 p.
- KEMPTHORNE, O.; CUNOW, R. N. The partial diallel cross. **Biometrics**, v. 17, n. 01, p. 229-250, 1961.
- LARISH, L. L. B.; BREWBAKER, J. L. Diallel analyses of temperate and tropical popcorns. **Maydica**, v. 44, n. 04, p. 279-284, 1999.
- MIRANDA, G. V. *et al.* Genetic variability and heterotic groups of Brazilian popcorn populations. **Euphytica**, v. 162, n. 02, p. 431-440, 2008.
- PINTO, R. J. B. *et al.* Análise dialélica parcial de linhagens de milho-pipoca. **Revista Brasileira de Milho e Sorgo**, v. 06, n. 03, p. 323-335, 2007.
- PINTO, R. J. B. *et al.* Análise dialélica de linhagens de milho-pipoca. **Revista Ceres**, v. 54, n. 315, p. 471-477, 2007.
- RODRIGUES-ARNON, R.; SCOTT, G. E.; KING, S. B. Maize yield losses caused by southern corn rust. **Crop Science**, v. 20, n. 03, p. 812-814, 1980.
- SANTOS, F. S. *et al.* Genetic gain prediction of the third recurrent selection cycle in a popcorn population. **Acta Scientiarum. Agronomy**, v. 30, n. 05, p. 651-655, 2008.
- SCAPIM, C. A. *et al.* Análise dialélica e heterose de populações de milho pipoca. **Bragantia**, v. 61, n. 03, p. 219-230, 2002.
- SCAPIM, C. A. *et al.* Correlations between the stability and adaptability statistics of popcorn cultivars. **Euphytica**, v. 174, n. 02, p. 209-218, 2010.
- SCHMILDT, E. R. *et al.* Avaliação de métodos de correção do estande para estimar a produtividade em milho. **Pesquisa Agropecuária Brasileira**, v. 36, n. 08, p. 1011-1018, 2001.
- SHURTLEFF, M. C. **Compendium of Corn Diseases**. St. Paul: APS Press, 1992. 105 p.
- SILVA, H. P. *et al.* Capacidade de combinação e heterose para resistência a *Puccinia polysora* Underw. em milho. **Scientia Agricola**, v. 58, n. 04, p. 777-783, 2001.
- SILVA, T. R. *et al.* Genetic divergence in popcorn genotypes using microsatellites in bulk genomic DNA. **Crop Breeding and Applied Biotechnology**, v. 09, n. 01, p. 31-36, 2009.
- SIMON, G. A. *et al.* Depressão por endogamia em populações de milho-pipoca. **Bragantia**, v. 63, n. 01, p. 55-62, 2004.
- TRINDADE, A. P. *et al.* Genetic diversity underlying heterotic groups of popcorn lines of tropical and temperate germplasm determined by SSR markers. **Electronic Journal of Biotechnology**, v. 13, n. 01, p. 1-9, 2010.
- VIEIRA, R. A. *et al.* Heterotic parametrization for economically important traits in popcorn. **Acta Scientiarum. Agronomy**, v. 31, n. 03, p. 411-419, 2009a.
- VIEIRA *et al.* Diallel analysis of leaf disease resistance in inbred Brazilian popcorn cultivars. **Genetics and Molecular Research**, v. 08, n. 04, p. 1427-1436, 2009b.
- VIEIRA, R. A. *et al.* Resistência de híbridos de milho-pipoca a *Exserohilum turcicum*, agente causal da helmintosporiose do milho. **Scientia Agraria**, v. 10, n. 05, p. 391-395, 2009c.
- ZIEGLER, K. E.; ASHMAN, B. POPCORN. In: Hallauer, A. R. **Specialty corns**. Boca Raton: CRC Press, 1994. cap. 7, p. 189-223.
- ZUMMO, N. Components contributing to partial resistance in maize to *Puccinia polysora*. **Plant Disease**, v. 72, n. 02, p. 157-160, 1988.