



Revista Ciência Agronômica

ISSN: 0045-6888

ccarev@ufc.br

Universidade Federal do Ceará
Brasil

Gonzatto Maciel, Caciara; Walker, Clair; Feliciano dos Santos, Ricardo; Brião Muniz,
Marlove Fátima; Lemos Brum, Daniele

Fusarium oxysporum and *F. verticillioides* associated with damping-off in *Pinus* spp.

Revista Ciência Agronômica, vol. 48, núm. 1, enero-marzo, 2017, pp. 134-141

Universidade Federal do Ceará
Ceará, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=195347767015>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

***Fusarium oxysporum* and *F. verticillioides* associated with damping-off in *Pinus* spp.¹**

Fusarium oxysporum e *F. verticillioides* causando damping-off em *Pinus* spp.

Caciara Gonzatto Maciel^{2*}, Clair Walker², Ricardo Feliciano dos Santos³, Marlove Fátima Brião Muniz² and Daniele Lemos Brum²

ABSTRACT - Occurrence of *Fusarium* spp. is one of the problems, most limiting to growth of seedlings, in nurseries. This pathogen can be transmitted via seeds and causes damages to the seedlings during pre- and post-emergence stages. The present study aimed to identify *Fusarium* spp. at the species level based on morphological and molecular characteristics and to verify the pathogenicity of these isolates in seeds lots of *Pinus elliottii* and *P. taeda*. For this, we used two *Fusarium* isolates and five lots of *Pinus* spp. seeds. Morphological characterization was performed based on a key, specific to *Fusarium* spp. identification, whereas, molecular identification was carried out by amplification and sequencing of the regions from internal transcribed spacer (ITS) and the elongation factor 1- α (*tef1*). The pathogenicity test was conducted through the contact of the seeds with fungal culture for 48 h, followed by sowing them in sand. The variables evaluated were emergency speed index, percentage of emergency, non-emergency seeds, symptomatic seedlings, and seedling damping-off. One isolate, F1UFSM, was identified as *F. verticillioides* and another isolate, F2UFSM, was identified as *F. oxysporum*. Both the isolates were pathogenic to the seeds of *Pinus* spp., causing a reduction in the percentage of emergence and seedling damping-off.

Key words: ITS. Elongation factor (*tef1*). *Pinus taeda*. *Pinus elliottii*.

RESUMO - A ocorrência de *Fusarium* spp. é um dos problemas mais limitantes em viveiros, esse patógeno pode ser transmitido via semente e causar danos em pré e pós-emergência. O presente trabalho objetivou identificar isolados de *Fusarium* spp. a nível de espécie e verificar a patogenicidade desses isolados em lotes de sementes de *Pinus elliottii* e *P. taeda*. Foram utilizados dois isolados do patógeno e cinco lotes de sementes de *Pinus* spp. Para caracterização molecular, sequenciaram-se as regiões genômicas: ITS e fator de elongação 1- α (*tef1*), para morfologia dos isolados utilizaram-se chaves de identificação específica para o gênero *Fusarium*. O teste de patogenicidade foi realizado através do contato das sementes com a cultura fúngica por 48 horas, seguida de semeadura em areia. As variáveis avaliadas foram: índice de velocidade de emergência, porcentagem de emergência, plântulas sintomáticas e tombamento de plântulas. O isolado F1UFSM foi identificado como *Fusarium verticillioides* e o F2UFSM como *F. oxysporum*. Ambos foram patogênicos a *P. taeda* e *P. elliottii*, causando redução no percentual de emergência e tombamento de plântulas.

Palavras-chave: ITS. Fator de elongação (*tef1*). *Pinus taeda*. *Pinus elliottii*.

DOI: 10.5935/1806-6690.20170015

*Autor para correspondência

¹Recebido para publicação em 30/09/2015 a 12/04/2016

Parte da Tese de Doutorado em Engenharia Florestal da primeira autora, apresentada à Universidade Federal de Santa Maria/UFSM

²Departamento de Defesa Fitossanitária, Universidade Federal de Santa Maria, Campus Universitário, Centro de Ciências Rurais Prédio 42, 2º Piso, Sala 3223A, Santa Maria-RS, Brasil, 97.105-900, caciaraconzatto@gmail.com, clairwalker@gmail.com, marlovemuniz@yahoo.com.br, daniellemosbrum@yahoo.com.br

³Departamento de Fitopatologia, Escola Superior de Agricultura "Luiz de Queiroz" USP/ESALQ/USP, Piracicaba-SP, Brasil, ricardofelicianodossantos@gmail.com

INTRODUCTION

The occurrence of *Fusarium* spp. is one of the problems that are the most limiting to the growth of seedlings in nurseries. This cosmopolitan pathogen is a native inhabitant of soil and is difficult to control. Several species of *Fusarium* are associated with seed rot and damping-off of seedlings of *Pinus* spp., for example, *F. sambucinum* in *P. elliottii* (MACIEL *et al.*, 2013), *F. circinatum* in *P. radiata* and *P. pinaster* (LANDERAS *et al.*, 2005) as well as *F. circinatum* in *P. maximinoi*, *P. patula* and *P. tecunumanii* (STEENKAMP *et al.*, 2012), and *P. taeda* and *P. elliottii* (LORI; SALERNO, 2003). Pfenning *et al.* (2014) identified *F. circinatum* associated with *Pinus* in Brazil, where this pathogen is considered a quarantine fungus.

F. oxysporum and *F. verticillioides* were observed to be pathogenic to seedlings of *P. nigra* in northwestern Spain, causing damping-off during pre-emergence stage and, therefore, in the reduction of germination potential (MARTÍN-PINTO; PAJARES; DÍEZ, 2008). Because of the diversity of hosts that are compromised by *Fusarium*, it is necessary to identify the pathogen at the species level in order to facilitate the applicability of control strategies.

In this context, O'Donnell (2000) and Leslie and Summerell (2006) suggested three features of species for identification of *Fusarium* spp.: morphological, based on the similarity of the observed characters, called morphological markers, biological, based on the sexual compatibility between members of the same species, and phylogenetic, based on the analysis of gene sequences. DNA analysis is an effective tool, rapid and accurate to detection of fungal pathogens to species or strain level (TSUI; WOODHALL; CHEN, 2011), and has been complementing the morphological characterization, giving more credibility to the process of identifying the species.

Considering the above background, the objective of the present study was to identify *Fusarium* spp. at the species level based on morphological and molecular characteristics and to verify the pathogenicity of these isolates in seeds lots of *P. elliottii* and *P. taeda*.

MATERIAL AND METHODS

Seeds of *P. elliottii* and *P. taeda* (harvested in 2013) used in this study had their origin in the municipality of Ijuí (28°23'16" S e 53°54'54" W), located in the northwest region of Rio Grande do Sul state, Brazil. Five lots were used: two of *P. elliottii* (Lot 1 and Lot 2) and two of *P. taeda* (Lot 3 and Lot 4) and a fifth lot *P. elliottii* (SP1) was from São Paulo city, São Paulo. For isolation of the pathogen from the lots, seeds were subjected to the "blotter test",

whereby, they were incubated for seven days at 25 °C under a 12 h light/dark photoperiod; the morphological characteristics of the putative *Fusarium* colonies on the seeds were observed after this period. The putative colonies were then subcultured on Petri dishes containing potato dextrose agar (PDA) medium supplemented with 0.5 g·L⁻¹ streptomycin sulfate and incubated under the same conditions mentioned above. The characteristics of the resulting colonies were compared to those described earlier for the genus *Fusarium* (GERLACH; NIRENBERG, 1982; LESLIE; SUMMERELL, 2006). Monosporic cultures were obtained (Table 1) according to the methodology of Alfenas and Mafia (2007) and stored in the mycological collection of the Laboratory of Plant Pathology, Universidade Federal Santa Maria, Santa Maria, RS, Brazil.

For molecular characterization of the fungal isolates, mycelium and spores were collected from the cultures grown on PDA medium for two weeks under the conditions described in the previous section. DNA from the pathogen was extracted by the cetyltrimethylammonium bromide (CTAB) method (DELLAPORTA; WOOD; HICKS, 1983). Samples of the extracted genomic DNA were used in polymerase chain reaction (PCR) for amplification of internal transcribed spacer (ITS) region of rDNA using the primer pair, ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (WHITE *et al.* 1990.) and the region of elongation factor 1 alpha (*tef1*) using the primer pair EF1-T (5'-ATGGGTAAGGARGACAAGAC-3') and EF1-1567R (5'-ACHGTRCCRATACCACCRATCTT-3') (REHNER; BUCKLEY, 2005). Each PCR mixture contained approximately 1 µL DNA, 10 µL 5X GoTaq Reaction Buffer (Promega, EUA), 1 µL dNTPs mixture, 1 µL of each primer, 0.2 µL GoTaq DNA polymerase (Promega, EUA), and autoclaved MiliQ water to make the volume to 50 µL. PCR was performed in a GeneAmp PCR System 2400 (Perkin Elmer, EUA) under the following cycling conditions: 94 °C for 2 min, 40 cycles of denaturation at 94 °C for 30 s, annealing at 50 °C for 2 min, and elongation at 72 °C for 1 min followed by a final extension at 72 °C for 4 min. At the end of the reaction, the PCR products were stored at 4 °C. A negative control, without DNA, was included in the PCR amplifications. After electrophoresis on a 0.8% agarose gel in 1X TAE buffer (0.04 M Tris-acetate, 1 mM EDTA) and staining of the gel with ethidium bromide (1 mg L⁻¹), the amplified fragments were visualized under ultraviolet light. A 1 kb Plus DNA ladder (Invitrogen, USA) was used as the molecular weight marker. The PCR products obtained were purified following the protocol described by Schmitz and Riesner (2006) using polyethylene glycol 6000 (PEG 6000).

Table 1 - Details of isolates *Fusarium* spp. utilized in this study

Isolate	Origin	Collection period (month/year)	Access code in GenBank	
			ITS	TEF1
F1UFSM	Seeds of <i>P. elliottii</i>	04/2013	KX710194	KU170704
F2UFSM	Seeds of <i>P. taeda</i>	04/2013	KX710195	KU170705

The obtained sequences were compared to those of *Fusarium* and *Gibberella* spp. (teleomorph), available in the GenBank. The sequences from the GenBank that showed the highest scores were selected and aligned with the sequences of the amplified products by the ClustalW algorithm. Furthermore, phylogenetic analysis was conducted, adopting the Neighbor-joining method with 1000 replicates in the MEGA program version 4 (TAMURA; DUDLEY; NEI, 2007). The similarity of the nucleotide sequences between the isolates was calculated using the Basic Local Alignment Search Tool-BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

For morphological characterization, sterilized leaf fragments (~3–5 mm²) of carnation (*Dianthus caryophyllus*) were infected with colonies of the pure isolates on 2% culture medium agar (NELSON; TOUSSON; MARASAS, 1983; VENTURA, 1999) and incubated for 25 days at 25 ± 2 °C under 12 h light/dark photoperiod, for scaling and photography of reproductive structures. Thirty conidia were evaluated in the isolates of *Fusarium* spp. by measuring their length and width. For staining of the culture, mycelia were aliquoted in test tubes containing PDA medium and incubated for 10 days under the conditions specified above. The color was determined visually (front plate) and compared with that described in the literature (LESLIE; SUMMERELL; 2006; NELSON; TOUSSON; MARASAS, 1983).

To determine the average mycelial growth of the isolates, we placed agar-solidified culture medium disks (12 mm) derived from the plates of the pure cultures in the center of a Petri dish containing PDA medium. The plates were incubated under the conditions specified earlier and the mycelial growth was observed by measuring the diameter of the colony every 24 h with the aid of a digital caliper, during seven days. The measurements were taken in two diametrically opposed directions. At the end of this experiment, the sporulation of each isolate was determined, by adding 20 mL of sterile distilled water to each Petri dish and scraping with the handle of a Drigalski spatula; the suspension was filtered through gauze and the concentration (conidia mL⁻¹) of conidia was estimated using a Neubauer chamber. Five replicate plates (90 mm)

of each isolate were used for determining the sporulation. The mycelial growth were submitted to regression analysis using the statistic program System Analysis of Variance for Balanced Data (SISVAR 5.3) (FERREIRA, 2010).

The pathogenicity of F1UFSM and F2UFSM isolates in *Pinus* seeds was determined by inoculating the seeds of five lots of *P. elliottii* (Lot1, Lot2 and SP1) and *P. taeda* (Lot3 and Lot4) with the pathogens. The seeds used in the tests remained stored in the freezer (-18 °C) for two weeks for breaking their dormancy (BRASIL, 2009). For testing the pathogenicity, initially, the seeds were sterilized with a solution of 70% (v/v) alcohol for 30s and then with a solution of sodium hypochlorite (1% v/v) for 1 minute. They were subsequently washed with sterile distilled water and dried on sterile filter paper. Each treatment used 100 seeds, divided into four replicates of 25 seeds each.

After the incubation period of the fungus (seven days at 25 ± 2 °C with a photoperiod of 12 h light/dark), inoculation was carried out by keeping the seeds in contact with the fungal culture for 48 h at 25 ± 2 °C and 12 h light/dark photoperiod. For the control, seeds were exposed only to the PDA medium under the same conditions. After inoculation, the test of emergence in sand was performed, in which the seeds were placed in plastic boxes (11 × 11 × 3.5 cm) containing sifted sand as a substrate; the sand was sterilized by autoclaving for 2 h (with interval of 24 h) 1 atm and 120 °C. The material remained incubated in a temperature-controlled room with a temperature of 25 ± 2 °C and with manual irrigation where necessary. Were used four plastic boxes per treatment, and each treatment corresponds to inoculation of *Fusarium* spp. (F1UFSM or F2UFSM) on lots of *Pinus* spp. seeds (Lot1, Lot2, Lot3, Lot4 or SP1).

The variables evaluated were: a) emergence speed index (ESI): daily count of emerged seedlings, considered when the hypocotyls were bigger than 1.0 cm, determining the ESI through the equation suggested by Maguire (1962); b) seedling emergence: counting the number of seedlings at 28 days; c) abnormal seedlings symptomatic: seedling with symptoms caused by the *Fusarium* were checked; d) non-emergence seeds: count of the seeds

with rotted aspect and of those that had not started the germination process. For all variables, except the ESI, the results were expressed as percentages. When the presence of damping-off was detected, the seedlings were collected and incubated in a moist chamber or placed in Petri dishes with PDA culture medium, with the goal of determining whether the damage was caused by the inoculated fungus, and then re-isolation was performed.

The trials were conducted in a completely randomized design, treatments with four repetitions, and each repetition consisting of one plastic boxes containing 25 seeds. Means comparison was done by Tukey's test at 5% probability; the software used was SISVAR 5.3 (FERREIRA, 2010).

RESULTS AND DISCUSSION

In this study, the isolates F1UFSM and F2UFSM were pathogenic to *Pinus*'s seeds. The seedling symptoms began approximately 14 days after inoculation and were described as follows: seed rot and damping off in pre-emergence and post emergence of seedlings (Figure 1).

Analyzing the values of ESI lots of study, with and without inoculation of *Fusarium* spp., notes that those who remained in contact with the pathogen before sowing

had a lower rate of emergence speed. This means that the average number of seedlings per day decreased due to colonization of seed by the pathogen, thus causing an increase in the inequality of emergence of seeds, associated with reduced vigor. According to the data presented in Table 2 reveals that the isolates were pathogenic seeds of *Pinus* spp., directly interfering on seedling emergence variable, except for the seeds lot of *P. elliottii* (SP1), where no significant difference was for this variable. In this lot (*P. elliottii* - SP1) the difference was observed for the variables symptomatic seedlings and damping-off in post-emergence when compared with the control treatment. This result may indicate that the presence of the pathogen did not affect the initial formation processes (germination and emergence), but acted in a negative way during seedling development.

The fusariosis, in the seedlings of different *Pinus* spp., is reported to be an important disease in the nursery causing discoloration of the needles, drying of the apical parts, browning of the roots, growth stagnation (GRIGOLETTI JUNIOR; AUER, 2006), wilting and low seedling-survival rate (MACIEL *et al.*, 2013.), and rot and reduction in the root development (OCAMB; JUZWIK; MARTIN, 2002).

DNA of *Fusarium* spp. isolates was amplified with primers and sequenced. For the *tefl* and ITS regions, 677 and 678 bp fragments were amplified, respectively. Both the analyzed regions were effective in the classification of the isolates belonging to the same clade based on the other sequences of *Fusarium* spp. deposited in the GenBank having high bootstrap values. To construct the phylogenetic tree, based on both the ITS and *tefl* amplicons, the GenBank accessions that showed the highest coverage ($\geq 95\%$) and similarity were chosen ($\geq 98\%$) (Figure 2). The F1UFSM isolate was allocated in the same clade with *F. verticillioides* (anamorph *Gibberella moniliformis*), with bootstrap values of 98 and 87, while F2UFSM was grouped with *F. oxysporum* with bootstrap values of 97 and 84, respectively, for ITS and *tefl*. According to Geiser *et al.* (2004), *tefl* has become the most common marker, since it is highly informative among the *Fusarium* single copy genes. Schoch *et al.* (2012) emphasized the importance of specific markers for *Fusarium*, such as elongation factor 1 - α and suggested the ITS region as a potential universal code for the identification of fungi. The ITS region was considered efficient in the identification and separation of *Fusarium* (MENEZES *et al.*, 2010). Molecular markers, ITS and *tefl*, are considered fast and low cost practical tools for efficient discrimination and identification of *Fusarium* spp. (ARIF; CHAWLA; ZAIDI, 2012).

According to macroscopic and microscopic characteristics of the fungal colonies, F1UFSM was allocated to the section *Liseola* while F2UFSM was framed in the section *Elegans*; these sections are distinguished

Figure 1 - Symptoms post-emergence observed in seedlings of *Pinus* spp. after the inoculation with *Fusarium* sp.: (A) *P. elliottii* Lot1 x F1UFSM; (B) *P. elliottii* Lot2 x F1UFSM; (C) *P. elliottii* SP1 x F1UFSM (C); and (D) without inoculation - control




Table 2 - Mean values of emergence speed index (ESI), emergency (E), non-germinated seeds (NGS) and symptomatic seedlings (SS) of *Pinus* spp. seeds (Lot1, Lot2, Lot3, Lot4 and SP1) after inoculation with *Fusarium* spp. (F1UFMS and F2UFMS)

Lot x isolates	Variables			
	ESI	E (%)	NGS (%)	SS (%)
<i>P. elliottii</i> Lot1 X F1UFMS	5.33 b*	38 b	62 b	50 a
<i>P. elliottii</i> Lot1 X F2UFMS	4.73 b	36 b	64 b	63 a
<i>P. elliottii</i> Lot1 - Control	16.3 a	78 a	22 a	11 b

Lot x isolates	Variables			
	ESI	E (%)	NGS (%)	SS (%)
<i>P. elliottii</i> Lot2 X F1UFMS	9.3 b	50 b	50 b	75 a
<i>P. elliottii</i> Lot2 X F2UFMS	5.4 b	39 b	61 b	57 a
<i>P. elliottii</i> Lot2 - Control	15.3 a	75.0 a	25 a	26 b

Lot x isolates	Variables			
	ESI	E (%)	NGS (%)	SS (%)
<i>P. taeda</i> Lot3 X F1UFMS	1.5 b	18 b	82 b	10 a
<i>P. taeda</i> Lot3 X F2UFMS	1.5 b	12 b	88 b	13 a
<i>P. taeda</i> Lot3 - Control	13.8 a	73.0 a	27 a	9.0 a

Lot x isolates	Variables			
	ESI	E (%)	NGS (%)	SS (%)
<i>P. taeda</i> Lot4 X F1UFMS	2.05 b	17.0 b	83 b	34 a
<i>P. taeda</i> Lot4 X F2UFMS	3.02 b	19 b	81 b	39 a
<i>P. taeda</i> Lot4 - Control	9.65 a	64.0 a	36.0 a	21 a

Lot x isolates	Variables			
	ESI	E (%)	NGS (%)	SS (%)
<i>P. elliottii</i> SP1 X F1UFMS	18.3 a	81 a	19 a	32 a
<i>P. elliottii</i> SP1 X F2UFMS	19.4 a	86 a	14 a	16 ab
<i>P. elliottii</i> SP1 - Control	13.6 a	81 a	19 a	0.0 b


*Means followed by the same letter in the column do not differ by Tukey test at 5% significance. Where: F1UFMS and F2UFMS are isolates of *Fusarium* spp. obtained from seeds of *P. elliottii* and *Pinus taeda*, respectively

according to the morphological similarities of the species (GERLACH; NIRENBERG, 1982; NELSON; TOUSSON; MARASAS, 1983) and provide a direction for the sequence during the identification process. The isolates F1UFMS and F2UFMS were identified as *F. verticillioides* (synonym *F. moniliforme*) and *F. oxysporum*, respectively. However, the confirmation of species was made possible only by the molecular determination (Figure 2). According to Leslie and Summerell (2006), the combinations of phylogenetic and biological concepts have a high potential for consistent characterization of the genus *Fusarium*. Furthermore, the use of molecular techniques for DNA analysis has allowed the development of fast, sensitive, and specific methods, complementary to the diagnosis of pathogens, for complementary to the morphological analysis (TEIXEIRA *et al.*, 2004).

The pigmentation of the F1UFMS isolate was light beige in the PDA culture medium, while that of the isolate F2UFMS was violet (Table 3). The mycelial growth of both isolates was 12.85 cm day⁻¹, completing the Petri dish in seven days (Figure 3). Gupta, Misra and Gaur (2010) reported that *F. oxysporum* f. sp. *psidii* and *F. solani* showed maximum growth of 7.25 cm at 28 °C, after seven days of incubation. Skovgaard *et al.* (2003) recorded an average daily increase of 5.1 cm for the isolates of *F. commune*.

For variable sporulation in the leave-carnation-agar (LCA), it was observed that the isolate F1UFMS showed higher sporulation than F2UFMS what ranged from 11.25 × 10⁶ to 4.44 × 10⁶ conidia mL⁻¹, respectively (Table 3). Species of *F. verticillioides* produce sparse macroconidia

Figure 2 - Phylogenetic dendrogram based on neighbor-joining method from the DNA sequences of the ITS region (A) and 1 α - Elongation factor (B). The numbers on the branches indicate the percentage of repetitions of the bootstrap analysis in which the repeats were observed (1000 repetitions). * Isolates of *Fusarium* spp. obtained in this study



while *F. oxysporum* has abundant production of these structures (LESLIE; SUMERELL, 2006). Silva and Teixeira (2012) found an average sporulation of 3.09×10^3 conidia mL⁻¹ for *F. solani*, using 12-hour photoperiod and PDA medium. On the other hand, Lazarotto *et al.* (2014), in tests with to the culture complex *Gibberella fujikuroi*


on LCA culture medium, found high levels of sporulation, reaching 12.68×10^6 conidia mL⁻¹ for one of their isolates (F5). The LCA medium favors the ultimate expression of the characteristics and sporulation of the genus *Fusarium* (NELSON; TOUSSON; MARASAS, 1983; VENTURA, 1999).

Table 3 - Morphological characteristics of the *Fusarium* spp. isolates obtained of seeds from *Pinus* sp

Characteristics	F1UFMS <i>Fusarium verticillioides</i>	F2UFMS <i>Fusarium oxysporum</i>
Color of colony ¹	Tone light beige	Tone violet
Measure of conidia ²	(27.5) 36.9 (42.5) x (1.25) 2.7 (3.75) µm	(10) 14.67 (17.5) x (2.5) 2.7 (5) µm
Shape macroconidia ²	Long and slender; slightly falcate	Apical cell morphology short, slightly hook; basal cell foot shaped
Sporulation ²	11.25 x 10 ⁶ conidia mL ⁻¹	4.44 x 10 ⁶ conidia mL ⁻¹
Presence of sporodochia ²	+	+
Presence of microconidia	+	+
Presence of chlamydospores	-	+

¹Potato dextrose agar culture medium, 25 ± 2 °C and a photoperiod of 12 hours; ² Leave-carnation-agar culture medium, 25 ± 2 °C and a photoperiod of 12 hours

Figure 3 - Regression analysis (p<0,05) to the colony diameter of *F. verticillioides* and *F. oxysporum* (F1UFMS and F2UFMS) on PDA incubated at 25 °C and photoperiod 12 hours



We believe that the present study contributes to the knowledge on the diversity and pathogenicity of *F. oxysporum* and *F. verticillioides* associated with *Pinus* seeds. Further analyses involving other *Fusarium* species associated with *Pinus* seeds would be beneficial to compare the differences between the species and their effects on the *Pinus* seedlings.

CONCLUSION

The *Fusarium* species identified were *F. verticillioides* and *F. oxysporum* and are pathogenic to *P. taeda* and *P. elliottii*, causing damping-off in pre-and post-emergence.

ACKNOWLEDGMENTS

This work were supported by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) Foundation (Brazil) and Conselho Nacional

de Desenvolvimento Científico e Tecnológico (CNPq), project number 473722/2013-5.

REFERENCES

- ALFENAS, A. C.; MAFIA, R. G. **Métodos em fitopatologia**. Viçosa: UFV, 2007. 382 p. BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. **Regra para análise de sementes**. Brasília, DF, 2009. 395 p.
- ARIF, M.; CHAWLA, S.; ZAIDI, N. W. *et al.* Development of specific primers for genus *Fusarium* and *F. solani* using rDNA sub-unit and transcription elongation factor (TEF-1α) gene. **African Journal of Biotechnology**, v. 11, n. 2, p. 444-447, 2012.
- DELLAPORTA, S. L.; WOOD, J.; HICKS, J. B. A plant DNA miniprep: version II. **Plant Molecular Biology Reporter**, v. 1, n. 4, p. 19-21, 1983.
- FERREIRA, D. F. **Sisvar**: sistemas de análises de variância para dados balanceados: programa de análises estatísticas e planejamento de experimentos. versão 5.3 (Biud 75). Lavras: Universidade Federal de Lavras, 2010.
- GEISER, D. M. *et al.* FUSARIUM-ID v.1.0: a DNA sequence database for identifying *Fusarium*. **European Journal Plant Pathology**, v. 110, n. 5, p. 473-479, 2004.
- GERLACH, W.; NIRENBERG, H. **The genus Fusarium**: - a pictorial atlas. Berlin: Biologische Bundesanstalt für Land- und Forstwirtschaft, 1982. 406 p.
- GRIGOLETTI JUNIOR, A.; AUER, C. G. **Fusariose em mudas de Pinus taeda**. Colombo: Embrapa Florestas, 2006. 3 p. (Comunicado técnico 166).
- GUPTA, V. K.; MISRA, A. K.; GAUR, R. K. Growth characteristics of *Fusarium* spp. causing wilt disease in *Psidium guajava* l. in India. **Journal of Plant Protection Research**, v. 50, n. 4, p. 451-462, 2010.
- LANDERAS, E. *et al.* Outbreak of pitch canker caused by *Fusarium circinatum* on *Pinus* spp. in Northern Spain. **Plant Disease**, v. 89, n. 9, p. 1015, 2005.

- LAZAROTTO, M. *et al.* Mycelia growth and sporulation of *Fusarium chlamydosporum* species complex under different culture conditions. **Revista de Ciências Agrárias**, v. 57, n. 1, p. 35-40, 2014.
- LESLIE, J. F.; SUMMERELL, B. A. **The Fusarium laboratory manual**. 1. ed. USA: Blackwell Publishing, 2006. 388 p.
- LORI, G. A.; SALERNO, M. I. *Fusarium* species on seeds of *Pinus taeda* L. and *Pinus elliottii* Engelm. in Argentina. **Journal of Plant Diseases and Protection**, v. 110, n. 5, p. 437-443, 2003.
- MACIEL, C. G. *et al.* First report of *Fusarium sambucinum* associated on *Pinus elliottii* seeds in Brazil. **Plant Disease**, v. 97, n. 7, p. 995, 2013.
- MAGUIRE, J. D. Speed of germination aid in selection and evaluation for seeding emergence and vigor. **Crop Science**, v. 2, n. 2, p. 76-177, 1962.
- MARTÍN-PINTO, P.; PAJARES, J.; DÍEZ, J. Pathogenicity of *Fusarium verticillioides* and *Fusarium oxysporum* on *Pinus nigra* seedlings in northwest Spain. **Forest Pathology**, v. 38, n. 2, p. 78-82, 2008.
- MENEZES, J. P. *et al.* Variabilidade genética na região its do rDNA de isolados de *Trichoderma* spp. (Biocontrolador) e *Fusarium oxysporum* f. sp. *chrysanthemi*. **Ciência e Agrotecnologia**, v. 34, n. 1, p. 132-139, 2010.
- NELSON, P. E.; TOUSSON, T. A.; MARASAS, W. F. O. **Fusarium species: an illustrated manual for identification**. Philadelphia: Pennsylvania State University Press, 1983. 193 p.
- O'DONNELL, K. Molecular phylogeny of the *Nectria haematococca*-*Fusarium solani* species complex. **Mycologia**, v. 92, n. 5, p. 919-938, 2000.
- OCAMB, C. M.; JUZWIK, J.; MARTIN, F. B. *Fusarium* spp. and *Pinus strobus* seedlings: root disease pathogens and taxa associated with seed. **New Forests**, v. 24, n. 1, p. 67-79, 2002.
- PFENNING, L. H. *et al.* First report and characterization of *Fusarium circinatum*, the causal agent of pitch canker in Brazil. **Tropical Plant Pathology**, v. 39, n. 3, p. 210-216, 2014.
- REHNER, S. A.; BUCKLEY, E. A *Beauveria* phylogeny inferred from nuclear ITS and EF1- α sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. **Mycologia**, v. 97, n.1, p. 84-98, 2005.
- SCHMITZ, A.; RIESNER, D. Purification of nucleic acids by selective precipitation with polyethylene glycol 6000. **Analytical Biochemistry**, v. 354, n. 2, p. 311-313, 2006.
- SCHOCH, C. L. *et al.* Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. **Proceedings of the National Academy of Sciences of the United States of America**, v. 109, n. 16, p. 1-6, 2012.
- SILVA, J. L.; TEIXEIRA, R. N. V. Esporulação e crescimento micelial de *Fusarium solani* em diferentes meios de cultura e regimes de luminosidade. **Agro@mbiente On-line**, v. 6, n. 1, p. 47-52, 2012.
- SKOVGAARD, K. *et al.* *Fusarium commune* is a new species identified by morphological and molecular phylogenetic data. **Mycologia**, v. 95, n. 4, p. 630-636, 2003.
- STEENKAMP, E. T. *et al.* *Fusarium circinatum* and pitch canker of *Pinus* in Colombia. **Australasian Plant Pathology**, v. 41, n. 5, p. 483-491, 2012.
- TAMURA, K.; DUDLEY, J.; NEI, M. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. **Molecular Biology and Evolution**, v. 24, n. 8, p. 1596-1599, 2007.
- TEIXEIRA, H.; VIEIRA, M.; MACHADO, J. C. Marcadores RAPD na análise da diversidade genética de isolados de *Acremonium strictum*. **Fitopatologia Brasileira**, v. 29, n. 6, p. 651-655, 2004.
- TSUI, C. K. M.; WOODHALL, J.; CHEN, W. Molecular techniques for pathogen identification and fungus detection in the environment. **Ima Fungus**, v. 2, n. 2, p. 177-189, 2011.
- VENTURA, J. A. Taxonomia de *Fusarium* e seus segredos: parte I - história, meios e procedimentos de cultivo. **Revisão Anual de Patologia de Plantas**, v. 7, n. 1, p. 271-298, 1999.
- WHITE, T. J. *et al.* Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: INNIS, M. A. *et al.* (Ed.). **PCR Protocols: a guide to methods and applications**. New York: Academic Press, 1990. p. 315-322.