



Biota Neotropica

ISSN: 1676-0611

cjoly@unicamp.br

Instituto Virtual da Biodiversidade
Brasil

Oliveira Dias, Cristina de; Costa Bonecker, Sérgio Luiz
New records of Monstrilloida Sars, 1901 (Crustacea, Copepoda) on the Brazilian northeastern coast
Biota Neotropica, vol. 7, núm. 2, mayo-agosto, 2007, pp. 281-285
Instituto Virtual da Biodiversidade
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=199114293030>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

New records of Monstrilloida Sars, 1901 (Crustacea, Copepoda) on the Brazilian northeastern coast

Cristina de Oliveira Dias & Sérgio Luiz Costa Bonecker

Biota Neotropica v7 (n2)

<http://www.biotaneotropica.org.br/v7n2/pt/abstract?short-communication+bn00407022007>

Recebido em 24/10/06

Versão reformulada recebida em 18/02/07

Publicado em 01/05/07

Laboratório Integrado de Zooplâncton e Ictioplâncton, Departamento de Zoologia, Instituto de Biologia,
Universidade Federal do Rio de Janeiro – UFRJ,
CCS, Bloco A, Ilha do Fundão, CEP 21941-590, Rio de Janeiro, RJ, Brasil
Corresponding author: Cristina de Oliveira Dias, e-mail: crcldias@biologia.ufrj.br,
Sérgio Luiz Costa Bonecker, e-mail bonecker@biologia.ufrj.br

Abstract

Dias, C. O. & Bonecker S. L. C. **New records of Monstrilloida Sars, 1901 (Crustacea, Copepoda) On The Brazilian Northeastern Coast.** *Biota Neotrop.* May/Aug 2007 vol. 7, no. 2 <http://www.biotaneotropica.org.br/v7n2/pt/abstract?short-communication+bn00407022007>. ISSN 1676-0603.

During a series of zooplankton surveys carried out from 2001 through 2005 off the coast of the state of Bahia, Brazil, 98 individuals of monstrilloid copepods were collected. These belong to five species (*Monstrilla grandis*, *Cymbasoma cf. longispinosum*, *Cymbasoma cf. rigidum*, *Cymbasoma gracilis*, and *Cymbasoma quadridens*). The first three are recorded for the first time in the Bahia coastal region. The geographical range of *C. quadridens* is expanded to the Brazilian northeastern coast. The results presented herein increase to nine the number of nominal species of Monstrilloida known from off Bahia; the environmental diversity of Caravelas Channel with highly productive areas and coral reef zones harbor an abundant and diverse monstrilloid fauna that should be surveyed in more detail.

Keywords: new occurrence, zooplankton, Southwest Atlantic, Brazilian coast, Bahia, taxonomy.

Resumo

Dias, C. O. & Bonecker S. L. C. **Novas ocorrências para Monstrilloida Sars, 1901 (Crustacea, Copepoda) na costa Nordeste do Brasil.** *Biota Neotrop.* May/Aug 2007 vol. 7, no. 2 <http://www.biotaneotropica.org.br/v7n2/pt/abstract?short-communication+bn00407022007>. ISSN 1676-0603.

Durante uma série de amostragens de zooplâncton, realizadas de 2001 a 2005, na costa do estado de Bahia, Brasil, 98 exemplares de copépodes Monstrilloida foram coletados. Os exemplares pertencem a cinco espécies (*Monstrilla grandis*, *Cymbasoma cf. longispinosum*, *Cymbasoma cf. rigidum*, *Cymbasoma gracilis* and *Cymbasoma quadridens*). As primeiras três espécies foram coletadas pela primeira vez na região costeira do estado da Bahia. A distribuição geográfica de *C. quadridens* foi expandida para a costa nordeste brasileira. Os resultados apresentados aumentam a nove o número de espécies Monstrilloida conhecidas para a costa da Bahia. A diversidade ambiental do Canal de Caravelas, com áreas altamente produtivas e a proximidade de uma região de recifes de coral, abriga uma abundante e diversa fauna de Monstrilloida, que deve ser examinada mais detalhadamente.

Palavras-chave: nova ocorrência, zooplâncton, Atlântico sudoeste, costa brasileira, Bahia, taxonomia.

Introduction

Monstrilloid copepods are protelean parasites of benthic macroinvertebrates such as polychaetes and mollusks (Caullery & Mesnil 1914, Davis 1984); most postnaupliar and preadult stages are endoparasitic. Adults are the most conspicuous stage, because they are free-living (Suárez-Morales & Ivanenko 2004), and are captured by plankton nets in zooplankton surveys of coastal-neritic systems at all latitudes (Suárez-Morales & Dias 2001a).

The zooplankton communities of the continental shelf off the state of Bahia, on the northeast coast of Brazil, are still poorly known (Bonecker 1995, Neumann-Leitão 1994/1995). A previous survey of the monstrilloid copepod fauna off northeastern Brazil yielded records of *Monstrilla grandis* Giesbrecht, 1891, *Monstrilla rugosa* Davis, 1947, *Cymbasoma* cf. *rigidum* Thompson, 1888, *Cymbasoma* cf. *longispinosum* Bourne, 1890, and *Cymbasoma* cf. *gracilis* Gurney, 1927. Four new species were described: *Monstrilla careli* Suárez-Morales & Dias, 2000, *Monstrilla brasiliensis* Suárez-Morales & Dias, 2000, *Monstrilla satchmoi* Suárez-Morales & Dias, 2001b and *Monstrilla bahiana* Suárez-Morales & Dias, 2001b (Dias 2005). Of these species, *Monstrilla careli*, *M. brasiliensis*, *M. bahiana*, *M. satchmoi* and *Cymbasoma gracilis* were collected in the oceanic region off Bahia.

As part of serial surveys of the marine fauna off the coast of Bahia, zooplankton samples were collected from 2001 through 2005. Several specimens of monstrilloid copepods were collected and sorted for identification. The aim of this study was to expand the knowledge of the composition and distribution of the monstrilloid species in the region.

Material and Methods

The biological material examined was obtained as part of a project to study the copepod fauna off northeastern Brazil. The program was carried out from October 2001 to February 2005, at two sites along the coast of Bahia. The sites with the occurrence of monstrilloids were located between Boipeba (13° 35' 27" S and 38° 54' 56" W) and the Marau peninsula (14° 06' 32" S and 39° 01' 08" W), on the adjacent inner continental shelf near Camamu Bay (sta. 1-7) at stations as far as the 30-m isobath, and in the estuarine area of the Caravelas Channel (17° 44' 47" and 17° 48' 56" S and 39° 10' 51 and 39° 14' 11" W) and the adjacent coastal region (sta. 1-4) out to the 10-m isobath (Figure 1).

The samples were collected during the day, by subsurface horizontal and vertical hauls, with a conical net of mesh size 200 µm and mouth diameter 60 cm, fitted with a calibrated flowmeter. Samples were fixed and preserved in 4% buffered formalin. Monstrilloid copepods were sorted out from the original samples.

All the specimens found were deposited in the zooplankton collection of the Integrated Zooplankton and Ichthyoplankton Laboratory of the Federal University of Rio de Janeiro (MONSTRILLOIDA DZUFJR 175 to 191). The individuals were measured from the anterior end of the cephalic somite to the posterior margin of the anal somite.

Results and Discussion

A total of 98 individual monstrilloid copepods, belonging to the species *Monstrilla grandis*, *Cymbasoma gracilis*, *C. cf. longispinosum*, *C. rigidum* and *C. quadridens* Davis, were found in the surveyed area.

Two males of *C. quadridens* (1.19 and 1.26 mm, respectively) were collected. This species had a total length similar to that found by Dias (2005) in the Brazilian coast, and by Davis (1947) in the

northwestern Atlantic Ocean (Biscayne Bay, Florida). Hitherto, this species has been reported from the Atlantic Ocean (Davis 1947, Dias 1996, Johnsson 1998), including southeastern Brazil. This is the first record of the species off the Brazilian northeastern coast. The occurrence of *C. quadridens* in the area expands the distribution of this species northwards.

A single male of *Monstrilla grandis*, with a total body length of 1.06 mm was collected. The distribution of *M. grandis* is very wide (Suárez-Morales 2000a); this is the first record from the Bahia coastal region. The length of this specimen is similar to that found by Dias (2005) on the northeastern (state of Pernambuco), central and south coast of Brazil, and shorter than those found by Scott (1904) off Scotland, by Rose (1933) off France and by Ramírez (1971) off Argentina, which ranged from 1.6 to 2.0 mm. Suárez-Morales (2000a) reported total lengths of 0.61-0.65 mm (France). The size of copepods increases with decreasing temperature and with depth into oceanic waters. The warm-water communities contain smaller copepods than temperate waters, with cold polar waters containing large species (Hopcroft et al. 2001).

Four specimens of *Cymbasoma* cf. *longispinosum* (two males, one damaged; and two females) with total body lengths of 1.45 mm (male) and 1.96 and 2.06 mm (females), and two males of *Cymbasoma* cf. *rigidum*, each with total body length of 0.93 mm, were also recorded. The length of *Cymbasoma* cf. *longispinosum* is similar to that reported by Dias (2005) and Duarte (1999) for other Brazilian specimens. Rose (1933), from France, and Sars (1921), from Norway, reported larger lengths (F: 2.3 to 3.16 mm, M: 1.8 to 2.3 mm). The length of the males of *C. cf. rigidum* is similar to that found by Dias (2005) on the Brazilian coast, but differs from values reported by Wilson (1950) in the Philippines and Rose (1933) in France (1.5 to 1.8 mm).

In the present study, these nominal species are reported for the first time from the Bahia coastal region. *Cymbasoma* cf. *longispinosum* and *C. cf. rigidum* were found from off northeastern (state of Rio Grande do Norte) to southern Brazil (Dias 1996, 2005, Johnsson 1998, Duarte 1999, Dias 2005). Comparing the specimens found in different areas along the Brazilian coast, there are not morphological differences between the specimens found in the present study with the specimens designated as *Cymbasoma* cf. *longispinosum* and *C. cf. rigidum* by Dias (1996).

The distributions of some monstrilloid species, such as *C. longispinosum* and *C. rigidum*, have been reported to be worldwide (Isaac 1975); however, some authors consider that these two taxa are actually species-complexes, and that the strict forms have a more limited distributional range. The wide distribution reported for these species could certainly be a result of overlooking closely related species. Suárez-Morales (2000b) discussed the subtle differences shown by several species, which can be easily confused during casual observation; the author commented that this kind of problem is not uncommon within the group, because closer morphological analyses of monstrilloid copepods have shown that the wide distributional patterns reported for some of the commonest species may be misleading. *Cymbasoma morii* Sekiguchi 1982 (Grygier 1994), *C. chelemense* Suárez-Morales & Escamilla 2001, *C. californiensis* Suárez-Morales & Palomares 1999, and *C. rochai* Suárez-Morales & Dias 2000 are closely related to *Cymbasoma longispinosum*; and *Cymbasoma germanicum* (Timm 1893) is very closely related to *Cymbasoma rigidum* and to other, undescribed forms (Suárez-Morales 2006). Because of this problem, in the present report the specimens of *C. cf. longispinosum* and *C. cf. rigidum* are regarded as probable members of taxonomic complexes formed by the nominal species *C. longispinosum* and *C. rigidum*; they might be undescribed taxa. Although the size variation within a species may result from differences in the size of the host and the number of individuals within a




Figure 1. Map of the study area indicating the sampling stations with the occurrence of monstrilloids: a) inner continental shelf off Camamu; and b) estuarine area of Caravelas Channel and adjacent coastal region.

Figura 1. Mapa da área de estudo indicando as estações de amostragem com a ocorrência de Monstrilloida: a) plataforma continental interna de Camamu; e b) região estuarina do Canal de Caravelas e região costeira adjacente.

host (Suárez-Morales 2000a), the lengths of the specimens are provided in order to contribute to studies and revisions of the specimens belonging to these species-complexes.

Eighty-nine females (3 damaged) of *Cymbasoma gracilis* (1.28-1.86 mm) were collected. The total length range is similar to that found by Dias (2005). This species has been found in tropical waters of the Atlantic and Indian oceans and in the Mediterranean and Red seas (Gurney 1927, Isaac 1975). Dias (2005) gave the local distribution of this species as from the northeastern (states of Bahia and Rio Grande do Norte) to the southern Brazilian coast.

The occurrence of these species expanded the distribution proposed by Dias (2005) in the tropical species associations in Brazilian coast, characterized by the presence of Tropical Water in the area.

Comparing the two sampling areas, Camamu Bay and Caravelas Channel, most of the monstrolloids (95%) were found in the area of the Caravelas Channel and the adjacent coastal region (with highest values of salinity up to an average of 37), in shallow waters down to the 10 m isobath. The coastal zone of Caravelas, off southern Bahia, contains diverse and productive ecological systems, including the Abrolhos coral reefs, considered the most important coral-reef system of the South Atlantic Ocean. Suárez-Morales (2001) suggested that monstrolloids are particularly rich and abundant in coastal and reef areas because their hosts are aggregated and abundant in these environments. The coastal zone and the oceanic region adjacent to Caravelas is the most productive of the Bahia coast (Andrade & Domínguez 2002). This is an area of high diversity and productivity, related to the coral reefs, and the large number of monstrolloids can be attributed to a local mass liberation of adults from aggregated benthic hosts (polychaetes and mollusks). The biotic community of the coral reefs off southern Bahia is still poorly known. In the Abrolhos region, it is composed mainly of cnidarians. The other components of the benthic fauna are the sponges, mollusks, polychaete annelids and echinoderms (Castro 1994).

Because of their scarcity in the zooplankton samples, the value of the occurrence of each specimen is high. Previous reports have shown that this area harbors a rich variety of species, several of which are undescribed (Suárez-Morales & Dias 2001a). This report increases to nine the number of species of Monstrolloida known from the Bahia region and emphasizes the relevance of coastal and reef-related areas as harboring an abundant and diverse array of monstrolloid copepods.

Acknowledgments

This study was conducted in the Integrated Laboratory of Zooplankton and Ichthyoplankton, Department of Zoology, Institute of Biology, Federal University of Rio de Janeiro (UFRJ). It is part of a monitoring program of the Caravelas region developed by CEPEMAR Ltda., and of Camamu Bay developed by ENSR International do Brasil Ltda. and KRITERIA Consultores Associados Ltda. We thank the ARACRUZ Celulose S.A., El Paso Óleo and Gás do Brasil Ltda. for placing some of their zooplankton material at our disposal. We thank Janet W. Reid for improving the English text.

References

ANDRADE, A.C. DA S. & DOMINGUEZ, J.M.L. 2002. Geologic-Geomorphological Informations As Subsidies For Environmental Analysis: The Example of Caravelas Coastal Plain, Bahia. *Bol. Par. Geoc.* 51:9-17.

BONECKER, S.L.C. 1995. Dinâmica do zooplâncton no sistema estuarino do rio Mucuri-BA. Tese de Doutorado, Ciências Biológicas (Zoologia), Universidade de São Paulo, São Paulo, 194pp.

BOURNE, G.C. 1890. Notes on the genus *Monstrilla* Dana. *Quekett J. Microsc. Sci.* 30(4):565-578, pl. 37.

CASTRO, C.B. 1994. Corals of Southern Bahia. In: *Corals of Southern Bahia* (B. Hetzel & C.B. Castro, eds.). Editora Nova Fronteira, Rio de Janeiro, p.161-176.

CAULLERY, M. & MESNIL, F. 1914. Sur deux Monstrolloides parasites d'Annelides (*Polydora giardi* Mesn. et *Syllis gracilis* Gr.). *Bull. Sci. Fran. Bel.* (7)48(1):15-29.

DAVIS, C.C. 1947. Two monstrolloids from Biscayne Bay, Florida. *Trans. Am. Microsc. Soc.* 66(4):390-395.

DAVIS, C.C. 1984. Planktonic Copepoda (including Monstrolloida). In: *Marine plankton life cycle strategies* (K.A. Steidinger & L.M. Walter, eds.). CRC Press, Boca Raton, Florida, p.67-91.

DIAS, C.O. 1996. Monstrolloida (Copepoda) off the Brazilian coast. *Hydrobiologia* 324(3):253-256.

DIAS, C.O. 2005. Taxonomy and geographic distribution of the Monstrolloida (Copepoda Crustacea) from the South Occidental Atlantic. *Biota Neotrop.* 5(2):<http://www.biotaneotropica.org.br/v5n2/pt/fullpaper?bn02005022005+en> (last accessed on 07/06/07).

DUARTE, A.K. 1999. Ocorrência de Monstrolloida (Copepoda) em águas costeiras do Sul do Brasil. *Nauplius* 7:201-202.

GIESBRECHT, W. 1891. Elenco dei Copepodi pelagici raccolti dal Tenente di vascello Gaetano Chierchia durante il viaggio della R. Corvetta 'Vettor Pisani' negli anni 1882-1885 e dal Tenente di vascello Francesco Orsini nel Mar Rosso, nel 1884. *Rendiconti delle Sedute della Real Accademia dei Lincei* 7(1):474-481.

GRYGIER, M.J. 1994. Nomenclature, redescription, and new record from Okinawa of *Cymbasoma morii* Sekiguchi, 1982 (Monstrolloida). *Hydrobiologia* 292/293: 23-29.

GURNEY, R. 1927. VIII. Report on the Crustacea Copepoda and Cladocera of the plankton. *Trans. Zool. Soc. Lond.* 22(2):139-171.

HOPCROFT, R. R., ROFF, J. C. & CHAVEZ, F. P. 2000. Size paradigms in copepod communities: a re-examination. *Hydrobiologia* 453-454: 133-141.

ISAAC, M.J. 1975. Copepoda. Sub-order Monstrolloida. Fiches d'Identification du Zooplankton, Conseil International pour l'Exploration de la Mer 144/145:1-10, figs.

JOHNSON, R. 1998. Maxillopoda - Copepoda, Monstrolloida. In: *Catalogue of Crustacea of Brazil* (P.S. Young, ed.). Museu Nacional, Série Livros, Rio de Janeiro, 6:261-262.

NEUMANN-LEITÃO, S. 1994/1995. Resenha literária sobre zooplâncton estuarino no Brasil. *Trab. Oceanogr. Univ. Fed.* 23:25-53.

RAMÍREZ, F.C. 1971. Nuevas localidades para *Monstrilla grandis* Giesbrecht, 1892 y *Monstrilla helgolandica* Claus, 1863 (Copepoda: Monstrolloida) hallados en aguas de la plataforma Argentina. *Physis* 30(81):377-383.

ROSE, M. 1933. Copépodes pelagiques. *Faune de France*, Paris, 26:1-374, figs. 1-456.

SARS, G.O. 1901. An account of the Crustacea of Norway with short descriptions and figures of all the species. Vol. IV. Copepoda Calanoida. The Bergen Museum, Bergen, p.1-28, figs. 1-16.

SARS, G.O. 1921. An account of the Crustacea of Norway with short descriptions and figures of all the species. Vol. VII. Copepoda Monstrolloida & Notodelphyoida. The Bergen Museum, Bergen, 91pp., figs. 1-37.

SCOTT, T. 1904. Notes of some rare and interesting marine Crustacea. Twenty-second Annual Report of the Fishery Board for Scotland, Being for the Year 1903. Part III. Scientific Investigations, 22(3):242-261, Pls. 13-15.

SEKIGUCHI, H. 1982. Monstrolloid copepods from Ago Bay, central Japan. *Proc. Jpn Soc. Syst. Zool.* 22: 24-34, figs. 1-6 (Japanese summary).

SUÁREZ-MORALES, E. 2000a. Taxonomic report on some monstrolloids (Copepoda, Monstrolloida) from Toulon Bay, France. *Bull. Inst. R. Sci. Nat. Belg.* 70:107-118.

SUÁREZ-MORALES, E. 2000b. A new species and new geographic records of *Monstrilla* (Copepoda: Monstrolloida) from the Philippines. *J. Crust. Biol.* 20(4):680-686.

SUÁREZ-MORALES, E. 2001. Taxonomic report on a collection of monstrolloids (Copepoda: Monstrolloida) from Banco Chinchorro, Mexico

- with description of a new species. An. Inst. Biol. Univ. Nac. Autón. Mex. Zool. 72(1):9-28.
- SUÁREZ-MORALES, E. 2006. Validation and redescription of *Cymbasoma germanicum* (Timm) (Crustacea: Copepoda: Monstrilloidea) from Helgoland with comments on *Cymbasoma rigidum* Thompson. Helgol. Mar. Res. 60(3):171-179.
- SUÁREZ-MORALES, E. & DIAS, C.O. 2000. Two new species of Monstrilloidea (Copepoda) from Brazil. J. Mar. Biol. Ass. U.K. 80(3670):1-9.
- SUÁREZ-MORALES, E. & DIAS, C.O. 2001a. A new species of *Monstrilla* (Copepoda: Monstrilloidea) from Brazil with notes on *M. brevicornis* Isaac. Proc. Biol. Soc. Wash. 114(1):219-228.
- SUÁREZ-MORALES, E. & DIAS, C.O. 2001b. Taxonomic report of some monstrilloids (Copepoda: Monstrilloidea) from Brazil with description of four new species. Bull. Inst. R. Sci. Nat. Belg. 71:65-81.
- SUÁREZ-MORALES, E. & ESCAMILLA, J.B. 2001. Taxonomic report on some monstrilloids (Copepoda, Monstrilloidea) from southeast Mexico with the description of a new species of *Monstrilla*. J. Nat. Hist. 35(10): 1433-1445.
- SUÁREZ-MORALES, E. & IVANENKO, V.N. 2004. Two new species of *Monstrillopsis* Sars (Crustacea: Copepoda: Monstrilloidea) from the White Sea and Norway, with comments on *M. dubia* Scott. Arctic 57(1): 37-46.
- SUÁREZ-MORALES, E., PALOMARES-GARCIA, R. 1999. *Cymbasoma californiense*, a new Monstrilloid (Crustacea: Copepoda: Monstrilloidea) from Baja California, Mexico. Proc. Biol. Soc. Wash. 112(1):189-198.
- THOMPSON, I.C. 1888. Copepoda of Madeira and the Canary Islands, with descriptions of new genera and species. J. Linn. Soc. London (Zool.) 20:145-156, pls.10-13.
- TIMM, R. 1893. *Monstrilla grandis* Giesbr. *M. helgolandica* Claus, *Thaumaleus germanicus* n. sp. Zool. Anz. 16(432):418-420.
- WILSON, C.B. 1950. Contributions to the biology of the Philippine Archipelago and adjacent regions. Copepods gathered by the United States Fisheries Steamer 'Albatross' from 1887 to 1909, chiefly in the Pacific Ocean. Bull. U.S. Nat. Mus. 100(14,4):141-441, pls. 2-36.