



Biota Neotropica

ISSN: 1676-0611

cjoly@unicamp.br

Instituto Virtual da Biodiversidade
Brasil

Leme da Cunha, Nicolay; Fischer, Erich; Ferreira Santos, Carolina
Bat assemblage in savanna remnants of Sonora, central-western Brazil
Biota Neotropica, vol. 11, núm. 3, julio-septiembre, 2011, pp. 197-201
Instituto Virtual da Biodiversidade
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=199121042016>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Bat assemblage in savanna remnants of Sonora, central-western Brazil

Nicolay Leme da Cunha^{1,3}, Erich Fischer² & Carolina Ferreira Santos¹

¹Programa de Pós-graduação em Ecologia e Conservação,
Universidade Federal de Mato Grosso do Sul - UFMS, CEP 79070-900, Campo Grande, MS, Brazil

²Departamento de Biologia, Universidade Federal de Mato Grosso do Sul,
UFMS, CEP 79070-900, Campo Grande, MS, Brazil

³Corresponding author: Nicolay Leme da Cunha, e-mail: nicolaycunha@gmail.com

CUNHA, N.L., FISCHER, E. & SANTOS, C.F. **Bat assemblage in savanna remnants of Sonora, central-western Brazil.** Biota Neotrop. 11(3): <http://www.biotaneotropica.org.br/v11n3/en/abstract?inventory+bn03311032011>

Abstract: The Cerrado (savanna) is a wide Neotropical formation, but the knowledge on the occurrence and distribution of bat species for this phytogeographic region is scarce, especially in the western portion. Here we address what are the bat species and their relative abundances in a western Cerrado site, municipality of Sonora, Mato Grosso do Sul, Brazil. Bats were mist-netted in eight non consecutive months over the course of two years. Sampled bats (n = 507) belonged to 18 species. The estimator Jackknife 1 estimated 22 species, and diversity (H') was 1.6. Richness and diversity of bats in Sonora were slightly higher than in the southern Cerrado, and similar to or lower than in the central Cerrado. Frugivorous species predominated, as expected, however richness and abundance of nectarivorous were unexpectedly high. *Carollia perspicillata* was the dominant species. The endemic bat *Lonchophylla dekeyseri* was highly abundant in Sonora, which enhances the importance of conserving the savanna remnants in this region.

Keywords: Cerrado, Chiroptera, diversity of species, *Lonchophylla dekeyseri*, Mato Grosso do Sul, *Tonatia bidens*.

CUNHA, N.L., FISCHER, E. & SANTOS, C.F. **Comunidade de morcegos em remanescentes de savana em Sonora, centro-oeste do Brasil.** Biota Neotrop. 11(3): <http://www.biotaneotropica.org.br/v11n3/pt/abstract?inventory+bn03311032011>

Resumo: O Cerrado (savana) é uma ampla formação neotropical, porém o conhecimento sobre a ocorrência e distribuição de espécies de morcegos é escasso para esta região fitogeográfica, principalmente na porção oeste. Estudamos aqui quais são as espécies de morcegos e suas abundâncias relativas em uma localidade da porção oeste do Cerrado, no município de Sonora, Mato Grosso do Sul, Brasil. Morcegos foram capturados em redes-de-neblina em oito meses não consecutivos ao longo de dois anos. Os morcegos amostrados (n = 507) pertenceram a 18 espécies. O estimador Jackknife 1 estimou 22 espécies, e a diversidade (H') foi 1,6. A riqueza e a diversidade de morcegos em Sonora foram pouco maiores que na porção sul do Cerrado, e semelhantes ou menores que no Cerrado central. Espécies frugívoras predominaram, como esperado, porém a riqueza e abundância de nectarívoras foram inesperadamente altas. *Carollia perspicillata* foi a espécie dominante. O morcego endêmico *Lonchophylla dekeyseri* foi muito abundante em Sonora, fato que aumenta a importância de conservar os remanescentes de savana dessa região.

Palavras-chave: Cerrado, Chiroptera, diversidade de espécies, *Lonchophylla dekeyseri*, Mato Grosso do Sul, *Tonatia bidens*.

Introduction

Brazil is the second richest country in bat species, roosting approximately 15% of the known species of bats in the world (Bernard et al. 2010). The Brazilian savanna (Cerrado) is among the widest phytogeographic formations in the Neotropics, covering approximately 2 million km² (Scariot et al. 2005). Nonetheless, the knowledge about bat species occurrence and distribution is still incipient in the Cerrado (Gregorin & Ditchfield 2005, Zortéa & Alho 2008, Santos et al. 2010), especially in the western portion which includes the states of Mato Grosso do Sul and Mato Grosso (Bernard et al. 2010). In spite of bats have been increasingly surveyed in Mato Grosso do Sul (e.g. Camargo & Fischer 2004, Bordignon 2006, Longo et al. 2007, Camargo et al. 2009, Cunha et al. 2009, Ferreira et al. 2010, Alho et al. 2011), there is a large gap of data in the northern Cerrado of this state (Bernard et al. 2010). This region urges for biological inventories and conservation initiatives, because natural areas have been rapidly transformed into cattle ranches, monocultures, hydroelectric dams, and sucroalcohol facilities (Bernard et al. 2010); (pers. obs.). Here, we asked what are the bat species and their relative abundances in savanna remnants in the municipality of Sonora, a northern region of Cerrado in Mato do Grosso do Sul.

Methods

The study was carried out in savanna remnants (17° 32' 19.80" S and 54° 26' 2.91" W) inserted in a matrix of pastures and plantations of soybean, sorghum and sugar cane, close to the Correntes river, municipality of Sonora, Mato Grosso do Sul, Brazil. Segments of rock cliffs from karstic formation occur along the riverbanks. The climate is Aw of Köppen, the mean annual temperature and rainfall are approximately 22 °C and 1500 mm, respectively (Instituto... 1992). The savanna remnants consist of "cerradão" (dense forests with shaded understory), "cerrado *stricto sensu*" (marked presence of shrubs and trees interspersed with open fields), "vereda" (marshlands with *Mauritia flexuosa* palm trees), and riparian forests (Instituto... 1992).

Bats were mist-netted in May (three nights), June (four nights), September (four nights), and November (four nights) of 2008, and February (two nights), June (four nights), August (four nights), and November (four nights) of 2010. In each sampling night, four to six mist-nets of 12 × 3 m and two of 9 × 3 m were set at ground level, opened at dusk and closed after 4-9 hours, depending on rainfall. Total sampling effort was 42,282 h.m² (cf. Straube & Bianconi 2002); and month efforts were 7,182 h.m² in May, 7,074 h.m² in June, 7,371 h.m² in September and 7,056 h.m² in November of 2008; 2,160 h.m² in February, 4,788 h.m² in May, 5,796 h.m² in August, and 3,420 h.m² in November of 2010.

Identifications of bat species were based on literature (Vizotto & Taddei 1973, Taddei et al. 1983, Taddei 1996, Gregorin & Taddei 2002); nomenclature follows Simmons (2005), except *Artibeus planirostris* (Larsen et al. 2007). To help identifications, bats' forearm length and mass were measured with caliper (±1 mm) and dynamometer (±1 g), respectively. They were marked with nontoxic ink for identifying recaptures in a same night, and then released in the site of capture. Twenty two specimens were collected and deposited as voucher in the zoological collection of the Universidade Federal de Mato Grosso do Sul (ZUFMS-M): *Anoura caudifer* (E. Geoffroy 1818) – 292 e 293, *Anoura geoffroyi* Gray 1838 – 246, *Artibeus cinereus* (Gervais 1856) – 389, *Carollia perspicillata* (Linnaeus 1758) – 242; *Glossophaga soricina* (Pallas 1766) – 247; *Lonchophylla dekeyseri* Taddei, Vizotto & Sazima 1983 – 280, 522 e 523; *Molossops temminckii* (Burmeister 1854) – 294 e 390; *Natalus stramineus* Gray 1838 – 295; *Phyllostomus discolor* (Wagner

1843) – 235, 236 e 467, *Platyrrhinus helleri* (Peters 1866) – 241, *P. lineatus* (E. Geoffroy 1810) – 239 e 240, *Pteronotus parnellii* (Gray 1843) – 233, *Tonatia bidens* (Spix 1823) – 237, *Trachops cirrhosus* (Spix, 1823) – 524.

We evaluated sufficiency of sampling using rarefaction curves based on the cumulative quantity of captured individuals, and estimated total expected richness with Jackknife 1 estimator (1000 randomizations) based on sampling nights. Shannon's diversity index (H'), using neperian logarithm and the 95% CI (bootstrap, 1000 randomizations), were calculated for the bat assemblage. Analyses were done in the Vegan package (Oksanen et al. 2010), R program (R Development Core Team 2010).

Results

We captured 507 individuals belonging to four families, 14 genera and 18 species (Table 1). The rarefaction curve (Figure 1) showed that capture effort was sufficient to include most of the species at the study site. The Jackknife 1 estimated 22 ± 1.9 (mean \pm SD) species of bats, thus our observed richness ($n = 18$) included approximately 82% of the bat species which could be captured in mist-nets. The diversity (H') was 1.6, with 95% CI between 1.4 and 1.7. Phyllostomidae was the richest and commonest family (15 species, 97% of the captures); other families presented one species only. Frugivorous species were more common ($n = 6$) than nectarivorous ($n = 4$), insectivorous ($n = 4$), carnivorous ($n = 2$), omnivorous ($n = 1$), and hematophagous ($n = 1$) (Table 1). *Carollia perspicillata* was the dominant species; its relative abundance was one order of magnitude higher than that of the other common bat species – *L. dekeyseri* (Figure 2a), *G. soricina*, *P. lineatus*, *D. rotundus*, and *A. planirostris*. Thirteen additional species were relatively rare as they accounted for less than 10 individuals captured; only one individual of *A. cinereus* (Figure 2b), *C. auritus*, and *N. stramineus* was captured.

Discussion

The richness of bat species ($n = 18$) in savanna remnants in Sonora is slightly higher than that found in sites of southern Cerrado (10 and 13 species; respectively Cunha et al. 2009, Camargo et al. 2009), but it is similar to or less than richness (16-28 species) found in





Figure 1. Rarefaction curve of bat species on cumulative number of captured individuals in Sonora, Mato Grosso do Sul (dotted line = 95% confidence interval).

Figura 1. Curva de rarefação de espécies de morcegos em relação ao número cumulativo de indivíduos capturados em Sonora, Mato Grosso do Sul (linha pontilhada = intervalo de confiança de 95%).

Table 1. Number of captured individuals of 18 species of bats in Sonora, Mato Grosso do Sul.**Tabela 1.** Número de indivíduos capturados de 18 espécies de morcegos em Sonora, Mato Grosso do Sul.

Families/Species	Guild ^a	Females						Males				Σ
		Y	A	P	L	PoL	Σ	Y	AT	ET	Σ	
Phyllostomidae												
<i>Carollia perspicillata</i>	Frug	12	82	31	48	-	173	5	38	90	133	306
<i>Lonchophylla dekeyseri</i>	Nect	4	11	3	3	1	22	1	2	16	19	41
<i>Glossophaga soricina</i>	Nect	3	8	1	4	1	17	-	2	13	15	32
<i>Platyrrhinus lineatus</i>	Frug	1	12	2	3	-	18	1	10	3	14	32
<i>Desmodus rotundus</i>	Hema	1	8	10	2	-	21	1	3	6	10	31
<i>Artibeus planirostris</i>	Frug	-	1	8	4	-	13	-	3	4	7	20
<i>Tonatia bidens</i>	Insec	-	1	-	2	-	3	-	2	2	4	7
<i>Phyllostomus discolor</i>	Omni	-	1	-	1 ^b	-	2	-	1	2	3	5
<i>Anoura caudifer</i>	Nect	1	-	-	-	-	1	-	-	3	3	4
<i>Artibeus lituratus</i>	Frug	-	-	1	-	-	1	-	-	3	3	4
<i>Platyrrhinus helleri</i>	Frug	-	-	-	-	-	-	-	3	1	4	4
<i>Anoura geoffroyi</i>	Nect	-	-	-	-	-	-	-	1	1	2	2
<i>Trachops cirrhosus</i>	Carn	-	-	-	1	-	1	-	1	-	1	2
<i>Artibeus cinereus</i>	Frug	-	-	-	1	-	1	-	-	-	-	1
<i>Chrotopterus auritus</i>	Carn	-	-	-	-	-	-	-	-	1	1	1
Molossidae		-	-	-	-	-	-	-	-	-	-	-
<i>Molossops temminckii</i>	Insec	-	-	3	1	-	4	-	1	3	4	8
Mormoopidae		-	-	-	-	-	-	-	-	-	-	-
<i>Pteronotus parnellii</i>	Insec	-	1	1	-	-	2	-	3	1	4	6
Natalidae		-	-	-	-	-	-	-	-	-	-	-
<i>Natalus stramineus</i>	Insec	-	-	-	1	-	1	-	-	-	-	1

^aFrug: Frugivorous, Nect: Nectarivorous, Hema: Hematophagous, Insec: Insectivorous, Omni: Omnivorous, Carn: Carnivorous. ^bIt was additionally pregnant.

**Figure 2.** a) *Lonchophylla dekeyseri* and b) *Artibeus cinereus* recorded in Sonora, Mato Grosso do Sul (photos: N. L. Cunha).**Figure 2.** a) *Lonchophylla dekeyseri* e b) *Artibeus cinereus* registrados em Sonora, Mato Grosso do Sul (fotos: N. L. Cunha).

central region of Cerrado (Marinho-Filho et al. 1998, Rodrigues et al. 2002, Bordignon 2006, Zortéa & Alho 2008, Zortéa et al. 2010). Likewise, diversity of bat species in Sonora ($H' = 1.6$) is comparable to that in southern Cerrado sites ($H' = 1.5$ -1.8) (Camargo et al. 2009, Cunha et al. 2009, Ferreira et al. 2010) but lower than in central Cerrado ($H' = 2.2$) (Zortéa & Alho 2008). Therefore, available data roughly indicate that richness and diversity of bats increase toward northern (or central region; based on Zortéa & Alho 2008) of Cerrado, a possibility that deserves studies. Overall, the increasing of diversity with decreasing of latitude is expected (Stevens 2004, Willig & Bloch 2006).

The predominance of fruit-eating bats at the study site is comparable to other regions of Cerrado, however, the richness and relative abundance of nectar-feeding bats is apparently high in Sonora (Gonçalves & Gregorin 2004, Bordignon 2006, Camargo et al. 2009, Cunha et al. 2009, Zortéa & Alho 2008). This result can partially be due to the high availability of chiropterophilous flowers (*Bauhinia unguilata*, *Calliandra* sp., *Caryocar brasiliense*, *Hymenaea stigonocarpa*, and *Lafoesia* sp.) and of roosts in rock cavities at the Correntes riverbanks (per. obs.). The four nectarivore species registered in Sonora are common users of caves as day roosts, which seems to be especially important for *L. dekeyseri* and *A. geoffroyi* (Baumgarten & Vieira 1994, Coelho & Marinho-Filho 2002, Esbérard et al. 2005). Insectivorous bats in Sonora are represented by few species when compared to sites in southern Cerrado or in savannas in Bolivia and Paraguay, where insectivorous bats predominate (Cunha et al. 2009, Aguirre 2002, Willig et al. 2000, Alho et al. 2011, Zortéa & Alho 2008). Only one Phyllostominae insectivorous species occurred in Sonora, *T. bidens*. However, additional insectivorous species of this subfamily would be expected, based on the occurrences reported for southern Cerrado and Pantanal (Cunha et al. 2009, Alho et al. 2011). On the other hand, the proportion of captured individuals of *T. bidens* at the study site ($n = 7/507$) was markedly higher than that recorded in sites in southern Cerrado and Pantanal ($n = 5/9037$) (Alho et al. 2011).

Carollia perspicillata has commonly been recorded as the dominant species in different vegetation formations, regenerating areas, rocky outcrops and caves (Bredt et al. 1999, Esbérard et al. 2005, Bordignon 2006, Bernard & Fenton 2007, Castro-Arellano et al. 2007). However, *C. perspicillata* is generally less abundant than *A. planirostris*, *P. lineatus* and *S. lillium* in the southern Cerrado (Camargo et al. 2009, Cunha et al. 2009, Alho et al. 2011). The relative abundance of this species in local assemblages, therefore, appears to increase towards the north or the center of Cerrado (Zortéa et al. 2010, Zortéa & Alho 2008, Bernard & Fenton 2007, this study). The opposite occurs for *A. planirostris*, a relatively less common species in Sonora and other sites in central and northern Cerrado than in the southern Cerrado and Pantanal (Pulchério-Leite et al. 1999, Bordignon 2006, Camargo et al. 2009, Cunha et al. 2009, Teixeira et al. 2009, Zortéa et al. 2010, Alho et al. 2011).

Lonchophylla dekeyseri is unusually abundant at the study site. It is endemic of Cerrado and globally classified as "near threatened" (International... 2010), which highlights the importance of conserving the savanna remnants in Sonora. Only six populations of this species are known, all associated with karstic regions (Aguirre et al. 2010). The high abundance of *L. dekeyseri* indicates that Sonora is close to the core region of population "B" (cf. Aguirre et al. 2010) and expands the distribution of this population to the west. *Lonchophylla dekeyseri* and *A. cinereus* were only found in northern Cerrado of Mato Grosso do Sul (Coelho 2005, this study), and were not included in bat species compilation for the state (Cáceres et al. 2008). *Trachops cirrhosus* was also recently recorded for the first time in Mato Grosso do Sul (Alho et al. 2011); our record in Sonora is the second for this state.

Considering *L. dekeyseri*, *A. cinereus*, *T. cirrhosus* and two other recently reported species (Santos et al. 2010, Silveira et al. 2011), the updated known fauna of bats in Mato Grosso do Sul reaches 70 species.

Acknowledgements

We thank to the company ANAMBI - Análise Ambiental for logistic support; to J. L. Holladay for English revision; to CAPES, FUNDECT and CNPq for grants to N. L. Cunha, C. F. Santos and E. Fischer, respectively. Bat captures were carried out under IBAMA license (no. 10615-1).

References

- AGUIAR, L.M.S., BRITO, D. & MACHADO, R.B. 2010. Do current vampire bat (*Desmodus rotundus*) population control practices pose a threat to dekeyser's nectar bat's (*Lonchophylla dekeyseri*) long-term persistence in the Cerrado? Acta Chiropterol. 12:275-282. <http://dx.doi.org/10.3161/150811010X537855>
- AGUIRRE, L.F. 2002. Structure of a Neotropical savanna bat community 2002. J. Mammal. 83:775-784. [http://dx.doi.org/10.1644/1545-1542\(2002\)083%3C0775:SOANSB%3E2.0.CO;2](http://dx.doi.org/10.1644/1545-1542(2002)083%3C0775:SOANSB%3E2.0.CO;2)
- ALHO, C.J.R., FISCHER, E., OLIVEIRA-PISSINI, L.F. & SANTOS, C.F. 2011. Bat-species richness in the Pantanal floodplain and its surrounding uplands. Braz. J. Biol. 71:311-320.
- BAUMGARTEN, J.E. & VIEIRA, E.M. 1994. Reproductive seasonality and development of *Anoura geoffroyi* (Chiroptera: Phyllostomidae) in central Brazil. Mammalia 58:415-422. <http://dx.doi.org/10.1515/mamm.1994.58.3.415>
- BERNARD, E. & FENTON, M.B. 2007. Bats in a fragmented landscape: Species composition, diversity and habitat interactions in savannas of Santarém, Central Amazonia, Brazil. Biol. Conserv. 134:332-343. <http://dx.doi.org/10.1016/j.biocon.2006.07.021>
- BERNARD, E., AGUIAR, L.M.S. & MACHADO, R.B. 2010. Discovering the Brazilian bat fauna: a task for two centuries? Mammal Rev. 41:23-39. <http://dx.doi.org/10.1111/j.1365-2907.2010.00164.x>
- BORDIGNON, M.O. 2006. Diversidade de morcegos (Mammalia, Chiroptera) do Complexo Aporé-Sucuriú, Mato Grosso do Sul, Brasil. Rev. Bras. Zool. 24:1002-1009. <http://dx.doi.org/10.1590/S0101-81752006000400004>
- BREDT, A., UIEDA, W. & MAGALHÃES, E.D. 1999. Morcegos cavernícolas da região do Distrito Federal, centro-oeste do Brasil (Mammalia, Chiroptera). Ver. Bras. Zool. 16:731-770. <http://dx.doi.org/10.1590/S0101-81751999000300012>
- CÁCERES, N.C., CARMIGNOTTO, A.P., FISCHER, E. & SANTOS, C.F. 2008. Mammals from Mato Grosso do Sul, Brazil. Check List 4:321-335.
- CAMARGO, G. & FISCHER, E. 2004. Primeiro registro do morcego *Mimon crenulatum* (Phyllostomidae) no Pantanal, sudoeste do Brasil. Biota Neotrop. 5:181-184. <http://www.biotaneotropica.org.br/v5n1/pt/abstract?short-communication+BN00705012005> (último acesso em 16/02/2011).
- CAMARGO, G., FISCHER, E., GONÇALVES, F., FERNANDES, G. & FERREIRA, S. 2009. Morcegos do Parque Nacional da Serra da Bodoquena, Mato Grosso do Sul, Brasil. Chiropt. Neotrop. 15:417-424.
- CASTRO-ARELLANO, I., PRESLEY, S.J., SALDANHA, L.N., WILLIG, M.R. & WUNDERLE, J.M. 2007. Effects of reduced impact logging on bat biodiversity in terra firme forest of lowland Amazonia. Biol. Cons. 138:269-285. <http://dx.doi.org/10.1016/j.biocon.2007.04.025>
- COELHO, D.C. 2005. Ecologia e conservação da quiroptero-fauna no corredor Cerrado-Pantanal. Tese de Doutorado, Universidade de Brasília, Brasília.
- COELHO, D.C. & MARINHO-FILHO, J. 2002. Diet and activity of *Lonchophylla dekeyseri* (Chiroptera, Phyllostomidae) in the Federal District, Brazil. Mammalia 66:319-330. <http://dx.doi.org/10.1515/mamm.2002.66.3.319>

Bats of savanna remnants

- CUNHA, N.L., FISCHER, E., CARVALHO, L.F.A.C. & SANTOS, C.F. 2009. Bats of Buraco das Araras reserve, southwestern Brazil. *Biota Neotrop.* 9:189-195. <http://www.biotaneotropica.org.br/v9n4/pt/abstract?inventory+bn02909042009> (último acesso em 16/02/2011).
- ESBÉRARD, C.L., MOTTA, J.A. & PERIGO, C. 2005. Morcegos cavernícolas da Área de Proteção Ambiental (APA) Nascentes do Rio Vermelho, Goiás. *Rev. Bras. Zool.* 7:311-325.
- FERREIRA, C.M.M., FISCHER, E. & PULCHÉRIO-LEITE, A. 2010. Fauna de morcegos em remanescentes urbanos de Cerrado em Campo Grande, Mato Grosso do Sul. *Biota Neotrop.* 10:155-160. <http://www.biotaneotropica.org.br/v10n3/pt/abstract?article+bn02910032010> (último acesso em 16/02/2011).
- GONÇALVES, E. & GREGORIN, R. 2004. Quirópteros da Estação Ecológica da Serra das Araras, Mato Grosso, Brasil, com o primeiro registro de *Artibeus gnomus* e *A. anderseni* para o cerrado. *Lundiana* 5:143-149.
- GREGORIN, R. & DITCHFIELD, A.D. 2005. A new genus and species of nectar-feeding bat in the tribe Lonchophyllini (Phyllostomidae: Glossophaginae) from northeastern Brazil. *J. Mammal.* 86:403-414. <http://dx.doi.org/10.1644/BRB-229.1>
- GREGORIN, R. & TADDEI, V.A. 2002. Chave artificial para a identificação de molossídeos brasileiros (Mammalia, Chiroptera). *Mastoz. Neotrop.* 9:13-32.
- INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA - IBGE. 1992. Manual técnico da vegetação brasileira. IBGE, Rio de Janeiro, p.1-92.
- INTERNATIONAL UNION FOR CONSERVATION OF NATURE - IUCN. 2010. IUCN Red List of Threatened Species. Version 2010.4. www.iucnredlist.org (último acesso em 05/06/2011).
- LARSEN, P.A., HOOFFER, S.R., BOZEMAN, M.C., PEDERSEN, S.C., GENOWAYS, H.H., PHILLIPS, C.J., PUMO, D.E. & BAKER, R.J. 2007. Phylogenetics and Phylogeography of the *Artibeus jamaicensis* Complex Based on Cytochrome- b DNA Sequences. *J. Mammal.* 88:712-727. <http://dx.doi.org/10.1644/06-MAMM-A-125R.1>
- LONGO, J.M., FISCHER, E., CAMARGO, G. & SANTOS, C.F. 2007. Ocorrência de *Vampyressa pusilla* (Chiroptera, Phyllostomidae) no Pantanal sul. *Biota Neotrop.* 7:369-372. <http://www.biotaneotropica.org.br/v7n3/pt/abstract?short-communication+bn02407032007> (último acesso em 16/02/2011)
- MARINHO-FILHO, J.S., RODRIGUES, F.H.G., GUIMARAES, M.M. & REIS M.L. 1998. Os mamíferos da Estação Ecológica de Águas Emendadas, Planaltina, DF. In *Vertebrados da Estação Ecológica de Águas Emendadas - História Natural e Ecologia em um fragmento de cerrado do Brasil Central*. Brasília (J. Marinho-Filho, F.H.G. Rodrigues & M. Guimaraes, eds). SEMA TECIEMA, p. 34-63.
- OKSANEN, J., BLANCHET, F.G., KINDT, R., LEGENDRE, P., O'HARA, R.B., SIMPSON, G.L., SOLYMOS, P., STEVENS, M.H.H. & WAGNER, H. 2010. Vegan: community ecology package. R package. version 1.17-4. <http://CRAN.R-project.org/package=vegan>.
- PULCHÉRIO-LEITE, A., MENEGHELLI, M. & TADDEI, A.V. 1999. Morcegos da região de Campo Grande, Estado do Mato Grosso do Sul, com ênfase para as espécies urbanas. *Ensaios e Cienc.* 3:113-129.
- R DEVELOPMENT CORE TEAM. 2010. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <http://www.R-project.org>.
- RODRIGUES, F.H.G., SILVEIRA, L., JÁCOMO, A.T.A., CARMIGNOTTO, A.P., BEZERRA, A.M.R., COELHO, D.C., GARBOGINI, H., PAGNOZZI, J. & HASS, A. 2002. Composição e caracterização da fauna de mamíferos do Parque Nacional das Emas, Goiás, Brasil. *Rev. Bras. Zool.* 19:589-600. <http://dx.doi.org/10.1590/S0101-81752002000200015>
- SANTOS, C.F., NOGUEIRA, M., CUNHA, N., CARVALHO, L.F.A.C. & FISCHER, E. 2010. Southernmost record of the Sanborn's big-eared bat, *Micronycteris sanborni* (Chiroptera, Phyllostomidae). *Mammalia* 74:457-460. <http://dx.doi.org/10.1515/MAMM.2010.041>
- SCARIOT, A., SOUSA-SILVA, J.C. & FELFILI, J.M. 2005. Cerrado: ecologia, biodiversidade e conservação. Ministério do Meio Ambiente, Brasília. p. 430.
- SILVEIRA, M., MUNIN, R.L., TOMAS, W.M., FISCHER, E., BORDIGNON, M. & SILVEIRA, G.A. 2011. The distribution of the spectral bat, *Vampyrus spectrum*, reaches the southern Pantanal. *Biota Neotrop.*
- SIMMONS, N.B. 2005. Order Chiroptera. In *Mammal species of the world: A taxonomic and geographic reference* (D.E. Wilson & D.M. Reeder, eds). 3rd ed. Johns Hopkins University Press, Baltimore, p.312-529.
- STEVENS, R.D. 2004. Untangling latitudinal richness gradients at higher taxonomic levels: familial perspectives on the diversity of New World bat communities. *J. Biogeogr.* 31:665-674. <http://dx.doi.org/10.1111/j.1365-2699.2003.01042.x>
- STRAUBE, F.C. & BIANCONI, G.V. 2002. Sobre a grandeza e a unidade utilizada para estimar esforço de captura com utilização de redes-de-neblina. *Chir. Neotrop.* 8:150-152.
- TADDEI, V.A. 1996. Sistemática de Quirópteros. *Bol. Inst. Pasteur* 1:3-15.
- TADDEI, V.A., VIZOTTO, L.D. & SAZIMA, I. 1983. Uma nova espécie de *Lonchophylla* do Brasil e chave para identificação das espécies do gênero (Chiroptera, Phyllostomidae). *Ciênc. Cult.* 35:625-629.
- TEIXEIRA, R.C., CORRÊA, C.E. & FISCHER, E. 2009. Frugivory by *Artibeus jamaicensis* (Phyllostomidae) bats in the Pantanal, Brazil. *Stud. Neotrop. Fauna Environ.* 44:7-15. <http://dx.doi.org/10.1080/01650520802692283>
- VIZOTTO, L.D. & TADDEI, V.A. 1973. Chave para determinação de quirópteros brasileiros. *Bol. Ciências.* 1:1-72.
- WILLIG, M.R. & BLOCH, C.P. 2006. Latitudinal gradients of species richness: a test of the geographic area hypothesis at two ecological scales. *Oikos* 112:163-173. <http://dx.doi.org/10.1111/j.0030-1299.2006.14009.x>
- WILLIG, M.R., PRESLEY, S., OWEN, R. & LÓPEZ-GONZALEZ, C. 2000. Composition and structure of bat assemblages in Paraguay: a subtropical-temperate interface. *J. Mammal.* 81:386-401. [http://dx.doi.org/10.1644/1545-1542\(2000\)081%3C0386:CASOBA%3E2.0.CO;2](http://dx.doi.org/10.1644/1545-1542(2000)081%3C0386:CASOBA%3E2.0.CO;2)
- ZORTÉA, M. & ALHO, C.J.R. 2008. Bat diversity of a Cerrado habitat in central Brazil. *Biodivers. Conserv.* 17:791-805.
- ZORTÉA, M., MELO, F.R.D., CARVALHO, J.C. & ROCHA, Z.D.D. 2010. Morcegos da bacia do rio Corumbá, Goiás. *Chir. Neotrop.* 16:610-616.

Received 24/04/2011

Revised 05/07/2011

Accepted 02/09/2011