



Biota Neotropica

ISSN: 1676-0611

cjoly@unicamp.br

Instituto Virtual da Biodiversidade
Brasil

Pérez, Carlos Daniel; Braga Gomes, Paula

First record of the fireworm *Hermodice carunculata* (Annelida, Polychaeta) preying on colonies of the
fire coral *Millepora alcicornis* (Cnidaria, Hydrozoa)

Biota Neotropica, vol. 12, núm. 2, 2012, pp. 217-219

Instituto Virtual da Biodiversidade
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=199123113022>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

**First record of the fireworm *Hermodice carunculata*
(Annelida, Polychaeta) preying on colonies of the fire
coral *Millepora alcicornis* (Cnidaria, Hydrozoa)**

Pérez, C.D. & Gomes, P.B.

Biota Neotrop. 2012, 12(2): 217-219.

On line version of this paper is available from:

<http://www.biotaneotropica.org.br/v12n2/en/abstract?short-communication+bn01712022012>

A versão on-line completa deste artigo está disponível em:

<http://www.biotaneotropica.org.br/v12n2/pt/abstract?short-communication+bn01712022012>

Received/ Recebido em 09/11/11 -

Revised/ Versão reformulada recebida em 09/06/12 - Accepted/ Publicado em 15/06/12

ISSN 1676-0603 (on-line)

Biota Neotropica is an electronic, peer-reviewed journal edited by the Program BIOTA/FAPESP: The Virtual Institute of Biodiversity. This journal's aim is to disseminate the results of original research work, associated or not to the program, concerned with characterization, conservation and sustainable use of biodiversity within the Neotropical region.

Biota Neotropica é uma revista do Programa BIOTA/FAPESP - O Instituto Virtual da Biodiversidade, que publica resultados de pesquisa original, vinculada ou não ao programa, que abordem a temática caracterização, conservação e uso sustentável da biodiversidade na região Neotropical.

Biota Neotropica is an electronic journal which is available free at the following site
<http://www.biotaneotropica.org.br>

A **Biota Neotropica** é uma revista eletrônica e está integral e gratuitamente disponível no endereço
<http://www.biotaneotropica.org.br>

First record of the fireworm *Hermodice carunculata* (Annelida, Polychaeta) preying on colonies of the fire coral *Millepora alcicornis* (Cnidaria, Hydrozoa)

Carlos Daniel Pérez^{1,3} & Paula Braga Gomes²

¹*Centro Acadêmico de Vitória, Universidade Federal de Pernambuco – UFPE, CEP 55608-680, Vitória de Santo Antão, PE, Brazil*

²*Departamento de Biologia, Universidade Federal Rural de Pernambuco – UFRPE, Dois Irmãos, Recife, PE, Brazil*

³*Corresponding author: Carlos Daniel Pérez, e-mail: cdperez@ufpe.br*

PÉREZ, C.D. & GOMES, P.B. **First record of the fireworm *Hermodice carunculata* (Annelida, Polychaeta) preying on colonies of the fire coral *Millepora alcicornis* (Cnidaria, Hydrozoa).** Biota Neotrop. 12(2): <http://www.biotaneotropica.org.br/v12n2/en/abstract?short-communication+bn01712022012>

Abstract: The polychaete *Hermodice carunculata* is a voracious predator of several benthic organisms and one of the preferential groups in its diet is that of the cnidarians. This note presents the first record of a fireworm preying on the calcified hydroid *Millepora alcicornis*. The fireworm engulfs the terminal branches of the fire coral and spends several minutes scraping the surface and removing the soft tissues. Despite this being one of the most common calcified hydroids of the Brazilian northern and northeastern regions (and home to a vast associated community), this kind of association had never been recorded. This discovery points to the need of studying such associations because the damage caused by this polychaete is irreversible and depending on the amount of predators might alter reefs' health or modify the components of the reef community.

Keywords: *Amphinomidae, calcified hydroid, predation, Brazil.*

PÉREZ, C.D. & GOMES, P.B. **Primeiro registro do verme de fogo *Hermodice carunculata* (Annelida, Polychaeta) predando colônias do coral de fogo *Millepora alcicornis* (Cnidaria, Hydrozoa).** Biota Neotrop. 12(2): <http://www.biotaneotropica.org.br/v12n2/pt/abstract?short-communication+bn01712022012>

Resumo: O poliqueta *Hermodice carunculata* é um voraz predador de vários organismos bentônicos e um dos grupos preferenciais na sua dieta é o dos cnidários. No presente trabalho se apresenta o primeiro registro de predação do verme de fogo sobre o hidrocoral *Millepora alcicornis*. O verme engolfa os ramos terminais do hidrocoral e fica vários minutos raspando a superfície e retirando os tecidos moles. Apesar deste hidrocoral ser um dos representantes mais comuns dos recifes do norte e nordeste brasileiros e de albergar uma vasta comunidade associada, nunca tinha sido registrada esta associação. Esta nova descoberta alerta na necessidade de se estudar esta associação já que geralmente os danos ocasionados pelo poliqueta são irreversíveis e dependendo da quantidade de predadores pode alterar a saúde dos recifes ou modificar os componentes da comunidade recifal.

Palavras-chave: *Amphinomidae, hidroide calcário, predação, Brasil.*

The amphinomid polychaete, *Hermodice carunculata* (Pallas 1776), is a widespread resident of coral reefs and littoral areas of the Caribbean and Western Atlantic Ocean (Lizama & Blanquet 1975). It is apparently omnivorous (Marsden 1963b), feeding on sedentary animals and algae. This fireworm is regarded as an important predator on coral reefs where it grazes upon zoanthids, anemones, gorgonids, hydrocorals, scleractinians and octocorals (Lizama & Blanquet 1975, Lewis & Crooks 1996, Souza et al. 2007). Of particular interest is the fact that this polychaete feeds on living hermatypic corals – more specifically *Porites porites* and *P. astreoides* (Marsden 1962, 1963b) – and on the colonial zoanthid *Palythoa mamilliosa* (Marsden 1968). Witman (1988) and Lewis & Crooks (1996) studied this worm feeding on colonies of the calcified hydroid *Millepora complanata*, and such ecological relationship seems to be quite common in the Caribbean Sea.




Figure 1. *Hermodice carunculata* **a** in a colony of *Millepora alcicornis* at Ponta de Mangue reef, Brazil; escale bar: 8,5 cm, and **b** engulfing a terminal branch of the fire coral; escale bar: 1,7 cm.

Figure 1. *Hermodice carunculata* **a** sobre uma colônia de *Millepora alcicornis* nos recifes de Ponta de Mangue, Brasil; escala: 8,5 cm, e **b** engolindo um ramo terminal do coral de fogo; escala: 1,7 cm.

Hermodice carunculata initiates feeding by the application and attachment to the prey of the everted buccal mass. Predigestion of the coral polyps apparently results from the liberation of digestive enzymes from the buccal cells, as extracts of this region possess lipolytic and proteolytic activity (Marsden 1963a).

In coastal reefs of Ponta de Mangue, Alagoas, Brazil, several fireworms were observed feeding on *Millepora alcicornis* Linnaeus, 1758 during the morning (Figure 1a). The polychaete engulfed the terminal branches of the colony for many minutes (between 15' and 30') and remained static while removing the tissue (Figure 1b). *Millepora alcicornis* is one of the most common calcified hydroids of tropical American reefs and provides an optimal environment for epibionts (Amaral et al. 2008). Garcia et al. (2008, 2009) found a lush macrofauna associated with the colonies of *M. alcicornis* in reefs of northeastern Brazil, yet did not record the presence of *H. carunculata*. This is the first time that such a prey-predator association is being recorded between these animals in tropical Atlantic waters.

This relationship may cause permanent damage to its prey due to excessive consumption of tissue or by initiating a sequence of algal colonization in feeding lesions (Ott & Lewis 1972, Witman 1988). Excessive damage by predators may affect the general health of reefs or alter the composition of reef communities (Witman 1988). An example is the case of the crown-of-thorns starfish *Acanthaster planci* (Linnaeus 1758) considered a major predator of corals (yet a normal member of coral communities), but widespread population explosions have caused dramatic reductions of coral cover in Indo-Pacific coral reefs (Endean 1982).

In this context, in order to estimate the extent of the damage that could be caused by *H. carunculata* to corals of northeastern Brazilian reefs, we suggest that research projects be carried out to monitor and evaluate population density, their foraging cycles and the amount of time spent feeding.

Acknowledgments

We thank MSc Janine Farias, MSc David Oliveira, MSc Vithor Macedo and the sailors Carlos Silva and Carlos Eduardo for the assistance in the field work. Also, we thank Andrea Steiner (technical translator) for translation of the manuscript.

References

- AMARAL, F.M.D., STEINER, A.Q., BROADHURST, M.K. & CAIRNS, S.D. 2008. An overview of the shallow-water calcified hydroids from Brazil (Hydrozoa: Cnidaria), including the description of a new species. *Zootaxa* 1930:56-68.
- ENDEAN, R. 1982. Crown-of-thorns starfish on the Great Barrier Reef. *Endeavour* 6:10-14. [http://dx.doi.org/10.1016/0160-9327\(82\)90004-7](http://dx.doi.org/10.1016/0160-9327(82)90004-7)
- GARCIA, T.M., MATTHEWS-CASCON, H. & FRANKLIN-JUNIOR, W. 2008. Macrofauna associated with branching fire coral *Millepora alcicornis* (Cnidaria: Hydrozoa). *Thalassas* 24:11-19.
- GARCIA, T.M., MATTHEWS-CASCON, H. & FRANKLIN-JUNIOR, W. 2009. *Millepora alcicornis* (Cnidaria: Hydrozoa) as substrate for benthic fauna. *Braz. J. Oceanogr.* 57:153-155.
- LEWIS, J.B. & CROOKS, R.E. 1996. Foraging cycles of the amphinomid polychaete *Hermodice carunculata* preying on the calcareous hydrozoan *Millepora complanata*. *B. Mar. Sci.* 58:853-857.
- LIZAMA, J. & BLANQUET, R.S. 1975. Predation on sea anemones by the amphinomid polychaete, *Hermodice carunculata*. *B. Mar. Sci.* 25:442-443.
- MARSDEN, J.R. 1962. A coral eating polychaete. *Nature* 193:598. <http://dx.doi.org/10.1038/193598a0>

- MARSDEN, J.R. 1963a. A preliminary report on digestive enzymes of *Hermodice carunculata*. Can. J. Zoolog. 41:159-164. <http://dx.doi.org/10.1139/z63-019>
- MARSDEN, J.R. 1963b. The digestive tract of *Hermodice carunculata* (Pallas) Polychaeta: Amphinomidae. Can. J. Zoolog. 41:165-184. <http://dx.doi.org/10.1139/z63-020>
- MARSDEN, J.R. 1968. Routes of excretion of particulate waste in the polychaete, *Hermodice carunculata*. Can. J. Zoolog. 46:619-624. <http://dx.doi.org/10.1139/z68-088>
- OTT, B. & LEWIS, J.B. 1972. The importance of the gastropod *Coralliophila abbreviata* (Lamarck) and the polychaete *Hermodice carunculata* (Pallas) as coral reef predators. Can. J. Zool. 50:1651-1656. <http://dx.doi.org/10.1139/z72-217>
- SOUZA, J.R.B., RODRIGUES, H.A., NEVES, B.M. & PÉREZ C.D. 2007. First report of bristleworm predator of the reef octocoral *Carijoa riisei*. Coral Reef. 26:1033. <http://dx.doi.org/10.1007/s00338-007-0290-2>
- WITMAN, J.D. 1988. Effects of predation by the fireworm *Hermodice carunculata* on Milleporid hydrocorals. B. Mar. Sci. 42:446-458.

Received 09/11/2011

Revised 09/06/2012

Accepted 15/06/2012