



Biota Neotropica
ISSN: 1676-0611
cjoly@unicamp.br
Instituto Virtual da Biodiversidade
Brasil

Alves D'Avila, Frederico; de Castro Gomes, Almério
Seasonality of *Mansonia Titillans* during dam construction, Biritiba-Mirim, São Paulo State, Brazil
Biota Neotropica, vol. 13, núm. 1, 2013, pp. 69-73
Instituto Virtual da Biodiversidade
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=199126390006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

**Seasonality of *Mansonia Titillans* during dam construction,
Biritiba-Mirim, São Paulo State, Brazil**

D'Avila, F.A. & Gomes, A.C.

Biota Neotrop. 2013, 13(1): 70-73.

On line version of this paper is available from:

<http://www.biotaneotropica.org.br/v13n1/en/abstract?article+bn01213012013>

A versão on-line completa deste artigo está disponível em:

<http://www.biotaneotropica.org.br/v13n1/pt/abstract?article+bn01213012013>

Received/ Recebido em 19/01/12 -

Revised/ Versão reformulada recebida em 09/09/12 - Accepted/ Publicado em 25/01/13

ISSN 1676-0603 (on-line)

Biota Neotropica is an electronic, peer-reviewed journal edited by the Program BIOTA/FAPESP: The Virtual Institute of Biodiversity. This journal's aim is to disseminate the results of original research work, associated or not to the program, concerned with characterization, conservation and sustainable use of biodiversity within the Neotropical region.

Biota Neotropica é uma revista do Programa BIOTA/FAPESP - O Instituto Virtual da Biodiversidade, que publica resultados de pesquisa original, vinculada ou não ao programa, que abordem a temática caracterização, conservação e uso sustentável da biodiversidade na região Neotropical.

Biota Neotropica is an electronic journal which is available free at the following site
<http://www.biotaneotropica.org.br>

A **Biota Neotropica** é uma revista eletrônica e está integral e gratuitamente disponível no endereço
<http://www.biotaneotropica.org.br>

Seasonality of *Mansonia titillans* during dam construction, Biritiba-Mirim, São Paulo State, Brazil

Frederico Alves D'Avila^{1,2} & Almério de Castro Gomes[†] (in memoriam)

¹Departamento de Epidemiologia, Faculdade de Saúde Pública, Universidade de São Paulo – USP, CEP 01246-904, São Paulo, SP, Brasil

²Corresponding author: Frederico Alves D'Avila, e-mail: frdavila@ib.usp.br

D'AVILA, F.A. & GOMES, A.C. Seasonality of *Mansonia titillans* during dam construction, Biritiba-Mirim, São Paulo State, Brazil. Biota Neotrop. 13(1): <http://www.biotaneotropica.org.br/v13n1/en/abstract?article+bn01213012013>

Abstract: A two and a half year survey was conducted at a dam in southeastern Brazil. Shannon Traps were used for sampling. Kruskal-Wallis test showed little relation between rainfall and abundance. The data clearly show three abundance peaks, all of them in the end of the dry season, in consonance with the scarce literature existent. Although Kruskal-Wallis Test did not find a clear preference for the dry season, Pairwise Wilcoxon Rank Test revealed a significant difference between Fall and Spring samples. *Ma. titillans* population had a peak on late winter/early spring, close to the begin of wet season.

Keywords: *mansoniini*, insect vector, artificial lake.

D'AVILA, F.A. & GOMES, A. C. Sazonalidade de *Mansonia titillans* durante a construção de uma represa, Biritiba-Mirim, Estado de São Paulo, Brasil. Biota Neotrop. 13(1): <http://www.biotaneotropica.org.br/v13n1/pt/abstract?article+bn01213012013>

Resumo: Uma pesquisa de dois anos e meio foi realizada em uma represa no sudeste do Brasil. Foram utilizadas Armadilhas de Shannon para coleta. O teste Kruskal-Wallis mostrou pouca correlação entre a precipitação e abundância. Os dados mostram claramente três picos de abundância, todos eles no final da estação seca, em consonância com a escassa literatura existente. Apesar do teste de Kruskal-Wallis não ter encontrado uma clara preferência pela estação seca, o Teste de Wilcoxon par-a-par revelou uma diferença significativa entre as amostras do Outono e da Primavera. A população de *Ma titillans* tem um pico no fim do inverno/início da primavera, próximo ao início da estação chuvosa.

Palavra-chave: *mansoniini*, inseto vetor, lago artificial.

Introduction

Culicidae is one of the most studied insect family. Since late 19th century, when its role on yellow fever transmission (Franco 1969) was discovered. Since then many others illness were attributed as mosquito-borne, such as malaria, filariasis, and most of the caused by arboviruses.

Mansonia titillans (Walker 1848) is widely distributed in the Americas, occurring in tropical and subtropical Western Hemisphere from Southern USA to Argentina, including Caribbean Islands (Walter... 2011). It has some medical importance, as an irritating nuisance and as possible vector of Venezuelan Equine Encephalitis, which may cause sudden death and/or central nervous system disorder in these animals and also humans (Carpenter & La Casse 1955, Méndez 2001).

Mansoniini tribe has, while immatures, an unique breathing form. After hatching, the larvae attach their siphons, and themselves, to submerged roots of aquatic plants, obtaining oxygen from its aerenchymas. The pupae also remain attached to the roots of the plants until time for emergence of the adults. Water lettuce (*Pistia stratiotes* L.) and water hyacinth (*Eichornia crassipes* Kint) are the principal host plant (Harbach 2011). Due to this specificity, Mansoniini species are quite dominant where aquatic plants are abundant, normally eutrophized lakes and lagoons, frequently artificials.

Although its abundance, studies on *Ma. titillans* abundance and seasonality are scarce. This study describes the population fluctuation of this mosquito in artificial dam surroundings, near a metropolis in southeastern Brazil.

Materials and Methods

A 32-month survey was conducted from May 2002 to December 2004 at Biritiba-Mirim Dam (23° 31' 74" S and 45° 57' 14" W), in construction at the time, as part of its environmental impact analysis. At homonymous city and river, this dam is part of a water provision and flood control complex, Producer System of High Tietê (SPAT, in portuguese), in the 'Rio Tietê', about 70 km east from São Paulo city. It is a tropical region characterized by seasonal flooding, with a well defined rainfall season from November to March, accounting more than 60% of the total annual rainfall of about 1,400 mm, and a dry season from April to October. Average annual temperature is 25.9 °C (Centro... 2011).

Native vegetation on High Tietê Basin is majorly 'Mata Atlântica' rainforest. At the dam surrounding is common to find a mosaic of native vegetation and human activities, such as agriculture, loam and sand mines and eucalyptus reforestation (Centro..., 2001).

Mosquito catches were realized monthly, in a forest fragment near the construction site. Captures were carried out with Shannon Traps (Shannon 1939), gas-lamp lighted, two men operated, during the first 4 hours after dusk. Chloroform-filled deadly tubes were used to capture each specimen, it was then stored in labeled entomological boxes and carried to 'Laboratório de Entomologia em Saúde Pública (LESP) – FSP/USP'. All specimen were identified according to Forattini key (2002).

Monthly records of rainfall were obtained from a weather station near the study site. Statistical analysis was performed using data from monthly catches and climatic data through Spearman's Rank Correlation Coefficient, Pairwise Wilcoxon Rank Sum Test and Kruskal-Wallis H tests (Lowry 2012, Sachs 1992, Siegel & Castellan 1988, Spearman 1904), non-parametric, to evaluate the significance level of seasonal differences in insect abundance, and assess the influence of climatic parameters on population dynamics. Shapiro-Wilk Test (Shapiro & Wilk 1965) was used for normality test. All analysis was performed using R Statistical Computing Environment.

Results

A total of 8,783 Culicidae were caught during the study period. Mansoniini totalized 4,098 specimens, from the following species: *Coquillettidia albicosta* Peryassui 1908, *Coquillettidia chrysonotum* Peryassui 1922, *Coquillettidia hermanoi* Lane & Hermanoi 1940, *Coquillettidia juxtamansonia* Chagas 1907, *Coquillettidia venezuelensis* Theobald 1912, *Mansonia flaveola* Coquillett 1905, *Mansonia indubitans* Dyar & Shannon 1925 and *Mansonia titillans* Walker 1848.

Ma. titillans (n = 2,401) was 27.3% of all culicideans and 58.8% of Mansoniini. Its abundance was equally important among the capture sites. *Ma. titillans* females (98,4%) were more frequently seen than males in the traps.

Interspecific competition with *Coquillettidia* and other *Mansonia* species did not play a major role in this study, since relative abundance of all species of this genus was consistently lower than *Ma. titillans* during the two and a half years studied.

The weather was typical for the region during the studied period. The dry season take place between late fall and early spring, during 4 months, with some period variation. Rainy season began in September-October, and the heaviest rainfall period was from December to February (Figure 1).

Three population peaks were recorded (Figure 2), all of them occurred in September [2002] or October [2003/2004], coinciding with the end of the dry season and beginning of the wet one at this region.

First of all, the normality of the samples distribution was tested through Shapiro-Wilk Test and, as expected, normality of the sample was rejected ($p < 0.001$, $W = 0,67$).

The Kruskal-Wallis Test was performed to search for preference among the seasons. The null hypothesis, that the samples from different seasons have the same median, was accepted ($H = 7.6779$, $p = 0.05316$, $df = 3$), indicating that the seasonality of *Ma. titillans* was not directly determined by rainfall.

Spearman's rank correlation coefficient showed little correlation between rainfall and *Ma. titillans* abundance ($p = 0.173$), and also was little when compared Other Mansoniini with rainfall ($p = 0.008$) and *Ma. titillans* and other Mansoniini abundance ($p = 0.017$).

On Pairwise Wilcoxon Rank Sum Test, a significant difference was found between the Fall and Spring samples (0.049), while all other comparisons were non-significant.

Discussion

Although statistical insignificance, Kruskal-Wallis' p-value was placed at the boundary of H_0 acceptance. This data is complemented by Pairwise Wilcoxon, which had significant difference. The combination of both tests can be interpreted by *Ma. titillans* having a relative stable abundance along the year, with one peaks of great abundance. So is clear that there are cyclical peaks of abundance, somehow related to the seasons of the year.

The statistical analysis of the data did not reveal a connection between abundance and rainfall; this result can be explained by its ecology. *Ma. titillans* immature stages live on permanent breeding sites, which abundance and quality are not directly dependent of rainfall allowing *Ma. titillans* to be abundant throughout the year. An explanation on abundance peaks can be its breathing technique; using aerenchymas allow Mansoniini, and *Ma. titillans* specifically, to find they optimum environment, while others, when also living on eutrophized waters, find hard times during dry season. De Kruijf (1975) also registered an abundance peak at similar season.




Figure 1. Rainfall data in the Biritiba-Mirim Dam, State of São Paulo, Brazil, May 2002 to December 2004.




Figure 2. Seasonal fluctuation of *Mansonia titillans* in the Biritiba-Mirim Dam, State of São Paulo, Brazil, May 2002 to December 2004.

Ma. titillans is dependent of aquatic plants, which are normally perennial, so dry season may be when interspecific competition is less important and less strong.

Remarkable differences have been described regarding abundance of *Ma. titillans* in modified and primeval areas, while Taddei (1996) found relative frequency of 94%, studies in other 'Mata Atlântica' areas found much lower abundances (Guimarães et al. 2004, Reis et al. 2010, Forattini et al. 1986). This data indicates that *Ma. titillans* is a species mainly found in man-made environment, bringing, commonly, nuisance more than bearable to man and cattle, like in Tucuruí.

In conclusion, when living in optimum environment, *Ma. titillans* shows a behavior with population peak in late winter/early spring, close to the begin of wet season.

Acknowledgements

We thank CNPq for the financial support that made this research possible, EMAE for supporting the research and offering all needed infrastructure. The biologists Márcia Bicudo de Paula, Paulo Roberto Urbinatti, Delsio Natal, Aristides Fernandes, Marcelo Sciarreta for identification and capture help.

References

- CARPENTER, S.J. & LACASSE, W.J. 1955. Mosquitoes of North America (north of Mexico). Univ. of Calif. Press, Berkeley, Los Angeles, p.360.
- CENTRO DE MONITORAMENTO AMBIENTAL DA SERRA DO ITAPETI - CEMASI. 2001. Biodiversidade nas Bacias do Rio Paraitinga e do Rio Biritiba-Mirim. SERH/DAEE/CEMASI, São Paulo. Relatório Técnico

- CENTRO DE PESQUISAS METEOROLÓGICAS E CLIMÁTICAS APLICADAS À AGRICULTURA - CEPAGRI. Clima dos Municípios Paulistas. Clima de Biritiba-Mirim. http://www.cpa.unicamp.br/outras-informacoes/clima_muni_075.html (último acesso em 15/12/2011).
- DE KRUIJF, H.A.M. 1975. The relations between rainfall and mosquito populations In Tropical Ecological Systems. Trends in Terrestrial and Aquatic Research (F.B. Golley & E. Medica, eds). Ecological Studies, n.11, p.62-65.
- FORATTINI, O.P. 2002. Culicidologia Médica v.2. EdUSP, São Paulo.
- FORATTINI, O.P., ALVES, A.C., NATAL, D. & SANTOS, J.L.F. 1986. Observações sobre atividade de mosquitos Culicidae em mata primitiva da encosta no Vale do Ribeira, São Paulo, Brasil. Rev. Saude Publ. 20(1):1-20. <http://dx.doi.org/10.1590/S0034-89101986000100001>
- FRANCO, O. 1969. História da febre amarela no Brasil. Rev. Bras. Malariol. Doenças Trop. 21: 315-520. PMID:4906976.
- GUIMARÃES, A.E., LOPES C.A., MELLO R.P. & ALENCAR J. 2000. Ecologia de mosquitos (Diptera: Culicidae) em áreas do Parque Nacional da Serra da Bocaina, Brasil. 1 - Distribuição por habitat. Cad. Saude Publica 34(3):1107-1116.
- HARBACH, R.E. 2011. Tribe Mansoniini. Mosquito Taxonomic Inventory. <http://mosquito-taxonomic-inventory.info/tribe-mansoniini-belkin-1962> (último acesso em 16/12/2011).
- LOWRY, R. 2012. Concepts and Applications of Inferential Statistics. <http://vassarstats.net/textbook/ch12a.html> (último acesso em 22/04/2012)
- MÉNDEZ, W., LIRIA, J., NAVARRO, J.-C., GARCÍA, C.Z., FREIER, J.E., SALAS, R., WEAVER, S.C. & BARRERA, R. 2001. Spatial Dispersion of Adult Mosquitoes (Diptera: Culicidae) in a Sylvatic Focus of Venezuelan Equine Encephalitis Virus. J. Med. Entomol. 38(6): 813-819. PMID:11761379. <http://dx.doi.org/10.1603/0022-2585-38.6.813>
- REIS, M., AZULIM, G.M. & MARCONDES, C.B. 2010. Inventário de mosquitos (Diptera: Culicidae) da Unidade de Conservação Ambiental Desterro, Ilha de Santa Catarina, Sul do Brasil. Biota Neotrop. 10(3) <http://www.biotaneotropica.org.br/v10n3/pt/abstract?inventory+bn01610032010> (último acesso em 05/12/2011).
- SHANNON, R. 1939. Methods for collecting and feeding mosquitoes in jungle yellow fever studies. Am. J. Trop. Med. Hyg. 19:131-140.
- SHAPIRO, S.S. & WILK, M.B. 1965. An analysis of variance test for normality (complete samples). Biometrika 52(3/4):591-611. <http://dx.doi.org/10.2307/2333709>
- SIEGEL, S. & CASTELLAN, JR.NJ. 1988. Nonparametric Statistics for the Behavioral Sciences. 2. ed. McGraw-Hill, New York.
- SPEARMAN, C. 1904. The proof and measurement of association between two things. Amer. J. Psychol. 15(1):72-101. <http://dx.doi.org/10.2307/1412159>
- TADDEI, W.P. 1996. O gênero *Mansonia* (Diptera: Culicidae) e a proliferação de mosquitos na usina hidrelétrica de Tucuruí. In Energia na Amazônia (S.B. Magalhães, R.C. Brito & Castro E.R., eds). MPEG/FPA/UNAMAZ, Belém, v.1, p.311-318
- WALTER REED BIOSSYSTEMATICS UNIT. Systematic Catalog of Culicidae. http://mosquitocatalog.org/taxon_descr.aspx?ID=40 (último acesso em 13/12/2011).

Received 19/01/2012

Revised 09/09/2012

Accepted 25/01/2013