



Biota Neotropica
ISSN: 1676-0611
cjoly@unicamp.br
Instituto Virtual da Biodiversidade
Brasil

Fonseca Mendes, Humberto; Andersen, Trond
Jururumberus, a new genus of Orthoclaadiinae (Diptera:Chironomidae) from Brazil
Biota Neotropica, vol. 13, núm. 4, octubre-diciembre, 2013, pp. 71-76
Instituto Virtual da Biodiversidade
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=199130048007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

***Jururumberus*, a new genus of Orthocladiinae (Diptera: Chironomidae) from Brazil**

Humberto Fonseca Mendes^{1,3} & Trond Andersen²

¹*Centro de Ciências Naturais e Humanas – CCNH, Universidade Federal do ABC – UFABC,
Bl. A, 638-3, Rua Santa Adélia 166, Bangu, CEP 09210-170, Santo André, SP, Brazil*

²*Department of Natural History, University Museum of Bergen, University of Bergen,
CP 7800, N-5020 Bergen, Norway*

³*Corresponding author: Humberto Fonseca Mendes, e-mail: orthocladiinae@gmail.com*

MENDES, H.F., & ANDERSEN, T. ***Jururumberus*, a new genus of Orthocladiinae (Diptera: Chironomidae) from Brazil.** 13(4): <http://www.biotaneotropica.org.br/v13n4/en/abstract?article+bn01213042013>

Abstract: *Jururumberus* n. gen. is erected based on the males of *J. arapuca* n. sp. from São Paulo State, in southeastern Brazil and *J. uatuma* n. sp. from the Amazonas in northern Brazil. The genus can be separated from other Orthocladiinae based on the combination of bare eyes, wing, and squama; frons U-shaped, extended ventrally; $AR \leq 0.3$, antenna without apical seta; palp short, five-segmented, with sensilla clavata in subapical sensillum coeloconicum; no acrostichals; costa strongly extended; R_{4+5} ending proximal to M_{3+4} ; anal point narrowly triangular with microtrichia only, sitting high on tergite IX; and virga composed of two spines.

Keywords: Chironomidae, Orthocladiinae, *Jururumberus*, new genus, new species, Brazil, Neotropical region, Mata Atlântica, Amazon.

MENDES, H.F. & ANDERSEN, T. ***Jururumberus*, um gênero novo de Orthocladiinae (Diptera: Chironomidae) do Brasil.** 13(4): <http://www.biotaneotropica.org.br/v13n4/pt/abstract?article+bn01213042013>

Resumo: *Jururumberus* n. gen. é descrito baseado em machos de *J. arapuca* n. sp. do Estado de São Paulo, no sudeste brasileiro e *J. uatuma* n. sp. do Estado do Amazonas no norte brasileiro. O gênero pode ser separado dos demais Orthocladiinae com base na combinação dos seguintes caracteres: olhos, asas e *squama* nus; fronte em forma de ‘U’, estendido ventralmente; $AR \leq 0.3$, antena sem seta apical; palpo encurtado, com cinco segmentos, com sensilas clavadas em uma sensila celocônica subapical; acrosticais ausentes; veia costal fortemente estendida; R_{4+5} terminando proximal em relação a M_{3+4} ; ponta anal estreita e triangular com apenas microtríquios, situada oralmente no tergito IX; virga composta de dois espinhos.

Palavras-chave: Chironomidae, Orthocladiinae, *Jururumberus*, gênero novo, espécie nova, Brasil, região Neotropical, Mata Atlântica, Amazônia.

Introduction

In their catalog of the Neotropical and Mexican chironomids Spies & Reiss (1996) recorded seven Orthoclaadiinae species from Brazil; one of these, *Ichthyoclaadius neotropicus* Fittkau, as uncertain, a species which later has proved not to occur in Brazil (Mendes et al. 2004). Ten years later Mendes et al. (2007) listed 37 species of orthoclads, while today about 100 species are known to occur in Brazil (Mendes & Pinho 2011). Several genera are so far only recorded from Brazil, viz. *Gynoclaadius* Mendes, Sæther et Andrade-Morraye, *Iporangomberus* Mendes et Andersen, *Lyrocladius* Mendes et Andersen, *Miambera* Andersen et Mendes, *Maximberus* Andersen et Mendes, *Oleia* Andersen et Mendes, *Saetheroclaadius* Andersen et Mendes, *Saetherocryptus* Andersen et Mendes, *Saetherolabis* Andersen et Mendes, *Saetherops* Andersen et Mendes, and *Ubatubaneura* Wiedenbrug et Trivinho-Strixino, but a wider distribution of several of these genera is to be expected.

However, a high number of new species still await description. When collecting in Brazil many of the orthoclads encountered do not readily fit into any described genus. Several of these might be terrestrial or semi-terrestrial and the larvae and pupae might be difficult to locate. Below we describe the males of two of these species and place them in a new genus. One of these species was taken in Boracéia Biological Reserve in São Paulo State in the coastal rainforest Mata Atlântica. The other species was collected in Uatuma Biological Reserve in the Amazonas State, in the Amazonian rainforest. Both species were collected in light traps.

Material and Methods

The specimens were mounted in Canada Balsam or Euparal following the procedures outlined by Sæther (1969). The general morphology follows Sæther (1980). Measurements are given as ranges.

The types will be deposited in Museu de Zoologia da Universidade de São Paulo (MZUSP) and in the Department of Natural History, University Museum of Bergen, University of Bergen, Bergen, Norway (ZMBN).

1. *Jururumberus* new genus

Type species: *Jururumberus arapuca* n. sp.

Other included species: *Jururumberus uatuma* n. sp.

Etymology: Deriving from the South American native language Tupi *yuru-ru* and *mberui*, meaning melancholic, sad and midge, mosquito, respectively, due to their gloomy facial expression. Gender of the genus name: masculine.

Diagnostic characters: The combination of bare eyes, wing, and squama; frons U-shaped, extended ventrally; $AR \leq 0.3$, antenna without apical seta; palp short, five-segmented, with sensilla clavata in subapical sensillum coeloconicum; no acrostichals; costa strongly extended; R_{4+5} ending proximal to M_{3+4} ; anal point narrowly triangular with microtrichia only, sitting high on tergite IX; and virga composed of two spines will separate the genus from all other orthoclads.

1.1. Description

Small sized species, wing length 0.67-0.87 mm.

Head. Eyes bare, reniform, without dorsomedian extension. Antenna with 12 flagellomeres, $AR \leq 0.30$; sparsely plumed; groove beginning on segment 1; sensilla chaetica present on flagellomeres 1-3 and ultimate; without subapical seta. Palp five-segmented, short; third palpomere with 5-7 sensilla clavata in subapical sensillum coeloconicum. Temporal setae in single row, all comparatively long. Frons U-shaped, extended ventrally; frontal tubercle absent.

Tentorium and stipes normal. Cibarial pump with anterior margin weakly convex. Clypeus with few setae.

Thorax. Anteprenotal lobes widely separated medially, with few, weak lateral anteprenotals. Acrostichals absent; dorsocentrals few, uniserial; prealars few; supraalar absent. Scutellum with few setae in single row.

Wing. Membrane without setae, with fine punctuation. Anal lobe weakly developed. Costa strongly extended; R_{2+3} running and ending midway between R_1 and R_{4+5} ; R_{4+5} ending proximal to M_{3+4} ; FCu distal to RM; Cu_1 slightly curved. Brachiolium with 1 seta, other veins bare. Squama bare. Sensilla campaniformia about 6 apically, 5 basally, and 2 above seta on brachiolium; 1 on RM; and 1 basally on R_1 .

Leg. Tibial spurs normal, comb of hind tibia composed of 9-12 setae. Tarsal pseudospurs and sensilla chaetica absent. Pulvilli vestigial.

Abdomen. Abdominal setation reduced. Tergite I bare, tergites II-III bare or with two irregular rows of few setae, tergites IV-VIII with two irregular rows of few setae. Sternites I-II bare, sternites II-IV bare or with single median setae, sternites V-VIII with single or few setae medially in two irregular rows.

Hypopygium. Anal point narrowly triangular, sitting high on tergite IX, with microtrichia only. Tergite IX with few long setae; laterosternite IX with few setae. Apodemes well sclerotized. Phallapodeme with normal aedeagal lobe. Transverse sternapodeme straight to weakly arched, oral projections distinct. Virga consisting of 2 separate spines. Gonocoxite with well developed inferior volsella, with posterior, free lobe. Gonostylus club-shaped, covered with microtrichia and with few, weak setae; crista dorsalis present; megaseta normal.

Female, pupa and larva. Unknown.

1.2. Systematics

The new genus will key to couplet 97 in the key to the genera of the Holarctic Region (Cranston et al. 1989), but no further as the costa is strongly extended and the anal point is hardly extending beyond the posterior margin of tergite IX nor originating from the posterior margin of the tergite. In the key to the Central American Chironomidae (Spies et al. 2009) the genus will key to couplet 122, but no further as the squama is lacking setae and scutum is without a medial tuft of microtrichia.

The genus is difficult to place, but most probably it groups with *Pseudosmittia* Edwards, further discussion on the group and its Neotropical members in Andersen & Mendes (2007).

2. *Jururumberus arapuca* n. sp. (Figures 1-6)

Type material: Holotype male: BRAZIL, São Paulo State, Salesópolis, Estação Biológica Boracéia, 10.xii.2001, H.F. Mendes et al. leg. (MZUSP). Paratypes: 1 male, same as holotype (MZUSP); 1 male, same as holotype except 12.xii.2001, C.G. Froehlich et al. leg. (ZMBN).

Etymology: From Tupi *ara* and *puka* meaning respectively light and trap due to the method the type material was sampled. The name is to be regarded as a noun in apposition.

Diagnostic characters: The species is larger than *J. uatuma* n. sp., the wing length is > 0.81 mm compared to 0.67 mm in *J. uatuma* n. sp. The posterior lobe of inferior volsella is broadly rounded, while it is nearly parallel-sided in *J. uatuma* n. sp. and the crista dorsalis is long and low, while it is triangular and placed subapically in *J. uatuma* n. sp.

2.1. Description

Male (n = 2-3). Total length 1.33-1.41 mm. Wing length 817-866 µm. Total length / wing length 1.59-1.71. Wing length / length of profemur 2.94-3.23.

Coloration. Head, pedicel, thorax and abdominal segments V-VIII dark brown; legs and antennal flagellum lighter brown; abdominal segments I-IV pale brown; wings translucent.

Head (Figure 1). Antennae with 12 flagellomeres; AR 0.28-0.30; ultimate flagellomere (Figure 2) 65-67 µm long. Temporal setae 7-8 including 4 inner verticals, 2-3 outer verticals, and 1-2 postorbital. Clypeus with 5-6 setae. Tentorium, stipes, and cibarial pump as in Figure 3. Tentorium 80-87 µm long; 11-13 µm wide. Stipes 52-63 µm long. Palp segment lengths / widths (in µm): 6-8 / 8-11; 12-14 / 18-19; 25-32 / 19-22; 23-28 / 12-14; 33-37 / 7-8. Third palpomere with 5-7 sensilla clavata in subapical sensillum coeloconicum, 12-14 µm long.

Thorax. Anteprenotum with 2 setae. Dorsocentrals 7-8; prealars 2-3. Scutellum with 4 setae.

Wing (Figure 4). VR 1.41-1.55. Costal extension 158-168 µm long. Brachiolum with 1 seta, remaining veins and cells bare. Squama bare.

Legs. Spur of fore tibia 25-26 µm long; spurs of mid tibia 14-15 µm and 12-14 µm long; spurs of hind tibia 38-40 µm and 15-17 µm long. Width at apex of fore tibia 18-20 µm; of mid tibia 17-21 µm; of hind tibia 28-33 µm. Comb with 11-12 setae; longest 32-36 µm long; shortest 19-21 µm long. Lengths and proportions of legs as in Table 1.

Abdomen. Strong setae on tergites I-VIII as: 0, 4-5, 6-8, 6-8, 7-11, 7-12, 6-10, 7-9. Strong setae on sternites I-VIII as: 0, 0, 0-1, 1-2, 2-6, 3-6, 7-9.

Hypopygium (Figures 12-13). Anal point covered with microtrichia; 15-22 µm long; 4-6 µm wide at base. Tergite IX with 6-8 setae; laterosternite IX with 2 setae. Transverse sternapodeme 50-53 µm long. Phallapodeme 48-50 µm long. Virga with 2 spines, longest 10-11 µm long. Gonocoxite 131-137 µm long. Inferior volsella with 7-8 µm long, 10-11 µm wide, broadly rounded lobe projecting caudad. Gonostylus 55-61 µm long, with long, low crista dorsalis, about 30 µm long, 3 µm high; megaseta 8-11 µm long. HR 2.25-2.38. HV 2.32-2.41.

2.2. Distribution and habitat

The type material was collected in light traps during an expedition to the Biological Station of Boracéia, Salesópolis municipality, São

Paulo State, in the Mata Atlântica forest. This nature reserve was created in 1954 and comprises an area of 96 hectares 110 km east of São Paulo city. It is composed of both primary and secondary forests under the protection of the Museu de Zoologia da Universidade de São Paulo (MZUSP: http://www.mz.usp.br/portugues/dci/estacao_biologica_boraceia/estacao_biologica_boraceia.html).

3. *Jururumberus uatuma* n. sp. (Figures 7-13)

Type material: Holotype male: BRAZIL, Amazonas State: Reserva Biológica Uatuma, Waterfall camp, 31.i.-02.ii.2008, light trap, N. Hamada *et al. leg.* (INPA).

Etymology: The species is named after the type locality. The name is to be regarded as a noun in apposition.

Diagnostic characters: See diagnostic characters for *J. arapuca* n. sp.

3.1. Description

Male (n = 1). Total length 1.28 mm. Wing length 670 µm. Total length / wing length 1.90. Wing length / length of profemur 2.73.

Coloration. Head, pedicel and abdominal segments V-VIII brown; thorax brown with slightly darker brown vitae, median anepisternum, preepisternum, scutellum and postnotum; legs and antennal flagellum lighter brown; abdominal segments I-IV pale brown; wings translucent.

Head. Antennae with 12 flagellomeres; AR 0.27; ultimate flagellomere 97 µm long. Temporal setae 8 including 5 inner verticals, 2 outer verticals, and 1 postorbital. Clypeus with 6 setae. Tentorium 82 µm long, 14 µm wide. Stipes 55 µm long. Palp segment lengths / widths (in µm): 6 / 8, 18 / 14, 30 / 22, 28 / 13, 26 / 8. Third palpomere with about 6 sensilla clavata in subapical sensillum coeloconicum, about 11 µm long.

Thorax (Figure 8). Anteprenotum with 2 setae. Dorsocentrals 6; prealars 3. Scutellum with 4 setae.

Wing (Figure 7). VR 1.47. Costal extension 135 µm long. Brachiolum with 1 seta, remaining veins and cells bare. Squama bare.

Legs (Figures 9-11). Spur of fore tibia 19 µm long, spurs of mid tibia 12 µm and 11 µm long, spurs of hind tibia 36 µm and 14 µm long. Width at apex of fore tibia 18 µm, of mid tibia 21 µm, of hind tibia 26 µm. Comb with 9 setae, longest 28 µm long; shortest 18 µm long. Lengths and proportions of legs as in Table 2.


Abdomen. Strong setae on tergites I-VIII as: 0, 0, 0, 1, 5, 7, 4, 5. Strong setae on sternites I-VIII as: 0, 0, 0, 0, 1, 2, 3, 7.

Table 1. Lengths (in µm) and proportions of legs of *Jururumberus arapuca* n. gen., n. sp., male (n = 2-3).


	fe	ti	ta ₁	ta ₂	ta ₃	ta ₄
p ₁	292-296	328-336	200-204	84-88	56-60	36-40
p ₂	356-360	332-340	148-160	60-68	44-48	31-33
p ₃	320-324	332-340	192-208	80-84	96-100	35-37
	ta ₅	LR	BV	SV	BR	
p ₁	29-34	0.61-0.62	3.80-4.02	3.06-3.10	2.66-3.12	
p ₂	27-29	0.44-0.47	4.24-5.04	4.35-4.70	2.44-3.44	
p ₃	31-33	0.57-0.61	3.46-3.47	3.19-3.40	3.20-3.40	

Table 2. Lengths (in µm) and proportions of legs of *Jururumberus uatuma* n. gen., n. sp., male (n = 1).

	fe	ti	ta ₁	ta ₂	ta ₃	ta ₄	ta ₅	LR	BV	SV	BR
p ₁	240	276	160	56	44	28	24	0.58	4.45	3.23	2.6
p ₂	296	280	120	48	40	24	20	0.43	5.27	4.80	2.2
p ₃	272	284	-	-	-	-	-	-	-	-	-



Figures 1-6. *Jururumberus arapuca* n. gen.; n. sp., male. 1) Head. 2) Ultimate flagellomeres of antenna. 3) Tentorium, stipes and cibarial pump. 4) Wing. 5) Hypopygium, dorsal view. 6) Hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.



Figures 7-13. *Jururumberus uatuma* n. gen.; n. sp., male. 7) Wing. 8) Thorax. 9) Apex of fore tibia. 10) Apex of mid tibia. 11) Apex of hind tibia. 12) Hypopygium, dorsal view. 13) Hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Hypopygium (Figures 12-13). Anal point covered with microtrichia, 22 µm long, 5 µm wide at base. Tergite IX with 7 setae including 3 sitting at base of anal point; laterosternite IX with 2 setae. Transverse sternapodeme 33 µm long. Phallapodeme 46 µm long. Virga of 2 weak spines, longest 5 µm long. Gonocoxite 104 µm long. Inferior volsella with 10 µm long, 6 µm wide, parallel-sided lobe projecting caudad. Gonostylus 43 µm long, with triangular subapical crista dorsalis, 7 µm wide, 5 µm high; megaseta 7 µm long. HR 2.42. HV 2.98.

3.2. Distribution and habitat

This species is known only from the holotype collected close to a waterfall in a federal nature reserve located in Presidente Figueiredo (REBIO Uatumã), Amazonas State, in the Amazon forest. The reserve was created in 1990 and comprises an area of 9500 hectares and has on its preservation list the Amazonian manatee, *Trichechus inunguis* (Natterer, 1883) and the freshwater sponge *Anheteromeyenia ornata* (Bonetto *et* Ezcurra de Drago, 1970). This freshwater sponge is protected only in this nature reserve and deserves special attention.

Acknowledgements

We are indebted to Dr. Claudio Gilberto Froehlich, FFCLRP/USP for arranging the expedition to the type locality of *Jururumberus arapuca* gen. n., n. sp. and Dr. Neusa Hamada, INPA, for the material of *J. uatuma* n. sp. Gladys Ramirez made the slide preparations. The study was partially funded by FAPESP (11/50162-1), partially by the University Museum of Bergen.

References

- ANDERSEN, T. & MENDES, H.F. 2007. Five enigmatic new orthoclad genera from Brazil (Diptera: Chironomidae, Orthoclaadiinae). In Contributions to the Systematics and Ecology of Aquatic Diptera – A tribute to Ole A. Sæther. (T. Andersen, ed.). The Caddis Press, Columbus, p.17-52.
- CRANSTON, P.S., OLIVER, D.R. & SÆTHER, O.A. 1989. The adult males of Orthoclaadiinae (Diptera: Chironomidae) of the Holarctic region. Keys and diagnoses. In Chironomidae of the Holarctic region. Keys and diagnoses. Part 3. Adult males (T. Wiederholm, ed.). Ent. scand., Suppl. 34:165-352.
- MENDES, H.F. & PINHO, L.C. 2011. Brazilian chironomid home page. <http://sites.google.com/site/brazilianchironomids/> (último acesso em 12/06/2013)
- MENDES, H. F., ANDERSEN, T. & PINHO, L.C. 2007. Taxonomic research on Brazilian Orthoclaadiinae (Diptera: Chironomidae) during the last decade. Entomol. Mex. 6:1173-1177.
- MENDES, H.F., ANDERSEN, T. & SÆTHER, O.A. 2004. New species of *Ichthyocladus* Fittkau, a member of the *Corynoneura*-group (Diptera: Chironomidae: Orthoclaadiinae), with a review of the genus. Stud. Neotrop. Fauna Environ. 39:15-35. <http://dx.doi.org/10.1080/01650520412331270936>
- SÆTHER, O.A. 1969. Some Nearctic Podonominae, Diamesinae and Orthoclaadiinae (Diptera: Chironomidae). Bull. Fish. Res. Board Canada 107:1-154.
- SÆTHER, O.A. 1980. Glossary of Chironomid morphology terminology (Diptera: Chironomidae). Ent. Scand., Suppl. 14:1-51.
- SPIES, M. & REISS, F. 1996. Catalog and bibliography of neotropical and Mexican Chironomidae (Insecta, Diptera). Spixiana, Suppl. 22:61-119.
- SPIES, M., ANDERSEN, T., EPLER, J.H. & WATSON JUNIOR, C.N. 2009. Chironomidae (Non-biting midges). In Manual of Central American Diptera (B.V. Brown, A. Borkent, J.M. Cumming, D.M. Wood, N.E. Woodley & M.A. Zumbado, eds.). NRC Research Press, Ottawa, p.437-480.

Received 06/12/2012

Revised 09/12/2013

Accepted 10/25/2013