



Psychology & Neuroscience

ISSN: 1984-3054

landeira@puc-rio.br

Pontifícia Universidade Católica do Rio de
Janeiro
Brasil

S. Ribeiro, Jefferson A.; da Silva, José A.; Y. Katayama, Bianka Y. N.; Jorge, Rodrigo
Pain in the treatment of vitreoretinal diseases
Psychology & Neuroscience, vol. 7, núm. 1, 2014, pp. 27-31
Pontifícia Universidade Católica do Rio de Janeiro
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=207029991005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Pain in the treatment of vitreoretinal diseases

Jefferson A. S. Ribeiro^{1,2}, José A. da Silva³, Bianka Y. N. Y. Katayama¹, and Rodrigo Jorge¹

1. University of São Paulo, Ribeirão Preto, SP, Brazil

2. University of the State of Amazonas, Manaus, AM, Brazil

3. University of São Paulo, São Paulo, SP, Brazil

Abstract

Treatment of vitreoretinal diseases comprises a range of options that has increased in the past several years, allowing more individuals to benefit from new therapeutic approaches. The use of lasers in the retina, new surgical techniques, and intravitreal drug administration constitute common treatment options. These retinal and vitreal procedures have both benefits and drawbacks. Among the drawbacks is discomfort associated with the procedure, constituting a relevant issue because it can compromise adherence to treatment. We reviewed aspects of pain associated with vitreoretinal procedures and the available options for its control. With regard to vitreoretinal procedures, laser photocoagulation is associated with a higher pain score compared with vitrectomy and intravitreal injection using traditional anesthesia methods, suggesting that laser photocoagulation still needs to be improved with regard to comfort during the procedure. In some cases, a combination of analgesic options may be the best choice for pain-sensitive individuals. Individual differences and the specific condition that is treated should be considered when deciding the best treatment option. **Keywords:** pain, vitrectomy, laser photocoagulation, intravitreal injection, treatment.

Received 10 April 2013; received in revised form 17 September 2013; accepted 05 November 2013. Available online 07 March 2014.

Introduction

Treatment of vitreoretinal diseases comprises a range of options that has increased in the past several years, allowing more individuals to benefit from new therapeutic methods. The use of lasers in the retina, new surgical techniques, and intravitreal drug administration constitute common treatment options. The increase in the number of invasive procedures in the retina has both benefits and drawbacks. Among the drawbacks is discomfort associated with the procedure, constituting a relevant issue because it can compromise adherence to treatment. Therefore, pain associated with vitreoretinal procedures and their treatment options have been the focus of several ophthalmological studies.

Ophthalmological procedures

Among the procedures most commonly used in the treatment of vitreoretinal diseases are pars plana vitrectomy, laser photocoagulation, and intravitreal drug administration.

Laser

Photocoagulation of the retina using lasers can be used to treat various vascular diseases in the retina such as diabetic retinopathy (Diabetic Retinopathy Study Research Group, 1978; Early Treatment Diabetic Retinopathy Study Research Group, 1991), vascular occlusions (Central Vein Occlusion Study Group, 1993; Branch Vein Occlusion Study Group, 1986), retina ruptures, and peripheral degeneration (Pollak & Oliver, 1981), and macular diseases such as diabetic macular edema (Early Treatment Diabetic Retinopathy Study Research Group, 1985).

Two laser wavelengths are most commonly used, infrared and green. The type of wavelength can be associated with greater discomfort caused by pain that occurs during the procedure. The use of infrared lasers is associated with greater pain intensity compared with green lasers (Lira, Nascimento, Arieta, de Carvalho, & Silva, 2010).

Pain is an important complication associated with retinal panphotocoagulation (Hitchings, 1980). The pain associated with this procedure tends to be more salient in the superior temporal quadrant, particularly the regions at 3, 9, and 12 hours that correspond to the long ciliary nerves and anterior to the equator.

Jefferson A. S. Ribeiro (M.D.), Bianka Y. N. Y. Katayama (M.D.), and Rodrigo Jorge (M.D., Ph.D.), Department of Ophthalmology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, Brazil. Jefferson A. S. Ribeiro, Superior School of Health Sciences, University of the State of Amazonas, Manaus, AM, Brazil. José A. Da Silva (Ph.D.), Department of Psychology, School of Philosophy, Sciences, and Literature, University of São Paulo, Brazil. Correspondence regarding this article should be directed to: Dr. Rodrigo Jorge, Division of Ophthalmology, Ribeirão Preto Medical School, Avenida Bandeirantes, 3900, Ribeirão Preto, São Paulo, 14049-900, Brazil. Phone: +55-16-36022523. Fax: +55-16-36022860. E-mail: rjorge@fmrp.usp.br

However, the treatment of conditions that affect the posterior pole such as diabetic macular edema has been associated with a lower incidence of pain (Weinberger, Ron, Lichter, Rosenblat, Axer-Siegel, & Yassur, 2000). Pain frequently irradiates in areas such as the ipsilateral upper eyelid and frontal, parietal, and temporal regions and is particularly intense in the superior orbital edge and temporal region immediately following the lateral canthus (Tamai & Mizuno, 1984). To reduce pain during retinal photocoagulation, the laser pulse has been modified with regard to the way the energy is delivered, in which the wave amplitude of the laser is changed (Friberg & Venkatesh, 1995). The exposure time can also be reduced with a proportional increase in potency such that a shorter laser exposure time is associated with a lower pain sensation. Importantly, this benefit can be obtained without compromising the efficacy of treatment (Al-Hussainy, Dodson, & Gibson, 2008).

Outpatient treatment with lasers is commonly performed with the use of local anesthesia by means of instilling eye drops. However, different drugs and administration routes have also been used to produce analgesia during laser photocoagulation, including the injection of drugs (e.g., lidocaine and ropivacaine) in peri- or retrobulbar areas (Wu et al., 2006; Kallio, Puska, Summanen, Paloheimo, & Maunukse, 1999) and intramuscular, subtenonian, and subconjunctival injections (Wu et al., 2006; Stevens, Foss, & Hamilton, 1993; Schlote, & Derse, 2001; Patel, Jenkins, Benjamin, & Webber, 2006; Tesha et al., 2010). Moreover, the local and oral administration of nonsteroidal antiinflammatory drugs (NSAIDs), anxiolytics (alone or in combination), and inhalant analgesics have been used (Wu et al., 2006; Weinberger et al., 2000; Vaideanu, Taylor, McAndrew, Hildreth, Deady, & Steel, 2006; Cook, Newsom, Mensah, Saeed, James, & Ffytche, 2002).

The administration of NSAIDs before the procedure is one option for alleviating pain. The use of diclofenac sodium (0.1%) eye drops prior to laser photocoagulation reduces pain associated with panphotocoagulation in the treatment of proliferative diabetic retinopathy (Weinberger et al., 2000). In another study, oral diclofenac potassium (50 mg) administered 45 min prior to retinal panphotocoagulation in individuals with diabetic retinopathy was superior to placebo in reducing procedural pain (de Faria Rodrigues, Messias, Da Silva, Ribeiro, Jorge, & Scott, 2010).

Intravitreal injection of drugs

The administration of drugs via intravitreal injection has become a common procedure in the age of pharmacomodulation (Anijeet, Hanson, Bhagey, & Bates, 2007; Fung, Rosenfeld, & Reichel, 2006; Aiello et al., 2004; Jager, Aiello, Patel, & Cunningham, 2004). It is a minimally invasive and efficacious intervention in the treatment of several vitreoretinal diseases (Jaissle, Szurman, & Bartz-Schmidt, 2005). The intravitreal injection represents a direct route of

drug administration in the posterior segment of the eye. The use of this technique results in immediate levels of high vitreal concentrations of various drugs including antibiotics, antiviral agents (Taskintuna et al., 1997), antifungal agents (Scott, Cruz-Villegas, Flynn, & Miller, 2004), steroids (Bonini-Filho, Jorge, Barbosa, Calucci, Cardillo, & Costa, 2005), and agents that act against vascular endothelial growth factor (VEGF; Gragoudas, Adamis, Cunningham, Feinsod, & Guyer, 2004; Rosenfeld, Moshfeghi, & Puliafito, 2005; Bonini-Filho, Costa, Calucci, Jorge, Melo, & Scott 2009).

The type of anesthetic used for intravitreal injections varies according to the experience of the surgeon. The methods employed for anesthesia during intravitreal injection include topical administration of eye drops and gels (Friedman & Margo, 2006; Kozak, Cheng, & Freeman, 2005), peribulbar and subconjunctival injections (Kishore, Conway, & Peyman, 2001; Kaderli & Avci, 2006), and the use of cotton-tipped applicators soaked with anesthetic and applied to the injection site for a short period of time prior to the injection (Kaderli & Avci, 2006). Due to the fact that multiple intravitreal injections may be necessary to maintain the desired effect in some conditions such as age-related macular degeneration (Gragoudas et al., 2004; Rosenfeld et al., 2006), the patient's comfort becomes a relevant issue.

To determine the best treatment options, studies have compared three types of anesthesia (Cintra, Lucena, Da Silva, Costa, Scott, & Jorge, 2009): local anesthesia using proximetacaine chloridrate (0.5%) drops applied using a cotton-tipped applicator soaked for 30 s, subconjunctival injection of 0.4 ml xylocaine (2%), and peribulbar injection of 4 ml xylocaine (2%). The pain that was exclusively associated with the intravitreal injection was lower in the group that received peribulbar anesthesia, whereas subconjunctival and local anesthesia had similar effects. However, the pain associated with the entire procedure (anesthesia + intravitreal injection) was higher in the peribulbar group, demonstrating that the local route with eye drops applied using an anesthetic-soaked cotton-tipped applicator may cause less pain during the entire procedure and reduce the frequency of subconjunctival hemorrhage compared with subconjunctival injection of the anesthetic.

The spectrum of indications of intravitreal injections has increased in recent years, making it an option or adjuvant for treatments that use laser photocoagulation for diabetic retinopathy and venous occlusions of the retina (Tonello, Costa, Almeida, Barbosa, Scott, & Jorge, 2008; Costa, Jorge, Calucci, Melo, Cardillo, & Scott, 2007). Thus, comparing the discomfort associated with these two treatment options would be interesting, which may guide the choice of the method used. A recent study investigated patients with proliferative diabetic retinopathy and persistent retinal neovascularization who could receive retreatment with laser or intravitreal anti-VEGF injection (Lucena et al., 2013). The study reported that retreatment with the injection was more comfortable than with the laser, in

addition to having beneficial effects on the regression of neovascularization and an improvement in visual acuity.

Pars plana vitrectomy

When vitrectomy surgery was initially introduced, the principal indications were severe and persistent vitreous hemorrhage and tractional retinal detachment that involved the center of the macula (Aaberg, 1977). With the advance of surgical techniques and reduction of severe complications, new indications have emerged (Ramsay, Knobloch, & Cantrill, 1986; Bustros, Thompson, Michels, & Rice, 1987). One of these technical improvements is the use of smaller incisions (i.e., 23 and 25 gauge) as opposed to 20 gauge, allowing faster surgeries and less manipulation of the ocular surface (Chen, 1996; Kwok et al., 1999a; Jackson, 2000; Theelen, Verbeek, Tilanus, & van den Biesen, 2003; Schmidt, Nietgen, & Brieden, 1999). A shift has occurred from the initial trend of performing vitreoretinal procedures under general anesthesia to the use of local anesthesia (e.g., peribulbar, retrobulbar, and subtenonian); (Rao, Wong, Groenewald, McGalliard, Jones, & Ridges, 1998; Kirkby, Benson, Callear, & Loo, 1999; Javitt, Addiego, Friedberg, Libonati, & Leahy, 1987; Duker et al., 1991; Demediuk, Dhaliwal, Papworth, Devenyi, & Wong, 1995; Davis & Mandel, 1986; Benedetti & Agostini, 1994; Batterbury, Wong, Williams, Kelly, & Mostafa, 1992; Stevens, Franks, Orr, Leaver, & Cooling, 1992; Kwok et al., 1999b). Some studies have reported the use of local anesthesia that produces adequate analgesia while maintaining ocular movement during the procedure (Theocharis, Alexandridou, & Tomic, 2007; Tang, Lai, Lai, Zou, Li, & Li, 2007).

Peribulbar, subtenonian, and retrobulbar anesthesia appears to be equally effective in pain control during surgical procedures (Roman, Chong Sit, Boureau, Auclin, & Ullern, 1997; Newson, Wainwright, & Canning, 2001; Lai et al., 2005). Another point that should be considered with regard to anesthesia during surgical procedures is preemptive analgesia for vitrectomy. Peribulbar injections of anesthetics prior to general anesthesia surgery appear to be a good option for decreasing postoperative discomfort in patients who undergo vitreoretinal surgery compared with general anesthesia alone (Schönfeld, Hierneis, & Kampik, 2012; Kristin et al., 2001).

Final considerations

With regard to vitreoretinal procedures, laser photocoagulation is associated with the highest pain score (Tesda et al., 2010; Hillier, Aboud, Thind, & Clark, 2009; de Faria Rodrigues et al., 2010; Lucena et al., 2013), whereas vitrectomy and intravitreal injections are less painful (Yau, Jackman, Hooper, & Sheidow, 2011; Blaha, Tilton, Barouch, & Marx, 2011; Lucena et al., 2013; Lai et al., 2005). Notably, such a comparison is made by measuring pain after completion of a procedure that used traditional methods

of anesthesia, suggesting that laser photocoagulation still needs to be improved with regard to procedural comfort.

The complexity of options for the treatment of retina and vitreous diseases requires knowledge of the various aspects of available therapeutics and methods that may be useful for facilitating patient adherence to treatment by reducing discomfort and pain. In some cases, a combination of analgesic options may be the best choice in individuals who are more susceptible to pain, and each case should be considered individually. Therapeutic options that cause less patient discomfort, regardless of the analgesic technique, may also be used, provided they are proven to be as effective as traditional treatments.

References

- Aaberg, T. M. (1977). Vitrectomy for diabetic retinopathy. In: H. Mackenzie Freeman, T. Hirose, & C.L. Schepens (Eds.), *Vitreous surgery and advances in fundus diagnosis and treatment* (pp. 297-313). New York: Appleton-Century-Crofts.
- Anijeet, D. R., Hanson, R. J., Bhagey, J., & Bates, R. A. (2007). National survey of the technique of intravitreal triamcinolone injection in the United Kingdom. *Eye*, 21, 480-486.
- Aiello, L. P., Brucker, A. J., Chang, S., Cunningham, E. T., Jr., D'Amico, D. J., Flynn, H. W., Jr., ... Trese, M. T. (2004). Evolving guidelines for intravitreal injections. *Retina*, 24(5 Suppl.), S3-S19.
- Al-Hussainy, S., Dodson, P. M., & Gibson, J. M. (2008). Pain response and follow-up of patients undergoing panretinal laser photocoagulation with reduced exposure times. *Eye*, 22, 96-99.
- Batterbury, M., Wong, D., Williams, R., Kelly, J., & Mostafa, S. M. (1992). Peribulbar anaesthesia: Failure to abolish the oculocardiac reflex. *Eye*, 6, 293-295.
- Benedetti, S., & Agostini, A. (1994). Peribulbar anesthesia for vitreoretinal surgery. *Retina*, 14, 277-280.
- Blaha, G. R., Tilton, E. P., Barouch, F. C., & Marx, J. L. (2011). Randomized trial of anesthetic methods for intravitreal injections. *Retina*, 31, 535-539.
- Bonini-Filho, M. A., Jorge, R., Barbosa, J. C., Calucci, D., Cardillo, J. A., & Costa, R. A. (2005). Intravitreal injection versus sub-Tenon's infusion of triamcinolone acetate for refractory diabetic macular edema: a randomized clinical trial. *Investigative Ophthalmology and Visual Science*, 46, 3845-3849.
- Bonini-Filho M, Costa R. A., Calucci D, Jorge R, Melo L. A. Jr, & Scott I. U. (2009). Intravitreal bevacizumab for diabetic macular edema associated with severe capillary loss: one-year results of a pilot study. *Am Journal of Ophthalmology*, 147, 1022-1030.
- Branch Vein Occlusion Study Group (1986). Argon laser scatter photocoagulation for prevention of neovascularization and vitreous hemorrhage in branch vein occlusion: A randomized clinical trial. *Archives of Ophthalmology*, 104, 34-41.
- Bustros, S., Thompson, J. T., Michels, R. G., & Rice, T. A. (1987). Vitrectomy for progressive proliferative diabetic retinopathy. *Archives of Ophthalmology*, 105, 196-199.
- Central Vein Occlusion Study Group (1993). Central vein occlusion study of photocoagulation therapy: Baseline findings. *Online Journal of Current Clinical Trials*, 95, 6021.
- Chen, J. C. (1996). Sutureless pars plana vitrectomy through self-sealing sclerotomy. *Archives of Ophthalmology*, 114, 1273-1275.
- Cintra, L. P., Lucena, L. R., Da Silva, J. A., Costa, R. A., Scott, I. U., & Jorge, R. (2009). Comparative study of analgesic effectiveness using three different anesthetic techniques for intravitreal injection of bevacizumab. *Ophthalmic Surgery, Lasers and Imaging*, 40(1), 13-18.
- Cook, H. L., Newsom, R. S., Mensah, E., Saeed, M., James, D., & Ffytche, T. J. (2002). Entonox as an analgesic agent during panretinal photocoagulation. *British Journal of Ophthalmology*, 86, 1107-1108.
- Costa, R. A., Jorge, R., Calucci, D., Melo, L. A., Jr., Cardillo, J. A., & Scott, I. U. (2007). Intravitreal bevacizumab (avastin) for central

- and hemicentral retinal vein occlusions: IBeVO study. *Retina*, 27, 141-149.
- Davis, D. B., 2nd, & Mandel, M. R. (1986). Posterior peribulbar anesthesia, an alternative to retrobulbar anesthesia. *Journal of Cataract and Refractive Surgery*, 12, 182-184.
- de Faria Rodrigues, A., Messias, A. M., Da Silva, J. A., Ribeiro, J. A., Jorge, R., & Scott, I. U. (2010). Diclofenac for panretinal photocoagulation pain. *Ophthalmology*, 117, 2441.e1-3.
- Demediuk, O. M., Dhaliwal, R. S., Papworth, D. P., Devenyi, R. G., & Wong, D. T. (1995). A comparison of peribulbar and retrobulbar anesthesia for vitreoretinal surgical procedures. *Archives of Ophthalmology*, 113, 908-913.
- Diabetic Retinopathy Study Research Group (1978). Photocoagulation treatment of proliferative diabetic retinopathy: The second report of diabetic retinopathy study (DRS) findings. *Ophthalmology*, 85, 82-106.
- Duker, J. S., Belmont, J. B., Benson, W. E., Brooks, H. L., Jr., Brown, G. C., Federman, J. L., Fischer, D. H., & Tasman, W. S. (1991). Inadvertent globe perforation during retrobulbar and peribulbar anesthesia: Patient characteristics, surgical management, and visual outcome. *Ophthalmology*, 98, 519-526.
- Early Treatment Diabetic Retinopathy Study Research Group (1985). Photocoagulation for diabetic macular edema: Early Treatment Diabetic Retinopathy Study Report No. 1. *Archives of Ophthalmology*, 103, 1796-1806.
- Early Treatment Diabetic Retinopathy Study Research Group (1991). Early photocoagulation for diabetic retinopathy: ETDRS Report Number 9. *Ophthalmology*, 98(5 Suppl.), 766-785.
- Friberg, T. R. & Venkatesh, S. (1995). Alteration of pulse configuration affects the pain response during diode laser photocoagulation. *Lasers in Surgery and Medicine*, 1995, 16, 380-383.
- Friedman, S. M., & Margo, C. E. (2006). Topical gel vs subconjunctival lidocaine for intravitreal injection: A randomized clinical trial. *American Journal of Ophthalmology*, 142, 887-888.
- Fung, A. E., Rosenfeld, P. J., & Reichel, E. (2006). The International Intravitreal Bevacizumab Safety Survey: Using the internet to assess drug safety worldwide. *British Journal of Ophthalmology*, 90, 1344-1349.
- Gragoudas, E. S., Adamis, A. P., Cunningham, E. T., Jr., Feinsod, M., & Guyer, D. R. (2004). Pegaptanib for neovascular age-related macular degeneration. *New England Journal of Medicine*, 351, 2805-2816.
- Hillier, R. J., Aboud, A., Thind, G., & Clark, D. I. (2009). Oral transmucosal fentanyl citrate: A novel analgesic agent for use in retinal photocoagulation. *Retina*, 29, 1506-1512.
- Hitchings, R. A. (1980). The symptom of ocular pain. *Transactions of the Ophthalmological Society of the United Kingdom*, 100, 257-259.
- Jackson, T. (2000). Modified sutureless sclerotomies in pars plana vitrectomy. *American Journal of Ophthalmology*, 129, 116-117.
- Jager, R. D., Aiello, L. P., Patel, S. C., & Cunningham, E. T., Jr. (2004). Risks of intravitreal injection: A comprehensive review. *Retina*, 24, 676-698.
- Jaissle, G. B., Szurman, P., & Bartz-Schmidt, K. U. (2005). Recommendation for the implementation of intravitreal injections: Statement of the German Retina Society, the German Society of Ophthalmology (DOG) and the German Professional Association of Ophthalmologists (BVA). *Klinische Monatsblätter für Augenheilkunde*, 222, 390-395.
- Javitt, J. C., Addiego, R., Friedberg, H. L., Libonati, M. M., & Leahy, J. J. (1987). Brain stem anaesthesia after retrobulbar block. *Ophthalmology*, 94, 718-723.
- Kaderli, B., & Avci, R. (2006). Comparison of topical and subconjunctival anesthesia in intravitreal injection administrations. *European Journal of Ophthalmology*, 16, 718-721.
- Kallio, H., Puska, P., Summanen, P., Paloheimo, M., & Maunukela, E. L. (1999). Retrobulbar/peribulbar block with 0.2% ropivacaine or 1% lidocaine for transscleral cyclophotocoagulation or retinal panphotocoagulation. *Regional Anesthesia and Pain Medicine*, 24, 341-346.
- Kirkby, G. R., Benson, M. T., Callear, A. B., & Loo, A. (1999). Local anaesthesia for vitreoretinal surgery: A case control study of 200 cases. *Eye*, 13, 122-123.
- Kishore, K., Conway, M. D., & Peyman, G. A. (2001). Intravitreal clindamycin and dexamethasone for toxoplasmic retinochoroiditis. *Ophthalmic Surgery and Lasers*, 32, 183-192.
- Kozak, I., Cheng, L., & Freeman, W. R. (2005). Lidocaine gel anesthesia for intravitreal drug administration. *Retina*, 25, 994-998.
- Kristin, N., Schönfeld, C. L., Bechmann, M., Bengisu, M., Ludwig, K., Scheider, A., & Kampik, A. (2001). Vitreoretinal surgery: Preemptive analgesia. *British Journal of Ophthalmology*, 85, 1328-1331.
- Kwok, A. K., Tham, C. C., Lam, D. S., Li, M., & Chen, J. C. (1999a). Modified sutureless sclerotomies in pars plana vitrectomy. *American Journal of Ophthalmology*, 127, 731-733.
- Kwok, A. K., Van Newkirk, M. R., Lam, D. S., & Fan, D. S. (1999b). Subtenons anesthesia in vitreoretinal surgery: A needleless technique. *Retina*, 19, 291-296.
- Lai, M. M., Lai, J. C., Lee, W. H., Huang, J. J., Patel, S., Ying, H. S., ... Handa, J. (2005). Comparison of retrobulbar and sub-Tenon's capsule injection of local anesthetic in vitreoretinal surgery. *Ophthalmology*, 112, 574-579.
- Lira, R. P., Nascimento, M. A., Arieta, C. E., de Carvalho, K. M., & Silva, V. B. (2010). Pain perception at laser treatment of peripheral retinal degenerations with green and infrared wavelengths. *American Journal of Ophthalmology*, 150, 726-730.
- Lucena, C. R. F. A., Ramos-Filho, J. A., Messias, A. M. V., Da Silva, J. A., Almeida, F. P. P. A., Scott I. U., ... Jorge R. (2013). Panretinal photocoagulation versus intravitreal injection retreatment pain in high-risk proliferative diabetic retinopathy. *Arquivos Brasileiros de Oftalmologia* (in press)
- Newsom, R. S. B., Wainwright, A. C., & Canning, C. R. (2001). Local anaesthesia for 1221 vitreoretinal procedures. *British Journal of Ophthalmology*, 85, 225-227.
- Patel, J. I., Jenkins, L., Benjamin, L., & Webber, S. (2006). Dilated pupils and loss of accommodation following diode panretinal photocoagulation with sub-tenon local anaesthetic in four cases. *Eye*, 16, 628-632.
- Pollak, A., & Oliver, M. (1981). Argon laser photocoagulation of symptomatic flap tears and retinal breaks of fellow eyes. *British Journal of Ophthalmology*, 65, 469-472.
- Ramsay, R. C., Knobloch, W. H., & Cantrill, H. L. (1986). Timing of vitrectomy for active proliferative diabetic retinopathy. *Ophthalmology*, 93, 283-289.
- Rao, G. P., Wong, D., Groenewald, C., McGalliard, J. N., Jones, A., & Ridges, P. J. (1998). Local anaesthesia for vitreoretinal surgery: a case control study of 200 cases. *Eye*, 12, 407-411.
- Roman, S. J., Chong Sit, D. A., Boureau, C. M., Auclin, F. X., & Ullern, M. M. (1997). Sub-Tenon's anaesthesia: An efficient and safe technique. *British Journal of Ophthalmology*, 81, 673-676.
- Rosenfeld, P. J., Moshfeghi, A. A., & Puliafito, C. A. (2005). Optical coherence tomography findings after an intravitreal injection of bevacizumab (Avastin) for neovascular age-related macular degeneration. *Ophthalmic Surgery, Lasers and Imaging*, 36, 331-335.
- Schmidt, J., Nietgen, G. W., & Brieden, S. (1999). Self-sealing, sutureless sclerotomy in pars plana vitrectomy. *Klinische Monatsblätter für Augenheilkunde*, 215, 247-251.
- Schlote, T., & Derse, M. (2001). Subconjunctival anesthesia in contact diode laser cyclophotocoagulation. *Ophthalmic Surgery and Lasers*, 32, 289-293.
- Schönfeld, C. L., Hierneis, S., & Kampik, A. (2012). Preemptive analgesia with ropivacaine for pars plana vitrectomy: Randomized controlled trial on efficacy and required dose. *Retina*, 32, 912-917.
- Scott, I. U., Cruz-Villegas, V., Flynn, H. W., Jr., & Miller, D. (2004). Delayed-onset, bleb-associated endophthalmitis caused by *Lecytophora mutabilis*. *American Journal of Ophthalmology*, 137, 583-585.
- Stevens, J. D., Foss, A. J., & Hamilton, A. M. (1993). No-needle one-quadrant sub-tenon anaesthesia for panretinal photocoagulation. *Eye*, 7, 768-771.
- Stevens, J. D., Franks, W. A., Orr, G., Leaver, P. K., & Cooling, R. J. (1992). Four-quadrant local anaesthesia technique for vitreoretinal surgery. *Eye*, 6, 583-586.
- Tang, S., Lai, P., Lai, M., Zou, Y., Li, J., & Li, S. (2007). Topical anesthesia in transconjunctival sutureless 25-gauge vitrectomy for macular-based disorders. *Ophthalmologica*, 221, 65-68.

- Taskintuna, I., Rahhal, F. M., Arevalo, J. F., Munguia, D., Banker, A. S., De Clercq, E., & Freeman, W. R. (1997). Low-dose intravitreal cidofovir (HPMPC) therapy of cytomegalovirus retinitis in patients with acquired immune deficiency syndrome. *Ophthalmology*, *104*, 1049-1057.
- Tesha, P. E., Giavedoni, L. R., Berger, A. R., Altomare, F., Chow, D. R., Navajas, E. V., Yoganathan, P., Wong, D. T., & Principe, A. (2010). Subconjunctival lidocaine before laser treatment: A randomized trial. *Ophthalmology*, *117*, 1810-1814.
- Theelen, T., Verbeek, A. M., Tilanus, M. A., & van den Biesen, P. R. (2003). A novel technique for self-sealing, wedge-shaped pars plana sclerotomies and its features in ultrasound biomicroscopy and clinical outcome. *American Journal of Ophthalmology*, *136*, 1085-1092.
- Theocharis, I. P., Alexandridou, A., & Tomic, Z. (2007). A two-year prospective study comparing lidocaine 2% jelly versus peribulbar anaesthesia for 25G and 23G sutureless vitrectomy. *Graefe's Archive for Clinical and Experimental Ophthalmology*, *245*, 1253-1258.
- Tamai, M., & Mizuno, K. (1984). Distribution of intra- and extraocular pain induced by argon laser photocoagulation. *Tohoku Journal of Experimental Medicine*, *142*, 427-435.
- Tonello, M., Costa, R. A., Almeida, F. P., Barbosa, J. C., Scott, I. U., & Jorge, R. (2008). Panretinal photocoagulation versus PRP plus intravitreal bevacizumab for high-risk proliferative diabetic retinopathy (IBeHi study). *Acta Ophthalmologica*, *86*, 385-389.
- Vaideanu, D., Taylor, P., McAndrew, P., Hildreth, A., Deady, J. P., & Steel, D. H. (2006). Double masked randomised controlled trial to assess the effectiveness of paracetamol in reducing pain in panretinal photocoagulation. *British Journal of Ophthalmology*, *90*, 713-717.
- Weinberger, D., Ron, Y., Lichter, H., Rosenblat, I., Axer-Siegel, R., & Yassur, Y. (2000). Analgesic effect of topical sodium diclofenac 0.1% drops during retinal laser photocoagulation. *British Journal of Ophthalmology*, *84*, 135-137.
- Wu, W. C., Hsu, K. H., Chen, T. L., Hwang, Y. S., Lin, K. K., Li, L.M., Shih, C.P., & Lai, C.C. (2006). Interventions for relieving pain associated with panretinal photocoagulation: A prospective randomized trial. *Eye*, *20*, 712-719.
- Yau, G. L., Jackman, C. S., Hooper, P. L., & Sheidow, T. G. (2011). Intravitreal injection anesthesia—comparison of different topical agents: A prospective randomized controlled trial. *American Journal of Ophthalmology*, *151*, 333-337.