



Revista Argentina de Microbiología

ISSN: 0325-7541

ram@aam.org.ar

Asociación Argentina de Microbiología
Argentina

Gornatti Churria, Carlos D.; Vigo, Germán B.; Origlia, Javier; Campos, Josefina; Caffer, María;
Píscopo, Miguel; Herrero Loyola, Miguel; Petruccelli, Miguel; Pichel, Mariana
Diagnosis of an outbreak of Salmonella Typhimurium in chinchillas (*Chinchilla lanigera*) by pulsed-field
gel electrophoresis

Revista Argentina de Microbiología, vol. 46, núm. 3, 2014, pp. 205-209

Asociación Argentina de Microbiología

Buenos Aires, Argentina

Available in: <http://www.redalyc.org/articulo.oa?id=213032482007>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



REVISTA ARGENTINA DE MICROBIOLOGÍA

www.elsevier.es/ram



INFORME BREVE

Diagnosis of an outbreak of *Salmonella* Typhimurium in chinchillas (*Chinchilla lanigera*) by pulsed-field gel electrophoresis[☆]

Carlos D. Gornatti Churria^{a,*}, Germán B. Vigo^b, Javier Origlia^a, Josefina Campos^c,
María Caffer^c, Miguel Píscopo^a, Miguel Herrero Loyola^a, Miguel Petruccelli^a,
Mariana Pichel^c

^a Cátedra de Patología de Aves y Pílijeros y Laboratorio de Diagnóstico de Enfermedades de las Aves y los Pílijeros, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina

^b Cátedra de Microbiología y Laboratorio de Diagnóstico e Investigaciones Bacteriológicas, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina

^c Servicio de Enterobacterias, Departamento de Bacteriología, Instituto Nacional de Enfermedades Infecciosas, ANLIS "Dr. Carlos G. Malbrán", Buenos Aires, Argentina

Received 17 February 2014; accepted 2 June 2014

KEYWORDS

Chinchilla lanigera;
Farmed chinchillas;
PFGE;
Salmonella
Typhimurium;
Zoonosis

Abstract

Adult chinchillas (*Chinchilla lanigera*) that had suddenly died in a commercial farm located in La Plata City, Buenos Aires Province, Argentina, in July 2012 were macroscopically, histopathologically, and microbiologically examined. *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) was isolated from the liver, spleen, heart, lungs, kidneys and intestines from each of the five animals evaluated. The five strains were susceptible to ampicillin, cephalotin, cefotaxime, nalidixic acid, gentamicin, streptomycin, chloramphenicol, fosfomicin, nitrofurantoin and trimethoprim-sulfamethoxazole, and resistant to tetracycline. Each of the five *S. Typhimurium* isolates was analyzed by *Xba*I- pulsed-field gel electrophoresis (PFGE), showing an identical electrophoretic profile with 15 defined bands, which was found to be identical to pattern ARJPXX01.0220 of the PulseNet Argentine National database of *Salmonella* PFGE patterns. This is the first work describing the postmortem diagnosis of an outbreak of salmonellosis in chinchillas by using molecular methods such as PFGE.

© 2014 Asociación Argentina de Microbiología. Published by Elsevier España, S.L. All rights reserved.

[☆] We would like to dedicate this article to Germán B. Vigo who passed away during its writing.

* Corresponding author.

E-mail address: danielgornatti@fcv.unlp.edu.ar (C.D. Gornatti Churria).

PALABRAS CLAVE

Chinchilla lanigera;
Chinchillas de granja;
PFGE;
Salmonella
Typhimurium;
Zoonosis

Diagnóstico de un brote de *Salmonella* Typhimurium en chinchillas (*Chinchilla lanigera*) mediante la electroforesis en gel de campo pulsado

Resumen

Empleando estudios anatomopatológicos y microbiológicos se examinó a un grupo de chinchillas (*Chinchilla lanigera*) adultas que murieron súbitamente en 2012 en una granja de la ciudad de La Plata (Buenos Aires, Argentina). Se aisló *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) del hígado, el bazo, el corazón, los pulmones, los riñones y los intestinos de los cinco animales evaluados. Los cinco aislamientos estudiados (uno por animal) fueron sensibles a ampicilina, cefalotina, cefotaxima, ácido nalidixico, gentamicina, estreptomycin, cloranfenicol, fosfomicina, nitrofurantoina y trimetoprima-sulfametoxazol, y resistentes a tetraciclina. El análisis de dichos aislamientos por electroforesis en gel de campo pulsado [*pulsed-field gel electrophoresis* (PFGE)] con *Xba*I mostró un perfil electroforético idéntico con 15 bandas, idéntico a su vez al patrón ARJPXX01.0220 del banco nacional argentino de datos de PulseNet, que cuenta con patrones de PFGE de *Salmonella*. El presente trabajo describe por primera vez el diagnóstico postmortem de un brote de salmonelosis en chinchillas usando un método molecular, como la electroforesis en gel en campo pulsado.

© 2014 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Salmonellosis is a widespread zoonotic and foodborne disease that represents one of the most important global veterinary and public health threats. *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) represents one of the main causes of salmonellosis in both humans and animals worldwide¹¹. Chinchillas (*Chinchilla lanigera*) are hystricomorphic mammals which belong to the order Rodentia and family Chinchillidae and are native of the South American Andes Mountains, having natural and restrictive distribution in Peru, Bolivia, and Chile⁶. The species *C. lanigera* was domesticated and now there are millions of chinchillas in captivity throughout the world. Chinchillas are raised and exhibited in zoos, used as exotic pets, and bred in farms for the commercial use of their pelts^{6,10}.

Salmonella spp. isolates can be characterized by phenotyping methods and by genotyping methods such as pulsed-field gel electrophoresis^{5,11} (PFGE). The PFGE method uses rare cutting restriction enzymes that generate a large number of DNA fragments, whose patterns are highly specific for molecular subtyping when separated by agarose gel electrophoresis^{5,13}. Thus, PFGE, which has not been previously applied to confirm the diagnosis of salmonellosis in chinchillas or to clarify epidemiologic and phylogenetic aspects of *Salmonella* strains associated with outbreaks or individual cases in this animal species, represents one of the best genotyping and subtyping methods for members of the genus *Salmonella*^{5,11,13}.

The objective of our study was to characterize the *S. Typhimurium* strains associated to an outbreak that occurred on a chinchilla farm.

In the last decades, *S. Typhimurium* and *Salmonella enterica* serovar Enteritidis (*S. Enteritidis*) have emerged as the main causes of human foodborne illnesses worldwide and have been recognized as the most prevalent causes of *Salmonella* infections in humans in the USA and Europe^{1,4}. According to the information reported by the Argentine

National Reference Laboratory and the WHO Global Foodborne Infections Network, the same situation was observed in Argentina, where *S. Typhimurium* has been the most common serovar isolated from human infections since 2007. Moreover, Argentina is known as one of the leading producers of chinchilla pelts worldwide and salmonellosis is one of the most important infectious diseases affecting chinchilla production².

Salmonellosis has been described in chinchillas since 1966¹⁴. The first report of salmonellosis in chinchillas mentioned an outbreak of *Salmonella enterica* serovar Dublin (*S. Dublin*) on a chinchilla farm in the United Kingdom¹⁴. This British study described that the affected chinchillas showed loss of body condition, enteritis, diarrhea, and sudden death. *S. Dublin* was isolated from three out of the nine chinchillas evaluated. Due to the fact that the strains showed antibiotic susceptibility to chloramphenicol, furazolidone, neomycin, ampicillin, and framycetin, antibiotic treatment with chloramphenicol was implemented and the outbreak was successfully controlled. The authors of this study concluded that the *S. Dublin* outbreak in British farmed chinchillas could have been associated with recently introduced carrier breeders and highlighted the possible infective role of one of the farmers because of its *S. Dublin* confirmed carrier status¹⁴. Another British study carried out in 1989 briefly summarized the isolation of *Salmonella enterica* subsp. *arizonae* recovered from a pet chinchilla with septicemia and identified as the rare serotype O-21 r,z⁸. In Slovakia, a surveillance of bacterial diseases in several commercial large-scale chinchilla farms was reported in 1994, and both *S. Enteritidis* and *Listeria* spp. infections were found to be the main causes of death in farmed chinchillas¹⁰. In Japan, a *S. Enteritidis* strain was isolated from most of the organs and intestinal content of a pet chinchilla affected by septicemia in 1997¹⁵. The postmortem and microscopic evaluations of this pet chinchilla revealed meningitis, panophthalmitis, necrotic enteritis, focal hepatic ne-

crotic lesions and microvascular thrombosis in the spleen and kidneys, which were associated with the bacterial infection since *S. Enteritidis* colonies were immunohistochemically detected in the affected pet chinchilla¹⁵. In Argentina, outbreaks of *S. Typhimurium* were reported on two chinchilla farms located in Jujuy Province and Santa Fe Province showing 12-30% of mortality in 1999². This Argentine study described farmed chinchillas suffering from anorexia, apathy, hyperthermia, diarrhea, abortion, and death. The postmortem examination of the affected animals revealed septicemic lesions in the liver, lungs, kidneys, stomach, intestines and uterus. The microbiological study allowed to isolate *S. Typhimurium* strains which were susceptible to a wide range of antibiotics in both cases. This study highlighted the control of both outbreaks with antibiotics such as furazolidone, chloramphenicol, and enrofloxacin and autogenous bacterins². In Turkey, an outbreak caused by both *S. Typhimurium* and *S. Enteritidis* infections on a chinchilla farm was reported in 2002⁷. This study described farmed chinchillas suffering from apathy, anorexia, tremors, paralysis, diarrhea, vaginal secretions, abortion, and death. The postmortem evaluation of two affected chinchillas revealed enteritis, hemorrhagic and purulent endometritis and septicemic lesions. This Turkish study reported the isolation of *Salmonella* strains which were susceptible to enrofloxacin, amoxicillin, and gentamicin, and resistant to erythromycin and ampicillin⁷. In Croatia, an outbreak of *S. Enteritidis* and the isolation of *S. Sofia* in farmed chinchillas was reported in 2003⁹. The postmortem examinations made in this Croatian study revealed gastritis, enteritis, enlarged spleens with or without focal necrosis, livers with focal necrosis and endometritis in chinchillas from five different commercial farms. The authors of this study reported the isolation of *S. Enteritidis* from the five commercial farms evaluated, and *S. Sofia* from one of them. All *S. Enteritidis* isolates were susceptible to ampicillin, chloramphenicol, neomycin, enrofloxacin, and trimethoprim-sulfamethoxazole, and resistant to sulfamethoxazole, whereas only three *S. Enteritidis* isolates were resistant to tetracycline. This study highlighted the successful results of the antibiotic treatments with enrofloxacin and discussed possible routes of *Salmonella* infection for both humans and animals, considering that most of the animal-derived food groups and human-animal working contacts could represent serious threats for public and veterinary health⁹. In Brazil, a retrospective study of postmortem examinations of chinchillas (1997-2011) including 202 necropsied animals was reported in 2012. This study briefly described one animal affected by a *Salmonella* spp. infection associated with hepatitis and cholecystitis⁶.

Eight previous works have described salmonellosis in both pet and farmed chinchillas^{2,6-10,14,15}, but none of them reported the application of molecular methods such as PFGE. Reports of outbreaks in farmed chinchillas or individual infections in pet chinchillas due to *S. Typhimurium*^{2,7}, *S. Enteritidis*^{7,9,10,15}, *S. Dublin*¹⁴, *S. Sofia*⁹, *S. enterica* subsp. *Arizonae*⁸, and *Salmonella* spp.⁶ allow to consider the high relevance of salmonellosis in chinchillas as a veterinary and public health threat worldwide, and the need for molecular methods such as PFGE to support its diagnosis and further investigations.

In our case, adult chinchillas suffered sudden death on a commercial farm located in La Plata City, Buenos Aires Province, Argentina, in July 2012. The outbreak lasted one week, during which 35% of the 3600 farmed chinchillas died. Each animal was housed in a stainless steel cage of standard size. The farmers provided running water using a hose system with one nipple per cage, and commercial chinchilla pelleted food which was stored in a peridomestic rodent-free covered container. Poor hygienic conditions were noted because abundant peridomestic rodent feces were observed close to several cages.

Five of the 1260 dead chinchillas were submitted to the Laboratorio de Diagnóstico de Enfermedades de las Aves y los Pilíferos (Cátedra de Patología de Aves y Pilíferos, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata) for their postmortem examination and diagnostic workup.

The postmortem examination revealed emaciated animals with hemorrhagic inflammation and presence of mucous and white nodules of approximately 2-3 mm in diameter in the duodenum and stomach. The livers showed pale, dark, and gray color changes. Splenomegaly and hemorrhagic pneumonia were also observed in all the animals examined. Samples of the intestine, stomach, lungs, liver, kidneys, heart, and spleen from the five animals were taken at necropsy, fixed in 10% buffered formalin for 48 h, embedded in paraffin, sectioned at 3 µm, and stained with hematoxylin and eosin (H&E) and periodic acid-Schiff (PAS). Smears of tissues were stained using Gram and modified Ziehl-Neelsen (Z-N) methods.

Tissue samples were also inoculated onto Hektoen enteric agar plates (Becton Dickinson France S.A., Le Pont-de-Claix, France), MacConkey agar plates (Becton Dickinson France S.A.), and trypticase soy agar plates (Becton Dickinson France S.A.) with 5% sheep blood and then incubated at 37°C for 48 h under aerobic conditions. In addition, these samples were inoculated onto Sabouraud agar plates (Merck KGaA, Darmstadt, Germany) and incubated at 28°C for 1 week under aerobic conditions. Presumptive *Salmonella* spp. colonies were chosen and biochemically tested. Species, subspecies and serotype were identified according to the White-Kauffmann-LeMinor scheme.

Susceptibility tests of *Salmonella* spp. isolates to ampicillin (10 µg/disk), cefotaxime (30 µg/disk), nalidixic acid (30 µg/disk), gentamicin (10 µg/disk), streptomycin (10 µg/disk), chloramphenicol (30 µg/disk), cephalotin (30 µg/disk), nitrofurantoin (300 µg/disk) and trimethoprim-sulfamethoxazole (1.25/23.75 µg/disk) were performed following standardized disk diffusion methods for *Enterobacteriaceae*³. The relatedness among the isolates obtained from each of the five farmed chinchillas was determined by PFGE by applying the *Xba*I enzyme (New England Biolabs, Inc., Ipswich, MA), following the PulseNet standard protocol¹² in a CHEF-DR III Pulsed Field Electrophoresis System (Bio-Rad Laboratories, Inc., Hercules, CA, USA). Gel images were obtained with a Gel-Doc 2000 Gel Documentation System (Bio-Rad Laboratories, Inc., Hercules, CA, USA). PFGE patterns were analyzed using BioNumerics software version 4.5 (Applied Maths, Kortrijk, Belgium). The relationship among the PFGE patterns was estimated using the same software, applying the Dice coefficient and

the UPGMA method with 1.5% tolerance and 1.5% optimization parameters as recommended by PulseNet usual analysis. PFGE profiles of the *S. Typhimurium* isolates obtained from each of the five chinchillas in the present case were compared with the PulseNet Argentine National database of *Salmonella* PFGE patterns, which contains 364 unique *Xba*I-PFGE patterns corresponding to 1291 isolates of *S. Typhimurium* from different sources (human, animals, and the environment), obtained in Argentina from 1968 to 2013.

The H&E microscopic examination of lung tissues showed edema and severe infiltration of mononuclear inflammatory cells (macrophages, lymphocytes, and plasma cells) with diffuse distribution in alveoli and blood vessel congestion. The livers showed sinusoidal and great blood vessel congestion, small foci of necrosis, and small focal areas of mononuclear inflammatory cell infiltration with diffuse distribution. The spleens showed moderate lymphoid depletion.

The stomachs had evident edema in the lamina propria, and severe mononuclear inflammatory cell infiltration in the mucous membrane, with epithelium hyperplasia. The intestines showed edema in the lamina propria and mononuclear inflammatory cell infiltration in the mucous membrane, with abundant bacterial colonies. The PAS and modified Z-N stains yielded negative results and no fungal growth was observed in Sabouraud agar plates.

The isolates obtained from each of the five farmed chinchillas were identified as *S. Typhimurium*. The isolates were susceptible to all the antimicrobial agents tested except to tetracycline. The five *S. Typhimurium* isolates were identical by *Xba*I-PFGE, showing an electrophoretic profile with 15 defined bands between 670 and 20 kilobases. This profile was found to be identical to pattern ARJPXX01.0220, representing 1.5% of the PulseNet Argentine National database of *Salmonella* PFGE patterns, which includes isolates from hu-




Figure 1 Selected *Xba*I enzyme-PFGE patterns from the PulseNet Argentine National database of *Salmonella* PFGE patterns including the chinchilla isolates.

man, animal, and food sources. The *Xba*I-PFGE pattern was also related to a foodborne outbreak in a hospital in Buenos Aires Province in 2013 (Fig. 1).

In our case, the isolation of *S. Typhimurium* from the liver, spleen, heart, lungs, kidneys and intestines from each of the five adult farmed chinchillas indicated septicemic infections. The macroscopic and microscopic lesions observed in our study were associated with acute salmonellosis, as previously reported in this animal species^{2,7,9,15}.

All isolates obtained in our study were resistant to tetracycline, which could be the result of overusing this antimicrobial agent against other bacterial infections. The *S. Typhimurium* strains isolated in our study showed low levels of antibiotic resistance, in accordance with the low levels of antibiotic resistance previously observed in Argentine and Turkish *S. Typhimurium* strains isolated from farmed chinchillas in 1999 and 2002, respectively^{2,7}, and in contrast to the high levels of antibiotic resistance observed in *S. Typhimurium* strains isolated from humans, other animal species and foodstuffs previously reported worldwide¹.

This is the first report confirming an outbreak of salmonellosis in chinchillas by molecular methods such as PFGE. The same clone of *S. Typhimurium* was isolated in the five farmed chinchillas evaluated in our study. Therefore, our results support the hypothesis that the animals acquired the infection from the same infectious source. Unfortunately, such source could not be established until now, but the presence of abundant feces of peridomestic rodents inside the farm could have played a key role in the transmission of the bacterial pathogen during this outbreak. Other possible routes of transmission could have been associated with the consumption of contaminated food or water, contact with human or animal carriers or combinations of these factors. The *Xba*I-PFGE pattern of the *S. Typhimurium* strains isolated from each of the five farmed chinchillas evaluated in our case has been identified since 2006 in samples from humans, domestic animals, and food from different regions of Argentina. This means that this clone is widely disseminated in Argentina. The PFGE pattern identified in the farmed chinchillas involved in this outbreak was immediately incorporated into the PulseNet Argentine National database of *Salmonella* PFGE patterns.

Due to the fact that people handling chinchillas or cleaning chinchilla accessories (e.g. cages) are likely to acquire *Salmonella* spp. infections, they should follow the recommended precautionary measures to reduce the transmission risk of this zoonotic microorganism.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Authorship

C. D. Gornatti Churria and G. B. Vigo contributed equally to this article and are joint first authors.

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

1. Busani L, Graziani C, Battisti A, Franco A, Ricci A, Vio D, Digiannatale E, Patterlini F, D'Incau M, Owczarek S, Caprioli A, Luzzi I. Antibiotic resistance in *Salmonella enterica* serotypes Typhimurium, Enteritidis and Infantis from human infections, foodstuffs and farm animals in Italy. *Epidemiol Infect.* 2004;132:245-51.
2. Caffer M, Abdala A, Romano F, Gianre V R, Alcain A, Terragno R. Salmonellosis en chinchillas. *Vet Argent.* 1999;16:423-6.
3. Clinical and Laboratory Standards Institute. Disk diffusion. Performance Standards for Antimicrobial Susceptibility Testing; 19th Informational Supplement, 2009; M100-S. Wayne, PA, USA.
4. European Food Safety Authority, European Centre for Disease Prevention and Control. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2011. *EFSA J.* 2013;11:3129.
5. Guerra B, Schrörs P, Mendoza MC. Application of PFGE performed with *Xba*I to an epidemiological and phylogenetic study of *Salmonella* serotype Typhimurium. Relations between genetic types and phage types. *New Microbiol.* 2000;23:11-20.
6. Lucena RB, Giarretta PR, Tessele B, Figuera R, Kommers G, Irigoyen L, Barros CSL. Diseases of chinchillas (*Chinchilla lanigera*). *Pesq Vet Bras.* 2012;32:529-35.
7. Misirlioğlu D, Çetin C, Kahraman MM, Caner V, Özyiğit M. *Salmonella* infection in a chinchilla farm. *Turk J Vet Anim Sci.* 2002;26:151-5.
8. Mountain A. *Salmonella* Arizona in a chinchilla. *Vet Rec.* 1989;125:25.
9. Naglič T, Šeol B, Bedeković M, Grabarević Ž, Listeš E. Outbreak of *Salmonella* Enteritidis and isolation of *Salmonella* Sofia in chinchillas (*Chinchilla laniger*). *Vet Rec.* 2003;152:719-20.
10. Novak S, Ruttkay D, Solar J. Results of screening of bacterial diseases in large-scale breeding of the chinchilla (*Chinchilla laniger*). *Slovenský Vet Časopis.* 1994;19:19-21.
11. Refsum T, Heir E, Kapperud G, Vardund T, Holstad G. Molecular epidemiology of *Salmonella enterica* serovar Typhimurium isolates determined by pulsed-field gel electrophoresis: comparison of isolates from avian wildlife, domestic animals, and the environment in Norway. *Appl Environ Microbiol.* 2002;68:5600-6.
12. Ribot E, Fair M, Gautom R, Cameron D, Hunter S, Swaminathan B, Barrett T. Standardization of pulsed-field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathog Dis.* 2006;3:59-67.
13. Vigo G, Leotta G, Caffer M, Salve A, Binsztein N, Pichel M. Isolation and characterization of *Salmonella enterica* from Antarctic wildlife. *Polar Biol.* 2011;34:675-81.
14. Watson W, Watson F. An outbreak of *Salmonella* Dublin infection in chinchillas. *Vet Rec.* 1966;78:15-7.
15. Yamagishi S, Watanabe Y, Tomura H, Sekine T, Mimura M, Iijima Y, Fujii M. Septic infection of a companion chinchilla with *Salmonella* Enteritidis. *J Jpn Vet Med Assoc.* 1997;50:345-8.