



Revista Argentina de Microbiología

ISSN: 0325-7541

ram@aam.org.ar

Asociación Argentina de Microbiología
Argentina

Villanueva, María P.; Medina, Gustavo; Fernández, Heriberto
Arcobacter butzleri survives within trophozoite of Acanthamoeba castellanii
Revista Argentina de Microbiología, vol. 48, núm. 2, abril-junio, 2016, pp. 105-109
Asociación Argentina de Microbiología
Buenos Aires, Argentina

Available in: <http://www.redalyc.org/articulo.oa?id=213046439004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



BRIEF REPORTS

Arcobacter butzleri survives within trophozoite of *Acanthamoeba castellanii*



María P. Villanueva^a, Gustavo Medina^b, Heriberto Fernández^{a,*}

^a Instituto de Microbiología Clínica, Universidad Austral de Chile, PO Box 567, Valdivia, Chile

^b Facultad de Ciencias de la Salud, Universidad Católica de Temuco, PO Box 15-D, Temuco, Chile

Received 2 March 2015; accepted 1 December 2015

Available online 11 March 2016

KEYWORDS

Arcobacter butzleri;
Acanthamoeba castellanii;
Survival;
Environmental
reservoir

Abstract The survival of three *Arcobacter butzleri* strains inside *Acanthamoeba castellanii* was assessed using axenic cultures of *A. castellanii* that were inoculated with the tested strains and incubated at 26 °C under aerobic conditions for 240 h. The behavior of bacteria in contact with amoebae was monitored using phase contrast microscopy. The bacterial survival rate within amoebae was assessed through counting colony forming units, using the gentamicin protection assay. All *A. butzleri* strains were able to survive during 240 h within the amoebae, thus suggesting that (i) *A. butzleri* resists the amoebic digestion processes at least for the analyzed time; (ii) that *A. castellanii* could serve as an environmental reservoir for this bacterium, probably acting as a transmission vehicle for *A. butzleri*.

© 2016 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Arcobacter butzleri;
Acanthamoeba castellanii;
Sobrevida;
Reservorio ambiental

Arcobacter butzleri* sobrevive en el interior de trofozoitos de *Acanthamoeba castellanii

Resumen Se determinó la sobrevivencia de 3 cepas de *Arcobacter butzleri* en el interior de trofozoitos de *Acanthamoeba castellanii* utilizando cultivos axénicos de la ameba, los que fueron inoculados con las cepas bacterianas e incubados a 26 °C en aerobiosis durante 240 h. La interacción bacterias/amebas fue monitorizada mediante microscopía de contraste de fase. Las tasas de sobrevivencia bacteriana en el interior de las amebas fueron establecidas a través del recuento de unidades formadoras de colonias mediante ensayos de protección con gentamicina. Todas las cepas de *A. butzleri* fueron capaces de sobrevivir en el interior de la ameba al menos durante 240 h, lo que sugiere, por un lado, que *A. butzleri* puede resistir los procesos de

* Corresponding author.

E-mail address: hfernand@uch.cl (H. Fernández).

digestión propios de la ameba, al menos por el lapso aquí analizado y, por el otro, que *A. castellanii* podría actuar como reservorio ambiental de esta bacteria y, probablemente, como vehículo de transmisión de *A. butzleri*.

© 2016 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la CC BY-NC-ND licencia (<http://creativecommons.org/licencias/by-nc-nd/4.0/>).

The genus *Arcobacter* was originally proposed by Vandamme *et al.*¹⁴ and includes bacteria previously considered as members of the genus *Campylobacter*. Today, the genus *Arcobacter* is represented by 21 species isolated from human, animals and environmental samples¹⁵. However, only four species have been isolated from human and animal samples and they are considered of medical interest: *Arcobacter cryaerophilus*, *Arcobacter butzleri*, *Arcobacter skirrowii* and *Arcobacter thereius*¹³.

A. butzleri is the species that is most frequently isolated from environmental water, food and clinical samples. From a public health point of view, it is considered an emerging foodborne and zoonotic enteropathogen transmitted by food and water. Furthermore, it has been ranked as the fourth most common member of *Campylobacteraceae* isolated from human feces^{2,5,13}. Thus, it has become increasingly important as an agent of diarrhea in human beings, recognizing as potential routes of infection the consumption and manipulation of contaminated raw or poorly cooked food of animal origin and untreated water^{2,5,13}.

One of the main reservoirs of this species is water in diverse environments² where they may interact with other microorganisms, which are natural inhabitants of hydric ecosystems, in particular with free-living amoebas.

Free-living amoebas of the Genus *Acanthamoeba* are widely distributed in water environments. Amoebic trophozoites are part of the indigenous inhabitants of natural water bodies, sharing this habitat with other microorganisms, mainly bacteria. Since these bacteria can survive for diverse periods within the amoeba, many researchers have suggested that these protozoa may be a reservoir for bacterial species¹. Because of the particular ability of some bacteria to survive within amoebae, the members of genus *Acanthamoeba* could play a role in the transmission of some enteric bacteria. The survival of *Campylobacter jejuni* inside *Acanthamoeba* and its transmission to chicken as an endosymbiont of *A. castellanii* and *Acanthamoeba polyphaga*¹ has been demonstrated.

Since *A. butzleri* is a bacterium that can be recovered from water environments and food products, such as mussels produced in estuarine environments^{2,6}, it could share this environment with free-living amoebas, probably establishing close relationships with these protozoa, similarly to those described in the genus *Campylobacter*^{1,11,12}.

In order to demonstrate if *A. butzleri* is capable of establishing similar close relations with free-living amoebas, we determined the survival capacity of *Arcobacter* inside *A. castellanii*.

Three strains of *A. butzleri* were used for this study. Strain F215 (isolated from child human stools), strain

Puar190 (isolated from chicken meat) and the reference strain LMG10828 (isolated from human feces) used as control strain and kindly provided by Dr. Peter Vandamme (Laboratory of Microbiology, Ghent University, Belgium). These strains were stored at -80°C in brain-heart infusion broth supplemented with 15% glycerol. Before each experiment, bacteria were grown on blood agar plates incubated at 26°C for 48 h under aerobic conditions.

A. castellanii strain BP91/2760 was originally isolated from the eye of a patient with keratitis and was kindly provided by Dr. Winiecka-Krusnell (Public Health Agency of Sweden). Amoebae were maintained axenically in peptone yeast extract glucose (PYG) medium at 26°C as monolayers in 25 cm^2 flasks and were examined under an inverted microscope before use. To assess the number of amoebae, counts were made using a Neubauer chamber.

To establish *A. butzleri*/*A. castellanii* early relationships, *A. castellanii* were grown to confluence in 25 cm^2 flasks with 5 ml of PYG medium. The flask were gently shaken and the PYG with non-adherent amoebae was removed and replaced with fresh medium; then the amoebae were incubated on ice for 30 min to weaken adherence. The suspensions were centrifuged at 203 g for 10 min and the pellet was washed three times with Page's saline solution. After the last centrifugation, the pellet was suspended in PYG and the total number of amoebae was estimated. Aliquots of 1 ml with approximately 6×10^4 trophozoites were placed in each well of a 24-well tissue culture dish. Bacterial strains suspended in Page's saline solution to a density of approximately 1.2×10^8 colony forming units (CFU), and 100 ml of this suspension was added to each well containing the amoebae and the plate was incubated at 26°C under aerobic conditions to induce the infection of the amoebas by *Arcobacter* strains in a 1:2000 ratio. As bacterial survival controls, three wells were inoculated with *A. butzleri* suspended in Page's saline solution without amoebae. Finally, the assessment of the early interactions between bacteria and amoebae was performed by taking aliquots of the suspension at 30 and 50 min and observed directly under phase contrast microscope.

Long-term interaction between bacteria and amoebae was studied at 24, 48, 72, 120, 168, and 240 h using a modification of the gentamicin protection assay proposed by Dirks and Quinlan³. In brief, after the early interaction assay, $100\text{ }\mu\text{g/ml}$ gentamicin were added to each well and incubated at room temperature for 1 h to kill extra-amoebic bacterial cells. Amoebas were harvested from the well and washed three times with Page's saline solution. After the third wash, an aliquot of $100\text{ }\mu\text{l}$ was taken and plated on blood agar plates to determine the number of extracellular bacteria that might have escaped the gentamicin treatment




Figure 1 Phase contrast microscopy. (A) Trophozoite of *A. castellanii* showing *A. butzleri* seen in close association with *A. castellanii* gathering at one pole of the amoeba cell known as food-cup (30 min of co-incubation). (B) Trophozoite of *A. castellanii* showing *A. butzleri* within the amoebic vacuoles (50 min of co-incubation).

at each sampling time. 100 μ l of 0.5% sodium desoxycholate was added to another aliquot for 25–30 min to perform the amoeba lysis. An aliquot of 100 μ l of the lysate suspension was taken to count the number of total bacteria by CFU counts (intracellular + extracellular bacteria). Abundance of viable intracellular bacteria was determined by subtracting the number of extracellular bacteria from the total count. All these experiments were carried out in triplicate. The amoebae/bacteria interaction was also monitored and documented through phase contrast microscopy and by bright-field microscopy.

The results of the assays were divided into two groups in accordance with the duration of the tests performed.

A. – Early interactions between *A. butzleri* and *A. castellanii*. In the first 30 min of co-incubation, *A. butzleri* could be seen in close association with *A. castellanii*, at this time most of the bacteria showed a tendency to gather at one pole of the amoeba cell (Fig. 1A). Soon after 50 min of co-culture, *A. butzleri* cells were detected inside the amoeba. The internalized *A. butzleri* were located in vacuoles and never in the amoeba cytoplasm (Fig. 1B). These observations are consistent with our previous studies^{7,8}, confirming an initial contact pattern in the interaction *A. butzleri*/*A. castellanii*.

The internalization of *A. butzleri* seems to be an active and probable metabolic process since all intracellular bacteria are located in vacuoles and never inside the amoeba cytoplasmic region. These images are similar to those previously reported by our group^{7,8}.

This early observation indicated that the interaction between bacteria and amoebae could be a common phenomenon in *Arcobacter*, which is time-efficient. Recently, Medina *et al.*⁸ have reported that the attachment process of *A. butzleri* to *A. castellanii* involves the participation of mannose-binding proteins and membrane-associated receptors of glucose and galactose present in the amoebae, whereas in their internalization, the protozoan actin polymerization plays an active role.

B. – Time course of the interaction of *A. butzleri* and *A. castellanii*. The long-term experiments demonstrated that *A. butzleri* strains F-215 and PUAr190 were able to survive for at least 240 h inside the amoebas. On the other hand, control assays done to establish whether the presence of *A. butzleri* could affect the survival of the amoebae, showed no significant differences in the survival rates of the amoebae with and without bacteria (data not shown).

Regarding the survival of *A. butzleri* within *A. castellanii* trophozoites at different time points, all the bacteria studied in the co-culture with the amoeba as well as control bacteria showed a decrease in CFU values until reaching the minimum count on the tenth day. There is a clear difference in the survival of these three strains of *A. butzleri* inside the amoebae. The data suggest that recently isolated *A. butzleri* strains F-215 and PUAr190 have better survival than the reference strain LMG10828. Their colony counts suggest that this strain does not multiply inside *A. castellanii*, not reaching the high numbers seen for the other two strains (Fig. 2). Furthermore, their UFC numbers drop in a




Figure 2 Average colony-forming units (CFU) of intracellular *A. butzleri* recovered after co-culture with *A. castellanii*. LMG-10828 control strain.

similar way as the number of viable bacterial cells decays in the control assay without amoebae. Whether those differences are related to the source of isolation, geographic origin or number of subcultures is unknown, requiring further investigations.

The methodology used in this study to estimate the intramoeba bacterial survival rate was selected because other methods, such as vital stains and counting by microscopy, express their results as percentages and do not allow to determine the bacterial number as accurately as the gentamicin protection assay does³.

The *A. butzleri* strains used in this study were not only able to infect *A. castellanii* but also capable of surviving inside the vacuoles, in which we confirmed that bacteria were actively motile. Through the phagolysosome formation analysis, Medina *et al.*⁸ concluded that the survival of *A. butzleri* inside the amoebae could be related to their ability to remain inside vacuoles not fused with lysosomes, or with their ability to retard phagolysosome formation. However, further investigations are needed to elucidate the survival mechanisms of *A. butzleri* inside free-living amoebae.

Transmissibility of enteropathogens, particularly those associated in zoonosis, has been difficult to identify and for the same reason, nearly impossible to prevent. In the last quarter of the 20th century and the beginning of this century, emerging pathogens have become a major health problem⁴. Recent observations showed that free-living amoebae may act as a reservoir and vehicle for the survival and transmission of *C. jejuni* and other enteropathogens, providing a unique form of self-perpetuation of these pathogens^{1,9,10,12}.

Our team has been very interested in the role of *Arcobacter* species as human pathogens and its association with animal sources^{5,6}. Different studies have indicated that *A. butzleri* can survive under several environmental conditions and can be isolated from diverse hosts like farm animals, seafood, and house pets^{2,5,6}. In the present study, the survival of *A. butzleri* inside *A. castellanii* strongly suggests that this protozoon can serve as a reservoir for *A. butzleri* and it might be an important vehicle in the transmission of this human emergent zoonotic pathogen, acting as a Trojan horse^{2,8,12}. Experimental transmissibility assays of *A. butzleri* inside *A. castellanii* into mice, currently performed in our laboratory (data not shown), allow to infer the eventual participation of this protozoon as transmission vehicle of *A. butzleri*. Further studies have to be carried out to elucidate this epidemiological angle of the *A. butzleri*/free-living amoebae relationships.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Financial support

Grants DID-UACH S-2007-37 and FONDECYT 1110202.

Conflict of interests

The authors declare that they have no conflict of interests.

Acknowledgements

To Professor Ph.D. Jadwiga Winięcka-Krusnell (Public Health Agency of Sweden, Sweden) for providing the *A. castellanii* T4 genotype strain; Professor Ph.D. Peter Vandamme (Ghent University, Belgium) for providing the *A. butzleri* reference strain LMG 10828; and Professor Ph.D. Guillermo Pérez-Pérez (New York University Langone Medical Center, U.S.A.) for their scientific advice and critical review.

References

1. Axelsson-Olsson D, Waldenström J, Broman T, Olsen B, Holmberg M. Protozoan *Acanthamoeba polyphaga* as a potential reservoir for *Campylobacter jejuni*. *Appl Environ Microbiol*. 2005;71:987–92.
2. Collado L, Figueras MJ. Taxonomy, epidemiology and clinical relevance of the genus *Arcobacter*. *Clin Microbiol Rev*. 2011;24:174–92.
3. Dirks BP, Quinlan JJ. Development of a modified gentamicin protection assay to investigate the interaction between *Campylobacter jejuni* and *Acanthamoeba castellanii* ATCC 30010. *Exp Parasitol*. 2014;140:39–43.
4. Duffy G, Lynch OA, Cagney C. Tracking emerging zoonotic pathogens from farm to fork. *Meat Sci*. 2008;78:34–42.
5. Fernández H, Flores S, Inzunza F. *Arcobacter butzleri* strains isolated from different sources display adhesive capacity to epithelial cells *in vitro*. *Acta Sci Vet*. 2010;38:287–91.
6. Fernández H, Villanueva MP, Mansilla I, Gonzalez M, Latif F. *Arcobacter butzleri* and *A. cryaerophilus* in human, animals and food sources, in southern Chile. *Braz J Microbiol*. 2015;46:147–57.
7. Fernández H, Villanueva MP, Medina G. Endosymbiosis of *Arcobacter butzleri* in *Acanthamoeba castellanii*. *Rev Argent Microbiol*. 2012;44:133.
8. Medina G, Flores-Martin S, Fonseca B, Otth C, Fernandez H. Mechanisms associated with phagocytosis of *Arcobacter butzleri* by *Acanthamoeba castellanii*. *Parasitol Res*. 2014;113:1933–42.
9. Olofsson J, Axelsson-Olsson D, Brudin L, Olsen B, Ellström P. *Campylobacter jejuni* actively invades the amoeba *Acanthamoeba polyphaga* and survives within non-digestive vacuoles. *PLoS ONE*. 2013;8:e78873. <http://dx.doi.org/10.1371/journal.pone.0078873>.
10. Siddiqui R, Khan NA. Biology and pathogenesis of *Acanthamoeba*. *Parasit Vectors*. 2012;5:1–13.
11. Snelling W, Stern N, Lowery C, Moore J, Gibbons E, Baker C, Dooley J. Colonization of broilers by *Campylobacter jejuni* internalized within *Acanthamoeba castellanii*. *Arch Microbiol*. 2008;189:175–9.
12. Vaerewijck MJM, Baré J, Lambrecht E, Sabbe K, Houf K. Interactions of foodborne pathogens with free-living protozoa: potential consequences for food safety. *Compr Rev Food Sci*. 2014;13:924–44.

13. Van den Abeele AM, Vogelaers D, Van Hende J, Houf K. Prevalence of *Arcobacter* species among humans, Belgium, 2008–2013. *Emerg Infect Dis.* 2014;20:1731–4.
14. Vandamme P, Vancanneyt M, Pot B, Mels L, Hoste B, Dewet-
tinck D, Vlaes L, Van den Borre C, Higgins R, Hommez J,
Kerstens K, Butzler JP, Goossens H. Polyphasic taxonomic study
of the emended genus *Arcobacter* with *Arcobacter butzleri*
comb. nov. and *Arcobacter skiwowii* sp. nov., an aerotolerant
bacterium isolated from veterinary specimens. *Int J Syst Bac-
teriol.* 1992;42:344–56.
15. Whiteduck-Léveillé K, Whiteduck-Léveillé J, Clotier M, Tam-
bong JT, Xu R, Topp E, Arts MT, Chao J, Adam Z, Lévesque CA,
Lapen DR, Villemur R, Talbot G, Khan IUH. *Arcobacter lanthieri*
sp. nov. isolated from pig and dairy cattle manure. *Int J Syst
Evol Microbiol.* 2015;65:2709–16.