



Revista Argentina de Microbiología

ISSN: 0325-7541

ram@aam.org.ar

Asociación Argentina de Microbiología
Argentina

Fellner, María Dolores; Durand, Karina A.; Solernou, Veronica; Bosaleh, Andrea;
Balbarrey, Ziomara; García de Dávila, María T.; Rodríguez, Marcelo; Irazu, Lucía; Alonio,
Lidia V.; Picconi, María A.

Epstein-Barr virus load in transplant patients: Early detection of post-transplant
lymphoproliferative disorders

Revista Argentina de Microbiología, vol. 48, núm. 2, abril-junio, 2016, pp. 110-118

Asociación Argentina de Microbiología
Buenos Aires, Argentina

Available in: <http://www.redalyc.org/articulo.oa?id=213046439005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



ORIGINAL ARTICLE

Epstein–Barr virus load in transplant patients: Early detection of post-transplant lymphoproliferative disorders



María Dolores Fellner^{a,*}, Karina A. Durand^a, Veronica Solernou^b,
Andrea Bosaleh^b, Ziomara Balbarrey^b, María T. García de Dávila^b,
Marcelo Rodríguez^c, Lucía Irazu^c, Lidia V. Alonio^a, María A. Picconi^a

^a Oncogenic Viruses Service, Virology Department, "Carlos G. Malbrán" National Institute of Infectious Diseases, Av. Vélez Sársfield 563, C1282AFF Buenos Aires, Argentina

^b Pathology Service, "Prof. Dr. Juan. P. Garrahan" National Pediatrics Hospital, Pichincha 1890, C1249ABP Buenos Aires, Argentina

^c Operational Team Quality Management, Parasitology Department, "Carlos G. Malbrán" National Institute of Infectious Diseases, Av. Vélez Sársfield 563, C1282AFF Buenos Aires, Argentina

Received 21 August 2015; accepted 21 February 2016

Available online 4 May 2016

KEYWORDS

Epstein–Barr virus;
Early PTLD detection;
Lymphoma;
Real-time PCR;
Transplant patients;
Viral load

Abstract High levels of circulating EBV load are used as a marker of post-transplant lymphoproliferative disorders (PTLD). There is no consensus regarding the threshold level indicative of an increase in peripheral EBV DNA. The aim of the study was to clinically validate a developed EBV quantification assay for early PTLD detection.

Transversal study: paired peripheral blood mononuclear cells (PBMC), plasma and oropharyngeal lymphoid tissue (OLT) from children undergoing a solid organ transplant with ($n=58$) and without ($n=47$) PTLD. Retrospective follow-up: 71 paired PBMC and plasma from recipients with ($n=6$) and without ($n=6$) PTLD history. EBV load was determined by real-time PCR. The diagnostic ability to detect all PTLD (categories 1–4), advanced PTLD (categories 2–4) or neoplastic PTLD (categories 3 and 4) was estimated by analyzing the test performance at different cut-off values or with a load variation greater than 0.5 log units.

The higher diagnostic performance for identifying all, advanced or neoplastic PTLD, was achieved with cut-off values of 1.08; 1.60 and 2.47 log EBV gEq/10⁵ PBMC or 2.30; 2.60; 4.47 log gEq/10⁵ OLT cells, respectively. EBV DNA detection in plasma showed high specificity but low (all categories) or high (advanced/neoplastic categories) sensitivity for PTLD identification. Diagnostic performance was greater when: (1) a load variation in PBMC or plasma was identified; (2) combining the measure of EBV load in PBMC and plasma.

The best diagnostic ability to identify early PTLD stages was achieved by monitoring EBV load in PBMC and plasma simultaneously; an algorithm was proposed.

© 2016 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: oncovir1@anlis.gov.ar (M.D. Fellner).

PALABRAS CLAVE

Virus *Epstein-Barr*;
Detección temprana
de PTLD;
Linfoma;
PCR en tiempo real;
Pacientes
trasplantados;
Carga viral

Carga de virus *Epstein-Barr* en pacientes trasplantados: detección temprana de desórdenes linfoproliferativos postrasplante

Resumen La carga alta del virus *Epstein-Barr* se utiliza como un marcador de desórdenes linfoproliferativos postrasplante (*post-transplant lymphoproliferative disorders* [PTLD]). El objetivo de este estudio fue validar clínicamente un ensayo de cuantificación del virus *Epstein-Barr* para la detección temprana de PTLD.

Se efectuó un estudio transversal en el que se analizaron muestras pareadas de células mononucleares periféricas (CMP), de plasma y de tejido linfoide orofaríngeo de niños con trasplante de órgano sólido, con PTLD (n = 58) y sin PTLD (n = 47). En el seguimiento retrospectivo se incluyeron 71 muestras pareadas de CMP y de plasma de trasplantados, con PTLD (n = 6) y sin PTLD (n = 6). La carga viral se determinó por PCR en tiempo real. Se estimó la capacidad diagnóstica para detectar PTLD (categorías: todas vs. avanzadas vs. neoplásicas) analizando diferentes valores de corte o una variación de carga mayor de 0,5 logaritmos.

El mayor desempeño diagnóstico para identificar todos los PTLD, los avanzados y los neoplásicos, se obtuvo con valores de corte de 1,08; 1,60 y 2,47 log copias/10⁵ en CMP y de 2,30; 2,60 y 4,48 log copias/10⁵ en células de tejido linfoide orofaríngeo, respectivamente. La detección del ADN del virus *Epstein-Barr* en el plasma mostró una especificidad alta, pero una sensibilidad baja (todas las categorías) o alta (categorías avanzadas o neoplásicas) para identificar PTLD. Se observó el desempeño diagnóstico más alto en las siguientes condiciones: 1) al identificar una variación de carga en CMP o en plasma; 2) combinando la medición de la carga viral en CMP y en plasma.

La mejor capacidad diagnóstica para identificar las etapas tempranas de los PTLD se logró mediante el seguimiento simultáneo de la carga viral en CMP y en plasma; se propone un algoritmo.

© 2016 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la CC BY-NC-ND licencia (<http://creativecommons.org/licencias/by-nc-nd/4.0/>).

Introduction

The *Epstein-Barr virus* (EBV) infection is associated with the development of post-transplant lymphoproliferative disorders (PTLD) in solid-organ and stem cell transplant recipients¹⁷.

PTLD encompass a wide range of disorders including benign to malignant lymphoproliferations²². Briefly, category 1 includes early benign PTLD; category 2, benign and neoplastic PTLD and categories 3 and 4, neoplastic PTLD. It had been noted that a therapeutic intervention, particularly during the early PTLD phases, could reverse lymphoproliferation and prevent progression to the irreversible lymphoma stage³.

The association of PTLD and elevated levels of circulating EBV were first described in the nineties¹⁸. Since then, different authors showed that PTLD, especially those occurring early after transplantation, are generally associated with an increase in EBV DNA in peripheral blood samples. This has led to using the EBV load as a risk marker for this disease^{7,8,10}.

Initially, many studies explored the clinical usefulness of these tests in a single sample in the presence of disease^{19,20}. Then, the importance of surveillance through frequent repetitive monitoring was brought up¹¹. At present, a high viral load is used for two purposes: an *early diagnosis* (indicating when to start the search for disease in a potential site), and as a *prevention strategy* (indicating when to therapeutically intervene)^{6,10}. Furthermore, some authors have proposed to calculate an average load in a set time, rather

than use a single cut-off value⁶. Thus, a reduction in PTLD morbidity and mortality was reported in centers that had implemented systematic EBV load monitoring, and the viral burden was maintained at levels defined as "low", by regulating the degree of immunosuppression¹².

In the literature, the definition of a threshold indicative of a "high load" is inconsistent because each laboratory has implemented its own cut-off values to distinguish PTLD from baseline levels¹³. The broad diversity of methods (type of quantitative PCR, amplified viral fragment), type and quantity of clinical sample analyzed (peripheral blood mononuclear cells, PBMC; whole blood; plasma), controls used to standardize assays, expression of results (number of viral genome equivalents per number of cells, DNA quantity, volume), sampling frequency, among others, have made it difficult to compare published data and extrapolate cut-off values from one laboratory to another. Therefore, the overall sensitivity, specificity, positive and negative predictive values obtained by different authors relied on the population studied (adult vs. pediatric), type of transplanted organ (solid organ vs. stem cells), pre-transplant infection status, among others.

An assay to quantify EBV by real-time PCR was developed in our laboratory to monitor transplant patients⁵. Thus, considering the lack of consensus on the EBV level that identifies PTLD, the optimal type of clinical sample, and using the logistics implemented to handle transplant patients' samples in our laboratory (already underway using a semi-quantitative PCR)⁴, the EBV load in this report was

analyzed in different clinical specimens, which also represent anatomical sites of viral persistence. Therefore, the aim of this study was to clinically validate the developed EBV quantification assay for the early detection of PTLD in transplant patients.

Materials and methods

Patients and samples

Pediatric recipients of solid organ transplants treated at "Prof. Dr Juan P. Garrahan" Pediatric Hospital, Austral University Hospital, Sor María Ludovica Children's Hospital and Favalaro Foundation between 2001 and 2010 were selected for inclusion in the present study. All patients were found to be infected with EBV, as established by the previous detection of IgG antibodies against viral capsid antigen and/or EBV-DNA in PBMC. Samples were obtained according to the monitoring of transplanted patients or treatment protocol and their selection was based on clinical data (presence or absence of signs and symptoms associated with EBV infection), previous EBV load results acquired by semi-quantitative PCR⁴ (including varying EBV load levels: low, high and very high) and histopathological examination of biopsy material (PTLD or a different diagnosis based on the corresponding lymphoid tissue).

An informed consent was obtained from all patients in accordance with the Helsinki declaration and other national or international regulations. All procedures were approved by the Ethical Research Committee, National Center of Medical Genetics, ANLIS "Dr. Carlos G. Malbrán".

Transversal study design

Paired peripheral blood and oropharyngeal lymphoid tissue (OLT) samples from 105 pediatric solid-organ transplant recipients were included. Fifty-eight of them were histologically diagnosed with PTLD, according to the World Health Organization classification: IARC, 2008²² (Table 1).

Retrospective follow-up study design

Seventy one paired PBMC and plasma samples were included from six organ recipients with a histological diagnosis of PTLD, and six transplant patients with a variety of clinical conditions associated with EBV, but not with PTLD.

PBMC and plasma were separated from 2.5 to 5 ml of EDTA-anticoagulated whole blood samples by centrifugation on a density gradient (Histopaque-1077, Sigma-Aldrich) as recommended, and then stored at -20°C . The OLT from patients with lymphoid hypertrophy were surgically removed and immediately stored at -80°C .

DNA extraction

DNA from PBMC and OLT samples was obtained as previously reported⁴.

Plasma DNA was extracted using the QIAmp DNA mini kit (QIAGEN, Hilden, Germany) following manufacturers' recommendations for free viral DNA extraction.

EBV DNA quantification

We applied a duplex real-time PCR assay that simultaneously amplifies an EBNA-1 coding gene portion and one of the human glyceraldehyde 3-phosphate dehydrogenase (GAPDH) encoding genes; the amplification was performed as previously described⁵.

The primers/probes were as follows:

EBV reaction: 5' CCGCTCCTACCTGCAATATCA 3' (forward primer) and 5' GGAAACCAGGGAGGCAAATC 3' (reverse primer); 5' VIC-TGCAGCTTTGACGATGG-MGB 3' (probe). They amplified a 73 bp fragment.

GAPDH reaction: 5' GGTGGTCTCCTCTGACTTCAACA 3' (forward primer); 5' GTGGTCGTTGAGGGCAATG 3' (reverse primer) and 5' FAM-CCACTCCTCCACCTTTGACGCTGG-3' TAMRA (probe). They amplified a 79 bp fragment.

Amplification was performed in a final reaction volume of 25 μl , containing 1 \times TaqMan Universal Master Mix with AmpErase UNG (Applied Biosystems, New Jersey, USA), 0.3 μM of EBV-primers, 0.05 μM of GAPDH-primers, 0.1 μM of EBV-probe and GAPDH-probe (Applied Biosystems, Foster city, USA) and the DNA to be amplified.

For each reaction around 366 ng of DNA were assayed (corresponding to the amount of DNA of 10^5 human cells⁴) from PBMC or OLT or the extracted DNA from 30 μl of plasma. The amplification was performed using a 7500 real-time PCR System (Applied Biosystems) under the following cycle conditions: 50°C 2 min; 95°C 10 min followed by 45 cycles of 95°C 15 s, 60°C 1 min.

Table 1 Description of the transplant population

Transplanted organ	Mean of the age at transplant (years)	Histology		Mean of the time between transplant and PTLD diagnosis (month)
		Non-PTLD ^a (n)	PTLD ^b (n)	
LIVER (n = 79)	3.0 (0.5–11)	42	37	66 (20–156)
KIDNEY (n = 24)	8.4 (2–12)	5	19	54 (6–132)
HEART (n = 2)	9.5 (9–10)	0	2	66 (60–72)

The age of the transplant population ranged between 1 and 18 years (mean: 9.0 years) and the male to female rate was 1.1.

^a All patients presented a histological diagnosis of follicular lymphoid hyperplasia.

^b The PTLD cases included: Category 1: 40 lymphoid hyperplasia and 2 infectious mononucleosis-like PTLD; Category 2: 8 polymorphic and polyclonal hyperplasia; Category 3: 4 immunoblastic and 2 Burkitt lymphomas, and Category 4: 2 Hodgkin lymphomas.

The EBV load measured in PBMC DNA was expressed as the number of EBV genome equivalents (gEq) per 10^5 PBMC. The normalized EBV load was estimated from the results of the EBNA-1 and GAPDH reactions (EBNA-1 load/GAPDH load $\times 10^5$). The EBV load in plasma was expressed as the number of EBV gEq per ml of plasma by multiplying the EBNA-1 load by a factor of 33.3 and correcting by extraction efficiency.

Kinetic variation was defined as a greater than 0.5 log unit increase or decrease with respect to the EBV load measured in a previous sample. This implies a load variation greater than the overall precision of the method⁵ and also relies on the concept that a difference higher than 0.5 log units is clinically significant⁹.

Clinical validation

Transversal study: it was done in order to analyze the diagnostic performance of different cut-off values as PTLD indicators. The diagnostic ability of the assay for detecting total PTLD (including categories 1–4), advanced PTLD (categories 2–4), or neoplastic PTLD (categories 3 and 4) cases (positive reference population) was evaluated analyzing the sensitivity and specificity of different EBV load levels. The organ recipients without PTLD were considered the negative reference population. Paired PBMC, plasma and OLT samples were analyzed. The viral load corresponding to the cross-point between sensitivity and specificity was considered the most efficient cut-off value.

Retrospective follow-up study: the purpose of the retrospective follow-up study was to analyze the usefulness of the EBV load variation as PTLD indicator. The diagnostic ability of the EBV load kinetic variation for detecting total PTLD was evaluated in samples taken before and at the moment of the diagnosis for PTLD cases, and between successive samples for organ recipients without a PTLD history. The analysis was run on paired PBMC and plasma samples.

Statistical analysis

Sensitivity, specificity, positive and negative predictive values were calculated applying the Describe program, version 2.41, from the software package WinPEPI, version 11.14.

Results

The EBV load measured in PBMC, plasma and OLT samples from transplant patients with and without PTLD diagnosis is shown in Figure 1A–C, respectively.

Clinical validation

Transversal design

Peripheral blood mononuclear cells. The analysis of the ability of PBMC-measured EBV load to identify PTLD is shown in Fig. 2. The highest diagnostic efficacy for the detection of total PTLD is achieved at a cut-off value of 1.08 log gEq per 10^5 PBMC (A); for advanced PTLD, at a cut-off of 1.60 log gEq per 10^5 PBMC (B), and for neoplastic PTLD, at a cut-off of 2.47 log gEq per 10^5 PBMC (C).

Plasma. Fig. 3 describes the sensitivity and specificity of EBV levels in plasma samples for PTLD diagnosis. The EBV load in plasma samples showed low sensitivity and high specificity to detect total PTLD (Fig. 3A). While increased sensitivity, maintaining high specificity, was observed for the detection of advanced PTLD (Fig. 3B), similar results were obtained when analyzing only neoplastic PTLD (data not shown).

Lymphoid tissue. The best relations between sensitivity (Se) and specificity (Sp) for the detection of total, advanced or neoplastic PTLD were obtained at cut-off values of 2.30 log (Se = 57.4 % and Sp = 55.5 %), 2.60 log (Se = 63.6 % and Sp = 66.7 %) or 4.48 log gEq per 10^5 cells (Se = 100 % and Sp = 100 %), respectively (data not shown).

Combination of several samples. In order to improve the diagnostic performance for PTLD detection, the combination




Figure 1 EBV load in the transplanted population. (A) EBV load in peripheral blood mononuclear cells, PBMC; (B) EBV load in plasma; (C) EBV load in oropharyngeal lymphoid tissue cells, OLT.




Figure 2 Analysis of the EBV load from PBMC samples. The best diagnostic performance of the PBMC EBV load to detect different PTLD categories required increasing cut-off values. (A) Total PTLD: included categories 1–4; (B) advanced PTLD: categories 2–4; (C) neoplastic PTLD: categories 3 and 4.

Table 2 Diagnostic performance of the EBV load measured on several sample types for PTLD detection

Cut-off value	Sensitivity ^a	Specificity ^b	PPV ^c	NPV ^d
$\geq \text{Log } 1.08 \text{ gEq}/10^5 \text{ PBMC} + \geq \text{Log } 2.00/\text{ml plasma}$	87.5	60.5	70.0	60.5
$\geq \text{Log } 1.08 \text{ gEq}/10^5 \text{ PBMC} + \geq \text{Log } 2.30 \text{ gEq}/10^5 \text{ OLT}^e \text{ cells}$	67.3	52.6	80.4	35.7
$\geq \text{Log } 1.08 \text{ gEq}/10^5 \text{ PBMC} + \geq \text{Log } 2.00 \text{ gEq}/\text{ml plasma} + \geq \text{Log } 2.30 \text{ gEq}/10^5 \text{ OLT}^e \text{ cells}$	72.2	47.4	79.6	34.6

gEq: EBV genome equivalents.

^a Sensitivity = [Number of true positives / (Number of true positives + Number of false negatives)] \times 100.

^b Specificity = [Number of true negatives / (Number of true negatives + Number of false positives)] \times 100.

^c Positive predictive value = [Number of true positives / (Number of true positives + Number of false positives)] \times 100.

^d Negative predictive value = [Number of true negatives / (Number of true negatives + Number of false negatives)] \times 100.

For calculations, the histological diagnosis was considered the gold standard to which the EBV quantification load was compared.

^e OLT: oropharyngeal lymphoid tissue.

of the EBV load measured in different anatomical sites of viral persistence was analyzed (Table 2). The most effective total PTLD detection was obtained by co-measuring the EBV load in PBMC and plasma. However, the incorporation of the OLT viral load did not improve diagnostic efficacy.

Retrospective follow-up

Results from the retrospective follow-up study are shown in Figure 4 and Tables 3 and 4. The determination of the kinetic load variation increased diagnostic efficacy compared to the use of a single cut-off value (Table 5). Moreover, the




Figure 3 Analysis of the EBV load from plasma samples. The plasma EBV load showed high specificity to detect all PTLD; while an increasing sensitivity was noted as PTLD stages progressed. Total PTLD: included categories 1–4 (A); advanced PTLD: categories 2–4 (B).




Figure 4 Kinetics of EBV load in PBMC and plasma from transplant patients. Recipients with PTLT history: (A–F) Patients without PTLT history: (G–L) FLH: follicular lymphoid hyperplasia. PH: plasmacytic hyperplasia. HL: Hodgkin lymphoma. DLCBL: Diffuse large B cell lymphoma. IS: immunosuppression. IM: infectious mononucleosis. KS: Kaposi's sarcoma. Tx: time of transplantation. EBV: detection of primary EBV infection. (■) EBV load from PBMC; (▲) EBV load from plasma; x axis: sampling times.

co-measurement of the kinetic variation in more than one clinical sample improved the diagnostic ability of the assay.

Discussion

Many of the publications in which EBV load was used to identify PTLT were based on patients diagnosed with neoplastic categories^{14,15}. This would explain the high threshold levels applied by different laboratories, which could be considered

consistent with those proposed in the present study for identifying neoplastic lymphoproliferations (transversal study). Thus, considering the present data, the highest EBV levels would be particularly associated with neoplasia. The definition of high load based on a cut-off value that detects neoplastic cases may not be useful for prevention purposes. In such sense, few reports exist that describe the amount of EBV associated with the different categories of PTLT²¹. According to the present results, early PTLT stages tend to show a number of EBV genomes in all anatomical sites

Table 3 Description of the kinetic variation of EBV load in PBMC and plasma from transplant patients at PTLT diagnosis

Patients	Histological diagnosis	Kinetic load variation ^a PBMC (log differences)	Kinetic load variation ^a Plasma (log differences)
A	PH	Negative (–1.34)	Positive (1.73)
	HL	Positive (0.70)	Negative (0.30) ^b
B	DLCBL	Negative (0.37)	Positive (1.16)
C	DLCBL	Positive (0.55)	Positive (2.47)
D	PH	Positive (0.56)	Positive (0.98)
E	DLCBL	Positive (0.83)	Negative (0.15)
F	PH	Positive (0.77)	NA

PH: plasmacytic hyperplasia (Category 1 of PTLT); HL: Hodgkin lymphoma (Category 4 of PTLT); DLCBL: Diffuse large B cell lymphoma (Category 3 of PTLT).

NA: Not applicable because EBV load remained not detectable.

^a Positive kinetic variation: ≥ 0.5 log rise in EBV load; Negative kinetic variation: < 0.5 log rise or a decrease (indicated by a minus sign) in EBV load.

^b Although EBV load did not show a positive kinetic variation, when HL was diagnosed it remained in high levels, similar to those detected at the moment of PH identification.

Table 4 Description of the kinetic variation of EBV load in PBMC and plasma from transplant patients without a PTLD diagnosis

Patients	Histological diagnosis	Kinetic load variation ^a PBMC (log differences)	Kinetic load variation ^a Plasma (log differences)
G	FLH	Negative (−0.96)	NA
		Positive (1.46)	NA
		Negative (−0.74)	NA
		Positive (0.64)	NA
		Negative (−0.03)	NA
		Negative (−0.93)	NA
H	FLH	Negative (−0.97)	NA
		Positive (2.45)	NA
		Negative (−1.33)	NA
		Negative (−0.60)	NA
I	FLH	Positive (0.54)	Positive (0.63)
		Negative (−0.11)	Negative (−0.63)
		Positive (1.48)	NA
		Negative (−0.34)	NA
J	FLH	Negative (−1.24)	Negative (−2.13)
		Negative (−1.73)	NA
		Positive (2.78)	NA
		Negative (−0.95)	NA
		Positive (0.13)	Negative (0.12)
K	FLH	Negative (−1.39)	Negative (0.37)
		Positive (0.74)	Negative (−0.37)
		Negative (0.31)	NA
		Negative (−0.62)	NA
		Negative (−0.61)	NA
L	FLH	Negative (0.12)	Negative (−0.37)
		Negative (−0.97)	Positive (1.10)
		Negative (0.40)	Negative (−1.10)
		Negative (0.04)	NA

FLH: follicular lymphoid hyperplasia.

NA: Not applicable because viral load remained not detectable.

^a Positive kinetic variation: ≥ 0.5 log rise in EBV load. Negative kinetic variation: < 0.5 log rise or a decrease (indicated by a minus sign) in EBV load.

of viral persistence, lower than that observed in neoplastic categories (transversal study).

Therefore, considering that the aim of the EBV quantification in transplant patients is to detect the onset of lymphoproliferation, we wondered which monitoring strategy would be the best. Thus, a cut-off of 2.48 log gEq per 10^5 PBMC would point to neoplastic PTLD patients, but many cases with less advanced stages would be missed. A cut-off value of 1.08 log gEq per 10^5 PBMC would detect the majority of organ recipients with PTLD, but would also include many without PTLD.

Table 5 Diagnostic performance of the kinetic variation of EBV load for PTLD detection

Kinetic load variation	Sensitivity	Specificity	PPV ^a	NPV ^b
PBMC	83.3	57.7	31.3	93.7
Plasma	80	80.8	44.4	95.5
PBMC + Plasma	83.3	81.8	55.6	94.7

^a Positive predictive value.^b Negative predictive value.

It has already been proposed that a high EBV load in plasma is a more specific PTLD marker than its determination in PBMC²³; however, it is also less sensitive¹. According to the present transversal study, the detection of EBV DNA in plasma samples appears to be a sensitive and specific indicator of advanced PTLD, albeit very insensitive to identify early phases. In addition, as noted in the retrospective follow-up (Fig. 4), two patients showed undetectable EBV load in plasma at PTLD diagnosis, one in category 1 (case F) and one even in a neoplastic stage (case E).

The levels of EBV in lymphoid tissue are not usually analyzed to detect PTLD since peripheral blood samples are less invasive and more easily available. In this respect, the present data showed that a high amount of EBV DNA in lymphoid tissue exhibits the highest ability for neoplastic PTLD detection, but not for less advanced categories.

In most reports, the determination of an increased EBV load in a single blood sample type is used for PTLD identification¹; however it has also been suggested that the simultaneous evaluation of whole blood and plasma samples allows for better diagnosis and monitoring, and may be used to rule out their presence²³. Thus, among all the combinations analyzed in the present report, the simultaneous

measurement of EBV load in PBMC and plasma resulted in a greater diagnostic efficacy regarding the quantification of a single sample type (transversal study).

Moreover, as the retrospective follow-up showed the kinetic variation of the EBV levels measured in PBMC or plasma, it evidenced a better diagnostic ability than the use of a single cut-off value; these results are consistent with already reported data⁶. Furthermore, an increased risk of PTLD was demonstrated when detecting viral level variations in patients with persistent high EBV loads².

Therefore, it has been concluded that the best strategy to monitor transplant patients, regarding the detection of the early stages of PTLD and the risk of progressing to neoplastic categories would include the following algorithm. Initially, it includes the monitoring of the EBV load in PBMC, taking into account that:

- (1) An EBV load below 1.08 log gEq per 10⁵ PBMC implies low risk. Thus, the patient's routine monitoring should be continued.
- (2) An EBV load between 1.08 log and 2.48 log gEq per 10⁵ PBMC involves an increased risk. A more frequent monitoring of the EBV load in PBMC is required to detect a kinetic variation, plus the monitoring of viral levels in plasma. The detection of a kinetic variation in PBMC and/or an EBV load in plasma greater than 2 log gEq/ml involves high PTLD risk.
- (3) An EBV load above 2.48 log gEq per 10⁵ PBMC plus the detection of viral levels greater than 2.52 log gEq/ml in plasma imply a high risk of neoplastic PTLD. The greater the EBV load and/or the simultaneous detection of a kinetic variation, the greater the risk of neoplastic PTLD stages.

It should be borne in mind that the algorithm requires some consideration. Firstly, the overlap in the EBV DNA levels in PBMC among transplant patients with and without PTLD. It should be noted that according to the present results and coinciding with other authors¹⁴, not all EBV loads defined as "high" imply the development of PTLD.

Carefully monitoring an organ recipient with a low or moderate EBV load is also suggested, since a small increase in viral levels may be associated with PTLD, as was observed in this study and previously reported ones¹⁶.

The suggested algorithm aims to identify PTLD in the most effective way; however there are patients with PTLD that remain undetected and recipients without PTLD who are included as risky. This points to the need for more specific markers that, when combined with the viral load, result in a more accurate selection of cases of incipient PTLD, allowing to reverse lymphoproliferation in its initial stages.

It should be mentioned that the study presented some limitations. One of them is the variety of transplanted organs included, as they show different PTLD prevalence. However, as the EBV levels in healthy carriers were known⁵, the initial point of rising load (shared by all the transplanted organ groups) could be identified. Furthermore, the number of patients included into each PTLD category was quite different, being the initial stages of PTLD (category 1) the broader. Such category 1 just depicted the beginning of the lymphoproliferation process, which was the aim of the present

study; however, all stages of the process were analyzed separately in order to describe the differences among them.

Finally, considering that the determination of the EBV load lacks a gold standard strategy worldwide, this paper provides an in-house methodology, which is analytical, clinically validated, less expensive and more accessible than commercial assays. Thus, it may represent an initial, but very helpful step for the monitoring of transplant patients, especially in low resource settings, where commercial kits are not affordable.

Ethical disclosures

Protection of human and animal subjects. The authors declare that the procedures followed were in accordance with the regulations of the responsible Clinical Research Ethics Committee and in accordance with those of the World Medical Association and the Helsinki Declaration.

Confidentiality of data. The authors declare that they have followed the protocols of their work centre on the publication of patient data.

Right to privacy and informed consent. The authors must have obtained the informed consent of the patients and/or subjects mentioned in the article. The author for correspondence must be in possession of this document.

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

The authors are indebted to S. Nuñez and J. Campos, for their technical assistance, Drs: C. Gonzalez Cambaceres, P. Bernáldez, J. Ibañez and A. Turconi ("Prof. Dr Juan P. Garrahan" Pediatric Hospital); J. Gregoresky and M. V. Ávila (Austral University Hospital); T. Gonzalez and S. Besga ("Sor María Ludovica" Children's Hospital), J. Trentadue (Favaloro Foundation) for their generous help during sampling, and their commitment to the prevention of PTLD in transplant patients. This project was partially supported by grants from ANLIS Carlos G. Malbrán (FOCANLIS/2009) (Buenos Aires, Argentina).

References

1. Bakker NA, Verschuuren EA, Veeger NJ, van der Bij W, van Imhoff GW, Kallenberg CG, Hepkema BG. Quantification of *Epstein-Barr virus*-DNA load in lung transplant recipients: a comparison of plasma versus whole blood. *J Heart Lung Transplant*. 2008;27:7–10.
2. Bingler MA, Feingold B, Miller SA, Michaels MG, Green M, Wadowsky RM, Rowe DT, Webber SA. Chronic high *Epstein-Barr virus* viral load state and risk for late-onset posttransplant lymphoproliferative disease/lymphoma in children. *Am J Transplant*. 2008;8:442–5.
3. Cesaro S, Pegoraro A, Tridello G, Calore E, Pillon M, Varotto S, Abate D, Barzon L, Mengoli C, Carli M, Messina C. A prospective study on modulation of immunosuppression for

- Epstein-Barr virus* reactivation in pediatric patients who underwent unrelated hematopoietic stem-cell transplantation. Transplantation. 2010;89:1533–40.
4. Fellner MD, Durand K, Correa M, Bes D, Alonio LV, Teyssié AR, Picconi MA. A semi-quantitative PCR method (SQ-PCR) to measure *Epstein-Barr virus* load: its application in transplant patients. J Clin Virol. 2003;28:323–30.
 5. Fellner MD, Durand KA, Rodríguez MA, Irazu L, Alonio LV, Picconi MA. Duplex realtime PCR method for *Epstein-Barr virus* and human DNA quantification: its application for post-transplant lymphoproliferative disorders detection. Braz J Infect Dis. 2014;18:271–80.
 6. Funk GA, Gosert R, Hirsch HH. Viral dynamics in transplant patients: implications for disease. Lancet Inf Dis. 2007;7:460–72.
 7. Gärtner B, Preiksaitis JK. EBV viral load detection in clinical virology. J Clin Virol. 2010;48:82–90.
 8. Green M, Webber S. Posttransplantation lymphoproliferative disorders. Pediatr Clin North Am. 2003;50:1471–91.
 9. Grys TE. Developing a quality system for quantitative laboratory-developed tests. Clin Microbiol Newsl. 2011;33:179–85.
 10. Gulley ML, Tang W. Using *Epstein-Barr viral* load assays to diagnose, monitor, and prevent posttransplant lymphoproliferative disorder. Clin Microbiol Rev. 2010;23:350–66.
 11. Jang JY, Kim KM, Lee YJ, Lee SG, Chi HS. Quantitative *Epstein-Barr virus* viral load monitoring in pediatric liver transplantation. Transplant Proc. 2008;40:2546–8.
 12. Kerkar N, Morotti RA, Madan RP, Shneider B, Herold BC, Dugan C, Miloh T, Karabicak I, Strauchen JA, Emre S. The changing face of post-transplant lymphoproliferative disease in the era of molecular EBV monitoring. Pediatr Transplant. 2010;14:504–11.
 13. Kimura H, Ito Y, Suzuki R, Nishiyama Y. Measuring *Epstein-Barr virus* (EBV) load: the significance and application for each EBV-associated disease. Rev Med Virol. 2008;18:305–19.
 14. Kinch A, Oberg G, Arvidson J, Falk KI, Linde A, Pauksens K. Post-transplant lymphoproliferative disease and other *Epstein-Barr virus* diseases in allogeneic haematopoietic stem cell transplantation after introduction of monitoring of viral load by polymerase chain reaction. Scand J Infect Dis. 2007;39:235–44.
 15. Meerbach A, Wutzler P, Häfer R, Zintl F, Gruhn B. Monitoring of *Epstein-Barr virus* load after hematopoietic stem cell transplantation for early intervention in post-transplant lymphoproliferative disease. J Med Virol. 2008;80:441–54.
 16. Orentas RJ, Schauer DW Jr, Ellis FW, Walczak J, Casper JT, Margolis DA. Monitoring and modulation of *Epstein-Barr virus* loads in pediatric transplant patients. Pediatr Transplant. 2003;7:305–14.
 17. Rickinson AB, Kieff E. *Epstein Barr virus*. In: Fields BN, Knipe DM, Howley PM, editors. Fields virology, vol. 2, 5th ed. Philadelphia: Lippincott: Williams and Wilkins; 2007. p. 2656–700.
 18. Riddler SA, Breinig MC, McKnight JL. Increased levels of circulating *Epstein-Barr virus* (EBV)-infected lymphocytes and decreased EBV nuclear antigen antibody responses are associated with the development of posttransplant lymphoproliferative disease in solid-organ transplant recipients. Blood. 1994;84:972–84.
 19. Rooney CM, Loftin SK, Holladay MS, Brenner MK, Krance RA, Heslop HE. Early identification of *Epstein-Barr virus*-associated post-transplantation lymphoproliferative disease. Br J Haematol. 1995;89:98–103.
 20. Rowe DT, Qu L, Reyes J, Jabbour N, Yunis E, Putnam P, Todo S, Green M. Use of quantitative competitive PCR to measure *Epstein-Barr virus* genome load in the peripheral blood of pediatric transplant patients with lymphoproliferative disorders. J Clin Microbiol. 1997;35:1612–5.
 21. Schubert S, Abdul-Khalik H, Lehmkuhl HB, Yegitbasi M, Reinke P, Kebelmann-Betz C, Hauptmann K, Gross-Wieltsch U, Hetzer R, Berger F. Diagnosis and treatment of post-transplantation lymphoproliferative disorder in pediatric heart transplant patients. Pediatr Transplant. 2009;13:54–62.
 22. Swerdlow SH, Webber SA, Chadburn A, Ferry JA. Post-transplant lymphoproliferative disorders. In: Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, Thiele J, Vardiman JW, editors. WHO classification of tumours of haematopoietic and lymphoid tissues. 4th ed. Lyon, France: International Agency for Research on Cancer; 2008. p. 343–50.
 23. Tsai DE, Douglas L, Andreadis C, Vogl DT, Arnoldi S, Kotloff R, Svoboda J, Bloom RD, Olthoff KM, Brozena SC, Schuster SJ, Stadtmayer EA, Robertson ES, Wasik MA, Ahya VN. EBV PCR in the diagnosis and monitoring of posttransplant lymphoproliferative disorder: results of a two-arm prospective trial. Am J Transplant. 2008;8:1016–24.