



Revista Brasileira de Fisioterapia

ISSN: 1413-3555

rbfisio@ufscar.br

Associação Brasileira de Pesquisa e Pós-
Graduação em Fisioterapia
Brasil

Chamochoy, Carla C. M.; Nunes, Fabiana R.; Guirro, Rinaldo R. J.; Guirro, Elaine C. O.
Comparison of active and passive forces of the pelvic floor muscles in women with and without stress
urinary incontinence

Revista Brasileira de Fisioterapia, vol. 16, núm. 4, julho-agosto, 2012, pp. 314-319

Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia
São Carlos, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=235023130003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Comparison of active and passive forces of the pelvic floor muscles in women with and without stress urinary incontinence

Comparação das forças ativa e passiva dos músculos do assoalho pélvico de mulheres com e sem incontinência urinária de estresse

Carla C. M. Chamocho¹, Fabiana R. Nunes², Rinaldo R. J. Guirro³, Elaine C. O. Guirro³

Abstract

Background: The reduction of the pelvic floor muscles (PFM) strength is a major cause of stress urinary incontinence (SUI). **Objective:** To compare active and passive forces, and vaginal cavity aperture in continent and stress urinary incontinent women. **Method:** The study included a total of thirty-two women, sixteen continent women (group 1 - G1) and sixteen women with SUI (group 2 - G2). To evaluate PFM passive and active forces in anteroposterior (sagittal plane) and left-right directions (frontal plane) a stainless steel specular dynamometer was used. **Results:** The anteroposterior active strength for the continent women (mean±standard deviation) (0.3 ± 0.2 N) was greater compared to the values found in the evaluation of incontinent women (0.1 ± 0.1 N). The left-right active strength ($G1=0.43\pm0.1$ N; $G2=0.40\pm0.1$ N), the passive force ($G1=1.1\pm0.2$ N; $G2=1.1\pm0.3$ N) and the vaginal cavity aperture ($G1=21\pm3$ mm; $G2=24\pm4$ mm) did not differ between groups 1 and 2. **Conclusion:** The function evaluation of PFM showed that women with SUI had a lower anteroposterior active strength compared to continent women.

Keywords: muscle strength; pelvic floor; physical therapy; urinary incontinence.

Resumo

Contextualização: A redução da força dos músculos do assoalho pélvico (MAP) é a maior causa da incontinência urinária de estresse (IUE). **Objetivo:** Comparar as forças ativa e passiva e a abertura da cavidade vaginal em mulheres continentas e com IUE. **Método:** O estudo incluiu um total de 32 mulheres, 16 mulheres continentas (grupo 1 - G1) e 16 mulheres com IUE (grupo 2 - G2). Um espécúlo dinamométrico de aço inoxidável foi usado para avaliar as forças ativa e passiva dos MAP nas direções ântero-posterior (plano sagital) e látero-lateral (plano frontal). **Resultados:** A força ativa ântero-posterior nas mulheres continentas ($0,3\pm0,2$ N) foi maior que nas mulheres com IUE ($0,1\pm0,1$ N). Não houve diferença entre o G1 e o G2 nos valores de força ativa látero-lateral ($G1=0,43\pm0,1$ N; $G2=0,40\pm0,1$ N), força passiva ($G1=1,1\pm0,2$ N; $G2=1,1\pm0,3$ N) e abertura da cavidade vaginal ($G1=21\pm3$ mm; $G2=24\pm4$ mm). **Conclusão:** A avaliação da função dos MAP mostrou que mulheres com IUE tiveram uma menor força ativa ântero-posterior quando comparadas com as mulheres continentas.

Palavras-chave: força muscular; assoalho pélvico; fisioterapia; incontinência urinária.

Received: 01/23/2012 – Revised: 01/24/2012 – Accepted: 01/24/2012

¹Department of Gynecology and Obstetrics, Faculty of Medicine of Ribeirão Preto, Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

²Department of Obstetrics and Gynecology, School of Medicine, Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

³Department of Biomechanics, Medicine and Rehabilitation of the Locomotor System, School of Medicine of Ribeirão Preto, USP, Ribeirão Preto, SP, Brazil

Correspondence to: Elaine Caldeira de Oliveira Guirro, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Curso de Fisioterapia, Av. dos Bandeirantes, 3900, Monte Alegre, CEP 14049-900, Ribeirão Preto, SP, Brasil, e-mail: ecguirro@fmrp.usp.br

Introduction ::::

Urinary incontinence (UI) is a highly prevalent condition¹⁻³. Its evaluation and treatment are costly and affects the individual's self-esteem, social and sexual activities⁴. Although surgery for stress urinary incontinence (SUI) is an option to treat this dysfunction⁵, the conservative treatment prioritizes pelvic floor muscle (PFM) strengthening, which highlights the need for assessment methods before and after intervention⁶⁻⁸.

Strength evaluation of PFM plays a decisive role in the physical therapy treatment for SUI. In addition, it represents an important tool to monitor the clinical results, and on occasions, it can be a method to demonstrate and motivate women to contract the PFM⁹⁻¹². Morin et al.¹³ studied anteroposterior PFM force and Verelst and Leivseth¹⁴ studied left-right force in dynamometers developed by each research group of authors. In another study, Peng et al.¹⁵ measured the pressure exerted by the PFM on vaginal cavity in four directions (anterior, posterior, left and right) in women with SUI and continence, and found a difference in PFM contraction pressure only in anterior and posterior direction. Based on this proposition and on the fact that no studies were found which would assess PFM strength in the anteroposterior and left-right directions concomitantly, the presented study proposed to appraise a new method in order to support the evaluation and the physical therapy treatment in a broad and more reliable way.

Therefore, the aim of this study was to compare active and passive (force at rest) anteroposterior and left-right strengths and vaginal cavity aperture in women with and without SUI using this equipment proved to be reliable¹⁶.

Method ::::

A total of 32 women participated in this study, sixteen continent women - group 1 (G1) – mean age of 37 ± 8 and body mass index (BMI) of 23 ± 1 kg/m²; and sixteen women with stress urinary incontinence - group 2 (G2) – mean age of 48 ± 7 and BMI of 26 ± 3 kg/m². The women from G1 were recruited at the university where the study was developed and an evaluation including some questions was performed to make sure that they did not have any symptoms of urinary incontinence. The women from G2 were recruited in a center specialized in urology and the diagnosis of stress urinary incontinence of these volunteers was obtained by clinical (questions about symptoms) and urodynamic tests (to exclude women with an overactive bladder). Exclusion criteria were: pregnant women, vaginal and urinary tract infection, history of urogynecological surgery, diseases which could interfere with the PFM force measurement, allergy to latex, neurological diseases, had previously performed exercises for the PFM, prolapse of pelvic organs, urge urinary

incontinence, had taken pain killers or muscle relaxants in a week before the data collection¹⁶⁻¹⁸. This study was approved by the Research Ethics Committee of the *Universidade Metodista de Piracicaba* (UNIMEP), Piracicaba, SP, Brazil number 56/07, and all women signed an informed consent form.

For the evaluation of the active and passive PFM forces and vaginal cavity aperture measurement, a stainless steel specular dynamometer was used. It captured the PFM strength in both anteroposterior (sagittal plane) and left-right directions (frontal plane). The dynamometric device (Figure 1) was fixed to a mobile apparatus for a more effective adjustment. The device reliability was performed by the test-retest evaluation over three consecutive weeks and it was considered good and excellent. The standard error of measurement (SEM) for the mean anteroposterior strength and the mean left-right strength was 1.96 N and 1.86 N, respectively. The system was calibrated by a linear regression with weights ranging from 0 to 4 kg¹⁶.

The volunteers were evaluated in gynecological position. Before recording the data, general explanations about the pelvic floor was given and in order to check the correct PFM contraction, a 2 second contraction was requested to the volunteers, considering this contraction valid only when an observable cranial movement of the perineum has occurred¹⁹.

The PFM evaluation was divided into three steps, conducted in a single day for each woman from G1 and G2. In the first step, the dynamometric speculum was positioned up to the maximum vaginal aperture that the volunteer could tolerate, without discomfort or pain. The volunteers were instructed to relax their PFM and the passive force of these muscles was registered for 4 seconds. In the second step, with the speculum in the same position, the maximum vaginal aperture was measured (mm) with a digital paquimeter with a resolution of 0.01 mm (Digimess®). After the registration of the passive force, with the speculum at the same position, the paquimeter was placed in the speculum in order to measure vaginal aperture, always by the same researcher. In the third step, the




Figure 1. Side view of the equipment.

PFM active strength was evaluated with a 4.9 N of passive force (determined in a pilot study with 10 women, where they felt that the equipment touched the vaginal cavity without pain or discomfort), where three maximum isometric contractions of 4 seconds each were performed with a command ("squeeze, squeeze, squeeze"), instructed always by the same researcher, with a 2 minute interval, as shown in Figure 2.

The volunteers were requested to perform the PFM contraction as isolated as possible, to minimize hip adductor, gluteus and abdominal muscles interferences. Abdominal and gluteus muscles were controlled by surface electromyography (EMG), to observe their activities during the PFM contractions, and when the contraction of these muscles was observed the PFM contraction was excluded.

The EMG was carried out by simple differential electrodes made by two silver parallel bars, spaced 10 mm apart, with a pre-amplifier circuit with 20 times gain ($\pm 1\%$), a signal noise ratio lower than $3 \mu\text{V RMS}$ and a common mode rejection rate higher than 100 dB. The skin was cleaned with a solution of 70% alcohol before placing the EMG electrodes on the rectus-abdominis and gluteus muscles²⁰ and the reference electrode was fixed on the anterosuperior iliac crest with hydrosoluble gel. The EMG signal was collected by the signal acquisition module (SAM) EMG1000 (Lynx[®]), with a sampling frequency of 2000 Hz, input range of $\pm 5 \text{ V}$, a resolution of 16 bits, with a band-pass filter from 20 to 1000 Hz (Butterworth filter) and an acquisition data software Aqdados 7.2[®] (Lynx[®]). SAM was supplied by a battery and connected to a personal computer by an optical fiber to isolate the equipment from the electric grid interferences²¹.

Before the evaluation the volunteers received instructions to empty their bladder and the data collection was not performed during the pre and menstrual periods to reduce hormonal interference²².

The PFM active and passive forces were evaluated with dynamometric equipment. The equipment was covered with Olla[®] condoms, lubricated with hydrosoluble gel.

The study can be considered as blinded because data was processed in specific routines implemented on the MatLab 7.1[®] software by a researcher who did not participate in the evaluation and data collection.

Data analysis

The sample size was calculated using the software GraphPad Statmate 2.0[®] based on standard deviation data of active and passive forces of women with and without UI from Morin et al.¹³ study, with an alpha error of 0.05 and power of 80%, resulting in 16 participants in each group.

The variables age and BMI were compared between groups using Chi-square tests.

Variables such as the mean active strength, passive force and vaginal cavity aperture in continent and incontinent women were compared by using Student's *t* test for independent samples. Mann-Whitney test was applied for the independent samples for the data that were not normally distributed. Data was analyzed using SPSS 13.0[®]. The significance level of 5% was considered in all analysis.

Results

With regards to age and BMI variables, there was only difference between the mean age of the groups ($p=0.0012$).

The anteroposterior active strength for continent women (mean \pm standard deviation) ($0.3 \pm 0.2 \text{ N}$) was greater compared to the values found in the evaluation of incontinent women




Figure 2. Flowchart with the three steps of the study.

(0.1 ± 0.1 N) and indicated a significant difference for the anteroposterior active strength ($p < 0.01$) - Figure 3; while left-right active strength ($G1 = 0.43 \pm 0.1$ N; $G2 = 0.40 \pm 0.1$ N) showed non-significant difference between groups ($p = 0.2$) - Figure 4. The passive force ($G1 = 1.1 \pm 0.2$ N; $G2 = 1.1 \pm 0.3$ N) ($p = 0.89$) and the vaginal cavity aperture ($G1 = 21 \pm 3$ mm; $G2 = 24 \pm 4$ mm) ($p = 0.06$) did not differ between G1 and G2.

Discussion ...

The use of dynamometers to measure pelvic floor muscle function is recent and there are only few studies^{13,14,23} comparing the muscle strength in women with or without urinary incontinence.




Figure 3. Mean and 95% confidence intervals (CI) of the anteroposterior active strength (APS) of the pelvic floor muscles of the continent and incontinent groups ($p < 0.05$).




Figure 4. Mean and 95% confidence interval (CI) of the left-right active strength (LRS) of the pelvic floor muscles of the continent and incontinent groups ($p > 0.05$).

The difference observed between the PFM anteroposterior active strength in continent and stress urinary incontinent women supports the hypothesis that the strength of these muscles is related to the continence function. These results were not in agreement with the study of Morin et al.¹³ and DeLancey et al.²³, which evaluated the anteroposterior active strength with a dynamometer and did not observe significant differences in relation to the maximum strength of continent and incontinent groups. It can be assumed that these disagreements could be related to the evaluation methodology applied as well as the difference in equipment used in both studies. In the Morin et al.¹³ study, statistical analyses were adjusted between group differences for age and parity. In addition, they looked at many different variables specific to force production, where they found differences in endurance and rate of force development, but not peak strength.

Strength evaluation in different directions simultaneously was only performed in one study²⁴, which had not compared continent and urinary incontinent women. According to Nishiyama et al.²⁵ pressure is directly proportional to strength even though they are different measures. Many studies^{11,26,27} with pressure manometers, also known as perineometers, compared the PFM pressure of continent and incontinent women and found a higher pressure in continent women corroborating with the results of the PFM anteroposterior active strength obtained in the present study.

Verelst and Leivseth²⁸ tested dynamometric equipment that measures strength in left-right direction. They reported that the resulting force applied by the PFM runs in an anteroposterior direction, therefore it would be the most appropriate direction to evaluate the PFM strength even though the strength applied in the left-right direction is important for a greater understanding of the PFM contraction behavior as the force applied by these muscles does not occur in one direction only. In 2007, the same authors¹⁴ compared the strength of women with continence and SUI using this equipment and did not observe any significant difference in left-right active strength between the groups, in agreement with results obtained in the present research. Peng et al.¹⁵ used a vaginal probe with four pressure sensors to measure the direction and amount of pressure applied in the vagina at rest, and during contractions of the PFM in 23 continent women and in 10 women with SUI. The authors observed that the pressure in the anteroposterior direction was significantly greater than that exerted on the left-right direction between the two groups, confirming the results obtained in this study.

The results found for the values of passive force between the continent and stress urinary incontinent groups were not in agreement Morin et al.¹³ study, which had evaluated the passive force and found lower values in incontinent

women when compared with the continent sample. Several factors may have influenced the values obtained from the passive force evaluation such as vaginal aperture, instruments used and the population studied.

Verelst and Leivseth¹⁴ and DeLancey et al.²³ had quantified the passive force in women with SUI and continence and had not observed any between-group differences, which corroborate with the results of passive force found in the present study.

Rahn et al.²⁹ reported that the vagina has a property called anisotropy, which means that the vagina is directionally dependent on the load applied to it, which may explain the differences found in strength in many studies as the instruments used were not the same. Besides, also based on that property, the way of the instrument was coupled to the vaginal cavity can also generate differences in the results observed³⁰.

The dynamometer used in this study measured the PFM strength in anteroposterior and left-right directions with a variable opening (in mm), while other studies with dynamometers used a predetermined opening of the equipment^{13,14,24}. This variable aperture in mm means that for each woman the opening (in mm) was one, and to maintain uniformity of the sample a value of the passive force was standardized (in N). Therefore, the recording of the active force was performed after opening the equipment to 4.9 N for all women ensuring that the force was not influenced by differences in the equipment fitted from one woman to another. As soon as determining the value of opening equal for all volunteers (mm), it probably

allows a better attachment in women who only need to apply a lesser contraction compared to women who the attachment was not very satisfactory³⁰.

The between-group difference in age is also a difficulty found in other studies. Morin et al.¹³ in their study comparing the strength of the PFM in the continent and incontinent groups reported that even with an identical procedure for both groups, significant differences were found in the variables age and parity. Verelst and Leivseth¹⁴ also found significant difference in age between the continent and incontinent women. Hannestad et al.³¹ demonstrated an increased prevalence of UI with advancing age.

One probably limitation of this study is the maximum PFM contraction that can include contraction of other muscles, especially rectus abdominis. In this study, we asked the volunteers to contract the PFM at the maximum contraction they could perform without the interference of other muscles, which could not stimulate the volunteers to perform a maximum contraction of the muscles studied as the purpose of the study was to evaluate the PFM itself without the influence of other muscle groups with the aim of achieving reliable data.

It would not be possible to evaluate women who have discomfort or pain while using the equipment to measure the PFM strength.

The current research highlights that there is a different behavior of PFM in women who are continent and incontinent, in relation to anteroposterior active strength.

References

- Keller SL. Urinary incontinence: occurrence, knowledge, and attitudes among women aged 55 and older in a rural Midwestern setting. *J Wound Ostomy Continence Nurs.* 1999;26(1):30-8.
- Guarisi T, Pinto Neto AM, Osís MJ, Pedro AO, Paiva LHC, Faúndes A. Incontinência urinária entre mulheres climatéricas brasileiras: inquérito domiciliar. *Rev Saúde Pública.* 2001;35(5):428-35.
- Kenton K, Mueller ER. The global burden of female pelvic floor disorders. *BJU Int.* 2006;98 Suppl 1:1-5.
- Wagner TH, Hu TW. Economic costs of urinary incontinence in 1995. *Urology.* 1998;51(3):355-61.
- DeLancey JO. The pathophysiology of stress urinary incontinence in women and its implications for surgical treatment. *World J Urol.* 1997;15(5):268-74.
- Dumoulin C, Bourbonnais D, Lemieux MC. Development of a dynamometer for measuring the isometric force of the pelvic floor musculature. *Neurourol Urodyn.* 2003;22(7):648-53.
- Bø K, Sherburn M. Evaluation of female pelvic-floor muscle function and strength. *Phys Ther.* 2005;85(3):269-82.
- Balmforth JR, Mantle J, Bidmead J, Cardozo L. A prospective observational trial of pelvic floor muscle training for female stress urinary incontinence. *BJU Int.* 2006;98(4):811-7.
- Peschers UM, Ginkelmaier A, Jundt K, Leib B, Dimpfl T. Evaluation of pelvic floor muscle strength using four different techniques. *Int Urogynecol J Pelvic Floor Dysfunct.* 2001;12(1):27-30.
- Bø K. Pelvic floor muscle strength and response to pelvic floor muscle training for stress urinary incontinence. *Neurourol Urodyn.* 2003;22(7):654-8.
- Amaro JL, Moreira EC, Gameiro MOO, Padovani CR. Pelvic floor muscle evaluation in incontinent patients. *Int Urogynecol J Pelvic Floor Dysfunct.* 2005;16(5):352-4.
- Rett MT, Simões JA, Herrmann V, Marques AA, Morais SS. Existe diferença na contratilidade da musculatura do assoalho pélvico feminino em diversas posições? *Rev Bras Ginecol Obstet.* 2005;27(1):12-9.
- Morin M, Bourbonnais D, Gravel D, Dumoulin C, Lemieux MC. Pelvic floor muscle function in continent and stress urinary incontinent women using dynamometric measurements. *Neurourol Urodyn.* 2004;23(7):668-74.
- Verelst M, Leivseth G. Force and stiffness of the pelvic floor as function of muscle length: a comparison between women with and without stress urinary incontinence. *Neurourol Urodyn.* 2007;26(6):852-7.
- Peng Q, Jones R, Shishido K, Omata S, Constantinou CE. Spatial distribution of vaginal closure pressures of continent and stress urinary incontinent women. *Physiol Meas.* 2007;28(11):1429-50.
- Nunes FR, Martins CC, Guirro ECO, Guirro RRJ. Reliability of bidirectional and variable-opening equipment for the measurement of pelvic floor muscle strength. *PM R.* 2011;3(1):21-6.
- Dumoulin C, Gravel D, Bourbonnais D, Lemieux MC, Morin M. Reliability of dynamometric measurements of the pelvic floor musculature. *Neurourol Urodyn.* 2004;23(2):134-42.
- Hundley AF, Wu JM, Visco AG. A comparison of perineometer to brink score for assessment of pelvic floor muscle strength. *Am J Obstet Gynecol.* 2005;192(5):1583-91.
- Bø K. Pelvic floor muscle exercise for the treatment of female stress urinary incontinence: validity of vaginal pressure measurements of pelvic floor muscle strength and the necessity of supplementary methods for control of correct contraction. *Neurourol Urodyn.* 1990;9(5):479-87.

20. Cram JR, Kasman GS, Haltz J. Introduction to surface electromyography. Maryland: Aspen Publishers; 1998.
21. Guirro RRJ, Forti F, Bigaton DR. Proposal for electrical insulation of the electromyographic signal acquisition module. *Electromyogr Clin Neurophysiol*. 2006;46(6):355-63.
22. Sarwar R, Niclos BB, Rutherford OM. Changes in muscle strength, relaxation rate and fatiguability during the human menstrual cycle. *J Physiol*. 1996;493(Pt 1):267-72.
23. DeLancey JOL, Trowbridge ER, Miller JM, Morgan DM, Guire K, Feener DE, et al. Stress urinary incontinence: relative importance of urethral support and urethral closure pressure. *J Urol*. 2008;179(6):2286-90.
24. Constantinou CE, Omata S. Direction sensitive sensor probe for the evaluation of voluntary and reflex pelvic floor contractions. *Neurourol Urodyn*. 2007;26(3):386-91.
25. Nishiyama M, Kimura Y, Nishiyama Y, Terazima M. Pressure-induced changes in the structure and function of the kinesin-microtubule complex. *Biophys J*. 2009;96(3):1142-50.
26. Mørkved S, Salvesen KA, Bø K, Eik-Nes S. Pelvic floor muscle strength and thickness in continent and incontinent nulliparous pregnant women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2004;15(6):384-90.
27. Thompson JA, O'Sullivan PB, Briffa NK, Neumann P. Assessment of voluntary pelvic floor muscle contraction in continent and incontinent women using transperineal ultrasound, manual muscle testing and vaginal squeeze pressure measurements. *Int Urogynecol J Pelvic Floor Dysfunct*. 2006;17(6):624-30.
28. Verelst M, Leivseth G. Force-length relationship in the pelvic floor muscles under transverse vaginal distension: a method study in healthy women. *Neurourol Urodyn*. 2004;23(7):662-7.
29. Rahn DD, Ruff MD, Brown SA, Tibbals HF, Word RA. Biomechanical properties of the vaginal wall: effect of pregnancy, elastic fiber deficiency, and pelvic organ prolapse. *Am J Obstet Gynecol*. 2008;198(5):590.e1-6.
30. Nunes FR, Guirro ECO, Martins CC, Guirro RRJ. Influence of visual feedback on pelvic floor muscle strength. *Eur J Obstet Gynecol Reprod Biol*. 2010;151(2):217-20.
31. Hannestad YS, Rortveit G, Sandvik H, Hunskaar S; Norwegian EPINCONT study. Epidemiology of Incontinence in the County of Nord-Trøndelag. A community-based epidemiological survey of female urinary incontinence: The Norwegian EPINCONT Study. *Epidemiology of Incontinence in the County of Nord-Trøndelag. J Clin Epidemiol*. 2000;53(11):1150-7.