



Revista Brasileira de Enfermagem

E-ISSN: 1984-0446

reben@abennacional.org.br

Associação Brasileira de Enfermagem  
Brasil

Milani Nespollo, Alice; Reschetti Marcon, Samira; Pollo de Lima, Nathalie Vilma; Lebre  
Dias, Tatiane; Martínez Espinosa, Mariano

Health Conditions and Memory Performance: a study with older adult women

Revista Brasileira de Enfermagem, vol. 70, núm. 3, mayo-junio, 2017, pp. 640-646

Associação Brasileira de Enfermagem

Brasília, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=267051078027>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## Health Conditions and Memory Performance: a study with older adult women

*Condições de saúde e desempenho da memória: um estudo com idosas*  
*Condiciones de la salud y del desempeño de la memoria: un estudio con mujeres mayores*

Alice Milani Nespollo<sup>I,II</sup>, Samira Reschetti Marcon<sup>III</sup>, Nathalie Vilma Pollo de Lima<sup>II</sup>,  
Tatiane Lebre Dias<sup>III</sup>, Mariano Martínez Espinosa<sup>IV</sup>

<sup>I</sup> Universidade Federal de Mato Grosso, Faculty of Nursing, Postgraduate Program in Nursing. Cuiabá, Mato Grosso, Brazil.

<sup>II</sup> Universidade Federal de Mato Grosso, Faculty of Nursing, Center for Mental Health Studies. Cuiabá, Mato Grosso, Brazil.

<sup>III</sup> Universidade Federal de Mato Grosso, Institute of Education, Postgraduate Program in Psychology. Cuiabá, Mato Grosso, Brazil.

<sup>IV</sup> Universidade Federal de Mato Grosso, Department of Statistics,  
Postgraduate Program in Public Health. Cuiabá, Mato Grosso, Brazil.

### How to cite this article:

Nespollo AM, Marcon SR, Lima NVP, Dias TL, Martínez Espinosa M. Health Conditions and Memory Performance: a study with older adult women. Rev Bras Enferm [Internet]. 2017;70(3):640-6. DOI: <http://dx.doi.org/10.1590/0034-7167-2016-0529>

Submission: 11-04-2016

Approval: 02-20-2017

### ABSTRACT

**Objective:** to verify the correlation between health condition and memory performance of older adult women in the community. **Method:** Analytical cross-sectional study developed with 28 older adult women living in Cuiabá-MT. They answered the Mini-Mental State Examination (MMSE), and a shortened Geriatric Depression Scale (GDS-15) to screen for dementia and depression symptoms. Memory skills were assessed through Rey Auditory Verbal Learning Test (RAVLT). **Results:** The mean age was 66.36 years and 75% of the participants had educational level higher than 7 years. The MMSE mean score was 28.45. The correlations found were: educational level and immediate memory ( $r = 0.49$ ;  $p = 0.008$ ); delayed recall and immediate memory ( $r = 0.71$ ;  $p < 0.001$ ); memory recognition and immediate memory ( $r = 0.43$ ;  $p = 0.021$ ) and recognition memory with delayed recall ( $r = 0.47$ ;  $p = 0.012$ ). **Conclusion:** High scores in the MMSE and a satisfactory health perception among the participants were evident. There was no correlation between memory performance and health perception. **Descriptors:** Aged; Cognition; Memory; Neuropsychological Tests; Health Condition.

### RESUMO

**Objetivo:** Verificar a correlação entre condições de saúde e desempenho da memória de idosas da comunidade. **Método:** Estudo transversal analítico, realizado com 28 idosas residentes em Cuiabá-MT. Essas foram submetidas a rastreio para demências e sintomas depressivos por meio do Miniexame do Estado Mental (MEEM) e Escala de Depressão Geriátrica Abreviada (GDS-15). A avaliação das habilidades de memória ocorreu por meio do Teste de Aprendizagem Auditivo-Verbal de Rey (RAVLT). **Resultados:** A idade média foi de 66,36 anos e 75% possuíam escolaridade maior que sete anos. A média do MEEM foi 28,45. As correlações encontradas foram: escolaridade e memória imediata ( $r = 0,49$ ;  $p = 0,008$ ); evocação tardia e memória de reconhecimento com memória imediata ( $r = 0,71$ ;  $p < 0,001$  e  $r = 0,43$ ;  $p = 0,021$ ) e memória de reconhecimento com evocação tardia ( $r = 0,47$ ;  $p = 0,012$ ). **Conclusão:** Evidenciou-se escore elevado no MEEM e percepção de saúde satisfatória entre os participantes. Não houve correlação entre desempenho da memória e percepção de saúde. **Descritores:** Idosos; Cognição; Memória; Testes Neuropsicológicos; Nível de Saúde.

### RESUMEN

**Objetivo:** Comprobar la interrelación entre las condiciones de salud y el desempeño de la memoria de mujeres mayores de la comunidad. **Método:** Estudio transversal analítico, realizado con 28 mujeres mayores residentes en Cuiabá-Mato Grosso. Dichas mujeres se sometieron a sondeo para demencias y síntomas depresivos por medio del Mini Test del Estado Mental (MEEM, sigla en inglés) y de la Escala de Depresión Geriátrica Abreviada (GDS-15). La evaluación de las habilidades de la memoria se realizó por medio de la Prueba de Aprendizaje Auditivo-Verbal de Rey (RAVLT). **Resultados:** La edad promedio era de 66,36

años y 75% poseían escolaridad mayor que siete años. La media del MEEM fue del 28,45. Las interrelaciones encontradas fueron las siguientes: escolaridad y memoria inmediata ( $r = 0,49$ ;  $p = 0,008$ ); evocación tardía y memoria de reconocimiento con memoria inmediata ( $r = 0,71$ ;  $p < 0,001$  y  $r = 0,43$ ;  $p = 0,021$ ) y memoria de reconocimiento con evocación tardía ( $r = 0,47$ ;  $p = 0,012$ ). **Conclusión:** Se hizo evidente el marcador elevado del MEEM y la percepción de salud satisfactoria entre los participantes. No hubo interrelación entre el desempeño de la memoria y la percepción de la salud.

**Descriptores:** Personas Mayores; Cognición; Memoria; Pruebas Neuropsicológicas; Nivel de Salud.

## CORRESPONDING AUTHOR

Alice Milani Nespollo

E-mail: alicenespollo@gmail.com

## INTRODUCTION

The decrease in fertility and infant mortality rate and the increase in the life expectancy of the world population have led to a change in the age pyramid in recent years and to a significant increase in older adults' population<sup>(1-2)</sup>. In 2010, the number of people over 65 years old was 524 million, representing 8% of the world's population. It is estimated that by the year 2050 this number will triple, reaching about 1.5 billion older adults and representing 16% of the world population<sup>(3)</sup>.

In Brazil, in 2010, the number of people aged 60 years old or over was 20,588,890, representing just over 10% of the country's population<sup>(4)</sup>. It is estimated that by 2025 the number of older adults will be 32 million, and in 2050 they will represent 22.71% of the Brazilian population<sup>(2)</sup>.

The increase in older adults' population, along with the consequent aging process, can cause small incapacities or major limitations to individuals in functional, psychic and cognitive aspects, directly interfering in the health conditions of the elderly<sup>(2)</sup>.

Health conditions are defined as the different health circumstances, with varying duration, that need answers from society, the healthcare system, professionals and users<sup>(5)</sup>. Several variables are related to and may affect health conditions of older adults, such as use of medications<sup>(6)</sup>, health self-perception, practice of physical activities<sup>(7)</sup>, physical illnesses and, especially, cognitive impairment<sup>(6-7)</sup>.

Some cognitive functions are more affected by advancing age, such as attention, executive function and memory<sup>(8)</sup>. The latter is considered the most susceptible to the aging process<sup>(9)</sup>. Memory is characterized by the ability to retain past information, store it and use it in the present. It is analyzed hierarchically, according to the time of retention of the stored information (short and long term) and the type of information<sup>(9)</sup>.

The decline of cognitive functions in the elderly is an expected event and does not significantly compromise daily living activities. However, this decline may be more severe, characterizing dementia, which affects the execution of simple daily tasks and directly interferes in the elderly's health conditions<sup>(10)</sup>.

A study conducted in the United States with 3,030 individuals between 34 and 85 years old - aimed at assessing the relationship between subjective memory complaints and the variables physical activity, cognitive activity, social activity and perceived health status - showed that, among the older adults, those individuals who presented subjective memory complaints had lower educational level and poor health perception<sup>(11)</sup>.

It is important to emphasize that elderly people with some level of cognitive impairment have a perception of their health

status different from cognitively healthy elderly<sup>(12)</sup>, meaning they overestimate their possible alterations. This way, cognitive evaluation through neuropsychological instruments, such as tests, scales and questionnaires, has been an efficient tool to assess overall cognitive ability and specific cognitive functions<sup>(8)</sup>, providing an accurate diagnosis of cognitive conditions. Neuropsychological examination assists in the diagnosis of cognitive decline (which usually precedes dementia) and in the differentiation and classification of dementia<sup>(13)</sup>.

Considering the above, assessing memory performance in its aging process is fundamental for the health professional to be able to identify early cognitive changes of older adults and interfere with actions to improve health conditions and, consequently, quality of life of this population.

## OBJECTIVE

To verify the correlation between health conditions and memory performance in older adult women in the community.

## METHOD

## Ethical aspects

This study is part of a quasi-experimental research approved by the Research Ethics Committee of the Universidade Federal de Mato Grosso - Campus Araguaia, according to Resolution 466/2012, that concerns research involving human beings. Anonymity of the participants was guaranteed, as well as the possibility to withdraw from the study at any time. Participants were informed about the condition of voluntary participation and signed an Informed Consent Form.

## Study design, location and period

This is an analytical cross-sectional study, conducted at the main campus of UFMT (Universidade Federal do Mato Grosso), in Cuiabá, capital of the state of Mato Grosso. The study was conducted from February to April 2016.

## Population or sample; inclusion and exclusion criteria

The sample consisted of women living in Cuiabá-MT and surrounding cities, invited to participate in an extension project developed by UFMT professors, called "Healthy Longevity Program".

The project is linked to the Older Adults University (UNATI - *Universidade da Terceira Idade*). It is open to the community and includes individuals over 60 years of both genders. It has about 300 older adults, of which approximately 95% are women, which determined the choice for the female gender

in the present study. The activities developed are water-based exercises, resistance training, among others, besides computer, language and singing classes.

Information was disseminated through social media in order to recruit individuals for the study. The selection had the following inclusion criteria: women; aged 60 years or over; participating for the first time in the "Healthy Longevity Program"; completed at least the first four years of formal education; present a score in the Mini-Mental State Examination (MMSE)<sup>(14)</sup> above the cutoff score for the educational level; present a score < 5 on the shortened Geriatric Depression Scale (GDS-15)<sup>(15)</sup>.

MMSE is an instrument used to screen for dementia by assessing orientation to time and to space, attention and calculation, registration recall and language. The cutoff score is related to educational level<sup>(16)</sup>. GDS-15 is also a screening scale, used for depression symptoms. It contains 15 questions for older adults, and scores over 5 suggest the presence of depressive symptoms<sup>(15)</sup>. Both tests were used to select participants in order to avoid a possible alterations in the neuropsychological test score, due to the presence of mild cognitive impairment, dementia and/or depression, considering that these pathologies might act as effect modifiers.

The following exclusion criteria were established: having a neurological history; using psychotropic medications; presence of auditory and/or visual limitations without corrective devices; difficulty in walking without the use of walking aids; and motor deficiency and/or musculoskeletal disorders that make practicing physical activities impossible.

### Study Protocol

According to the inclusion and exclusion criteria, a total of 41 volunteers were screened through application of the MMSE, the GDS-15 and a questionnaire addressing sociodemographic characteristics and health conditions. The questionnaire was divided in three parts: sociodemographic characteristics (age and educational level); health conditions (health perception and health problems); living habits (practice of physical activities). Time of application of the screening instruments was approximately 30 minutes. In this stage, two candidates were excluded due to their use of antidepressants and anxiolytics, one for having symptoms of depression and five because they were illiterate. Therefore, the sample was composed by 33 participants.

After the selection, neuropsychological evaluations were applied individually in an appropriate room in the Applied Psychology Service of the UFMT, with an evaluator trained and supervised by a UFMT teaching psychologist and member of the research. In this stage, five participants withdrew themselves from the research due to incompatibility of schedules or personal reasons. Finally, the final sample consisted of 28 older adult women.

The instrument used to evaluate memory was the Rey Auditory-Verbal Learning Test (RAVLT), adapted and validated for Brazilian older adults population<sup>(17)</sup>. It consists of a list of 15 words (list A), read aloud and followed by an immediate recall test (A1). This procedure is repeated five times (A2, A3, A4 and A5). After that, another list of words (list B) is read and followed by another recall test (B1). After application of list B, the participant is asked to remember as many words

as possible from list A (A6). This process is repeated after a 20-minute interval (A7). Finally, a list of 50 words containing list A, list B and 20 other words, semantically or phonetically similar to those of both lists, is presented to the participant, who is asked to recognize those belonging to the first list; in this step the recognition memory is evaluated<sup>(17)</sup>.

This test is recognized in the literature for its sensitivity in detecting memory deficits and, therefore, it is widely used. In this study the variables of delayed recall (A7) and recognition memory (REC) were used for the evaluation of long-term memory. Short-term memory was evaluated by the variable immediate memory (A1)<sup>(17)</sup>. These memory cognitive abilities are susceptible to the aging process<sup>(8)</sup>.

### Result Analysis and statistics

Data were analyzed in the software Statistical Package For The Social Sciences (SPSS) 17.0. To verify the distribution of the data, the Shapiro-Wilk test was applied. Non-normality was found for the immediate memory variable, and, therefore, non-parametric tests were used for the bivariate analysis.

The results of the continuous variables were presented as mean  $\pm$  standard deviation, according to distribution. Categorical variables were expressed as proportions. The Spearman Correlation Coefficient was used to analyze the correlation between numerical and nominal variables. The level of significance was set at 5%.

### RESULTS

The mean age of the 28 women evaluated was 66.36 years and 75% of them had seven years or more of formal education. MMSE mean score was 28.45. A total of 60.7% of the women considered their health as good and 25% as great; 78.6% of them reported having at least one health problem. Regarding physical activity, 53.6% of them reported not practicing any kind of exercise (Table 1).

**Table 1** – Health conditions characteristics of the sample, Cuiabá, Mato Grosso, Brazil, 2016 (N = 28)

	Mean $\pm$ SD**	n (%)
Age	66.36 $\pm$ 6.12	
MMSE*	28.45 $\pm$ 1.60	
Educational level		
1–3		1 (3.6)
4–7		6 (21.4)
> 7		21 (75.0)
Perception of health		
Regular		4 (14.3)
Good		17 (60.7)
Great		7 (25.0)
Health problems		
None		6 (21.4)
One or more		22 (78.6)

Note: \*MMSE: Mini-Mental State Examination; \*\*SD: Standard Deviation.



The descriptive results of the neuropsychological tests are shown in Table 2. The mean of number of words recalled in immediate memory test was 6.04; in the delayed recall test the mean was 8.57; and for memory recognition of the word list the mean was 12.82.

Table 3 presents the correlations between neuropsychological tests and the variables of health conditions, verifying correlation between educational level and immediate memory ( $R = 0.49$ ,  $p = 0.008$ ); delayed recall and immediate memory ( $R = 0.71$ ,  $p < 0.001$ ); recognition memory and immediate memory ( $R = 0.43$ ,  $p = 0.021$ ); and recognition memory and delayed recall ( $R = 0.47$ ,  $p = 0.012$ ).

**Table 2 –** Descriptive statistics of performance in the neuropsychological tests, Cuiabá, Mato Grosso, Brazil, 2016 (N = 28)

Tests	Mean $\pm$ SD*	Minimum	Median	Maximum
Immediate memory	6.04 $\pm$ 1.83	2	6	9
Delayed recall	8.57 $\pm$ 2.96	3	8	13
Memory recognition	12.82 $\pm$ 2.39	6	13	15

Note: SD\*: Standard Deviation.

**Table 3 –** Correlation of the variables age, educational level, health perception and neuropsychological tests, Cuiabá, Mato Grosso, Brazil (N = 28)

	Age	Educational attainment	Immediate memory	Delayed recall	Memory recognition
Educational level	-0.33	-			
Immediate memory	-0.25	0.49**	-		
Delayed recall	-0.27	0.36	0.71**	-	
Memory recognition	0.01	0.35	0.43*	0.47*	-
Perception of health	-0.18	0.22	0.19	0.27	0.24

Note: \* Correlation is significant at the 0.05 level (2-tailed); \*\* Correlation is significant at the 0.01 level (2-tailed)

## DISCUSSION

The participants of the present study had high scores in the MMSE and educational level mostly over seven years, corroborating evidence from the literature that MMSE performance is strongly influenced by the years of formal education<sup>(14,16,18)</sup>. Educational level is also considered an aspect of neuronal protection<sup>(18)</sup>.

This fact is explained by studies with Brazilian population, such as a study conducted in the city of Uberlândia-MG with adults and older adults, which verified that people with higher educational levels obtain higher scores in the MMSE. The scores varied from 25.56 for 1-4 years of formal education to 28.35 for individuals with more than 12 years

of formal education<sup>(18)</sup>. Another study, with 20 older adults in the Geriatric Ambulatory Services of the *Hospital das Clínicas* of the *Faculdade de Medicina de São Paulo* showed a mean MMSE score of 28.35 correlated to a mean of 7.75 years of formal education<sup>(19)</sup>, similar to the results found in the present study.

The variable “educational level” was also compared to MMSE scores in international studies. A study conducted in Japan with 47 participants with mean educational level of 13 years<sup>(20)</sup> and another one, conducted in Spain with 20 participants with mean education level of 13.43 years<sup>(21)</sup>, found mean MMSE scores of 28.20 and 28.86 respectively, showing results similar to the present research, even though educational level was way higher.

Considering the above, the five-year difference of educational level compared to the international studies does not seem to have a significant influence in the MMSE scores. The lack of impact in the test result of this difference in formal education might have happened due to the low mean age of the population of the present study. Studies indicate that age is also directly related to MMSE scores, since aging is one of the major risk factor for dementia<sup>(1,22)</sup>; therefore, higher ages make the individual more likely to have a lower performance in the MMSE. In this case, the sample of these international studies had mean ages higher than the sample of this study. Therefore, the findings reinforce the evidences found in the literature regarding influence of educational levels and age in MMSE performance<sup>(14,23)</sup>.

Although the results of the MMSE are presented in this paper - and this instrument was used only for the purpose of screening for cognitive impairment, and not as the main instrument of evaluation of the object studied - the discussion of the findings is important since studies conducted in Brazil use the test extensively, but still present contradictory results due to the non-existence of a standardized version<sup>(24)</sup>. Also, Brazilian studies addressing cognitive evaluation generally use the MMSE as the only assessment instrument, unlike international studies, that conduct cognitive examination using specific instruments for each function. However, this study was concerned with the specificity of the evaluation of memory performance, using a unique instrument to assess this function, unlike the majority of studies conducted with the same population in Brazil.

Regarding health perception of the elderly, most of the participants classified their health as “good”. This finding is in agreement with the literature, since evidence found in a study suggests that a negative health self-perception is closely related to cognitive decline<sup>(25)</sup>. The high percentage of “good” and “great” health self-assessments (80%) in this study is explained by the fact that the sample did not present evidence of cognitive deficit, had functional capacity preserved and were active and socially engaged.

A household survey conducted in three cities of different Brazilian regions, aimed at analyzing factors associated to a negative health perceptions of Brazilian older adults population, found a significant association between a negative health self-perception and use of two or more medications, hypertension and functional disability<sup>(26)</sup>. Therefore, health self-perception can be considered a reflection of the overall health situation of the elderly and directly related to autonomy and independence<sup>(26)</sup>. Furthermore, it is an important indicator of mortality, since those that perceive their health as bad have higher chances of dying than those that report health as good<sup>(27)</sup>.

When asked about health problems, most of the participants reported having one health problem or more. Interestingly, even though the participants had some type of disease, positive health perception was prevalent. The same result was found in a study that aimed to evaluate the health self-perception of 274 older adults in the city of Passo Fundo-RS<sup>(28)</sup>. This finding can be explained by the fact that these individuals do not have disabilities or limitations, and most of the times they live without any type of symptoms<sup>(29)</sup>. This corroborates the idea that, for the older adults, a bad health condition is associated with functional decline and need for help in the execution of activities, associating bad health condition with dependence and not necessarily with the presence and/or number of diseases<sup>(26)</sup>.

In the evaluation of immediate memory, the mean score found in this study is similar to the results of the validity study of the RAVLT test for Brazilian women between 65 and 69 years old<sup>(17)</sup>. The mean scores of delayed recall and memory recognition were respectively lower and higher than the scores obtained in the reference study. Another study conducted in the State of Parana with 57 individuals of both genders aged 65 or more found a mean of immediate memory of 5.1, a delayed recall mean of 7.1 and a memory recognition mean of 12.2, all lower to the results found in the present study<sup>(30)</sup>.

A positive correlation between educational level and immediate memory performance in the RAVLT test ( $p < 0.01$ ) was found in this study. This means that when the level of formal education increases the number of words recalled after the first reading of the word list also increases. This finding is similar to the result obtained in a study with 158 Brazilian older adults from Juiz de Fora-MG. This study aimed at verifying the correlation between cognitive tests obtained by the CERAD (Consortium to Establish a Registry for Alzheimer's Disease) and sociodemographic variables. A correlation ( $p < 0,001$ ) between educational level and immediate memory was also found<sup>(23)</sup>. The study by Magalhães and Hamdam<sup>(30)</sup> also indicated the same correlation.

Although not fully understood yet, the educational level seems to be a protective factor for the decline not only of memory performance, but also of cognitive function in general. This assertion can be explained by the cognitive reserve model, meaning that a more complex and effective neuronal network, attained through a higher education time, might compensate for neuronal and synaptic losses related to aging<sup>(22)</sup>.

Regarding the analysis of the correlation between delayed recall and recognition memory and between immediate memory and recognition memory, the high results obtained were corroborated in a study with Brazilian older adults ( $p \leq 0.05$  e  $p \leq 0.001$ )<sup>(23)</sup>. This means that as the number of words recalled in the first reading of the word list increases, number of words recalled after the 20-minute delay and distinguished among the 50 words in the recognition list also increase. Interestingly, even with the use of different instruments for the evaluation of the same variables, the results found are similar.

Health perception presented a positive correlation with educational level and immediate memory in the study conducted in Juiz de Fora-MG<sup>(23)</sup>. However, in the present study, no correlation was found between the variable "health perception" and the results of the neuropsychological test.

This study showed no correlation between the performance on the neuropsychological test and health perception, unlike a study conducted in Belo Horizonte-MG with 98 older adults that found that individuals with no cognitive impairment found in the MMSE tended to present a bad health perception ( $p = 0,006$ )<sup>(25)</sup>. A possibility for the correlations of these studies might be the fact that, among the older adults in the study from Belo Horizonte-MG, some presented cardiovascular problems, reaffirming the influence of the presence of chronic diseases in the perception of health conditions.

A relevant point is that, while the literature has been discussing the correlations between subjective complaints of memory and health conditions<sup>(11)</sup>, this study measured the performance of memory through a neuropsychological test. This gives greater reliability to the findings, since evaluation of this function considering subjectivity becomes abstract and vulnerable to the psychic and cognitive status of the individual interviewed<sup>(26)</sup>, as mentioned previously.

### Study limitations

This study presents some limitations regarding the comparison and discussion of the findings, since few national studies make a systematic evaluation of memory performance in the older adults' population and correlate it with health conditions. Another limitation is the fact that the absence of correlation between the health perception variable and the results of the neuropsychological test can be due to the reduced number of participants in the study, a fact that demonstrates the need for new studies on this subject.

### Contributions to the area of nursing, health or public policy

The contribution of the present study, as far as the authors were able to show, is due to the fact that it presents results little explored in Brazil. The early identification of memory alterations and their association with health conditions can help prevent cognitive decline, directly influencing the autonomy and independence of the elderly, factors that determine the quality of life. In this way, the nursing professional can use in his professional practice instruments that allow the systematization of evaluations, including neuropsychological tests.

Based on the knowledge of the variables correlated with memory performance and its possible changes, the health professional, in particular the nurse, can identify problems and direct the planning of care to older adults, as well as the monitoring, considering that cognitive loss occurs gradually in this population.

## CONCLUSION

This study evidence that the participating population presented a high score in the MMSE and a satisfactory health perception, despite reporting one or more health problems. Regarding the neuropsychological test that evaluated the memory

performance, there was no correlation of the variables of this test with the health perception.

## FUNDING

Project funded by the *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior. Edital 071/2013 – Programa Nacional de Cooperação Acadêmica 2013.*

## ACKNOWLEDGMENT

To the *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior*, master's degree scholarship.

## REFERÊNCIAS

1. Chaimowicz F. Saúde do Idoso [Internet]. Belo Horizonte: NESCOM UFMG; 2013 [cited 2016 Jun 09]. 167p. Available from: <https://www.nescon.medicina.ufmg.br/biblioteca/imagem/3836.pdf>
2. Moraes EN. Atenção à saúde do idoso: aspectos conceituais [Internet]. Brasília: Organização Pan-Americana da Saúde; 2012 [cited 2016 Jun 12]. Available from: <http://pesquisa.bvsalud.org/porta1/resource/pt/lil-713328>
3. World Health Organization. NIA/NIH/US Department of Health and Human Services. Global aging and health [Internet]. WHO; 2011 [cited 2016 Jun 10]. 32p. Available from: [http://www.who.int/ageing/publications/global\\_health.pdf](http://www.who.int/ageing/publications/global_health.pdf)
4. Brasil. Instituto Brasileiro de Geografia e Estatística-IBGE. Censo demográfico: resultado final: total de pessoas de 0 a 14 anos, 15 a 29 anos, 55 anos ou mais e 60 anos ou mais [Internet]. 2010 [cited 2016 Aug 14]. Available from: <http://www.ibge.gov.br/apps/snig/v1/?loc=0&cat=-1,-2,-3,77,128&ind=4707>
5. Mendes EV. Redes de atenção à saúde [Internet]. Brasília: Organização Pan-Americana de Saúde; 2011 [cited 2016 Jul 02]. Available from: [http://www.saude.sp.gov.br/resources/ses/perfil/gestor/documentos-de-planejamento-em-saude/elaboracao-do-plano-estadual-de-saude-2010-2015/textos-de-apoios/redes\\_de\\_atencao\\_mendes\\_2.pdf](http://www.saude.sp.gov.br/resources/ses/perfil/gestor/documentos-de-planejamento-em-saude/elaboracao-do-plano-estadual-de-saude-2010-2015/textos-de-apoios/redes_de_atencao_mendes_2.pdf)
6. Vagetti GC, Barboza Fo VC, Moreira NB, Oliveira V, Mazzardo O, Campos W. Condições de saúde e variáveis sociodemográficas associadas à qualidade de vida em idosas de um programa de atividade física de Curitiba, Paraná, Sul do Brasil. Cad Saúde Pública [Internet]. 2013 [cited 2016 Jul 19];29(5):955-69. Available from: <http://www.scielo.br/pdf/csp/v29n5/13.pdf>
7. Benedetti TRB, Mazo GZ, Borges LJ. Condições de saúde e nível de atividade física em idosos participantes e não participantes de grupos de convivência de Florianópolis. Ciênc Saúde Colet [Internet]. 2012 [cited 2016 Jul 19];17(8):2087-93. Available from: <http://www.scielo.br/pdf/csc/v17n8/19.pdf>
8. Bharer L. Cognitive plasticity in older adults: effects of cognitive training and physical exercise. Ann N Y Acad Sci [Internet]. 2015 [cited 2016 May 29];1337:1-6. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/nyas.12682/pdf>
9. Abrisqueta-Gomez J. Memória e envelhecimento cognitivo saudável. In: Malloy-Diniz LF, Fuentes D, Consenza RM, (orgs). Neuropsicologia do Envelhecimento: uma abordagem multidimensional. Porto Alegre: Artmed; 2013. p. 171-96.
10. Morrison JH, Baxter MG. The aging cortical synapses: hallmarks and implication for cognitive decline. Nat Rev Neurosci [Internet]. 2012 [cited 2016 Jul 19];13(4):240-50. Available from: <http://www.nature.com/nrn/journal/v13/n4/pdf/nrn3200.pdf>
11. Lee PL. The relationship between memory complaints, activity and perceived health status. Scand J Psychol [Internet]. 2014 [cited 2016 Jul 17];55(2):136-41. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/sjop.12107/epdf>
12. Freitas SR, Fernandes MH, Coqueiro RS, Reis Júnior WM, Rocha SV, Brito TA. Functional capacity and associated factors in the elderly: a population study. Acta Paul Enferm [Internet]. 2012 [cited 2016 Jun 21];25(6): 933-9. Available from: [http://www.scielo.br/pdf/ape/v25n6/en\\_v25n6a17.pdf](http://www.scielo.br/pdf/ape/v25n6/en_v25n6a17.pdf)
13. Malloy-Diniz LF, Abreu N, Bertola L, Fuentes D, Antunes AM, Paula JJ, et al. O exame neuropsicológico do idoso. In: Malloy-Diniz LF, Fuentes D, Consenza RM, organizadores. Neuropsicologia do Envelhecimento: uma abordagem multidimensional Porto Alegre: Artmed; 2013. p. 243-264.
14. Bertolucci PHF, Brucki SMD, Campacci SR, Juliano Y. O mini-exame do estado mental em uma população geral: impacto da escolaridade. Arq Neuro-Psiquiatr [Internet]. 1994 [cited 2016 Aug 29];52(1):1-7. Available from: <http://www.scielo.br/pdf/anp/v52n1/01.pdf>
15. Yesavage JA, Sheikh JL. 9/Geriatric Depression Scale (GDS): recent evidence and development of a shorter version. Clin Gerontol [Internet]. 1986 [cited 2016 Aug 29];5(1-2):165-73. Available from: [http://www.tandfonline.com/doi/pdf/10.1300/J018v05n01\\_09?needAccess=true](http://www.tandfonline.com/doi/pdf/10.1300/J018v05n01_09?needAccess=true)

16. Almeida OP. Mini Exame do Estado Mental e o diagnóstico de demência no Brasil. *Arq Neuropsiquiatr* [Internet]. 1998 [cited 2016 Aug 22];56(3-B):605-12. Available from: <http://www.scielo.br/pdf/anp/v56n3B/1774.pdf>
17. Malloy-Diniz LF, Lasmar VAP, Gazinelli LSR, Fuentes D, Salgado JV. The Rey Auditory-Verbal Learning Test: applicability for the Brazilian elderly population. *Rev Bras Psiquiatr* [Internet]. 2007 [cited 2016 Sep 03];29(4):324-9. Available from: <http://www.scielo.br/pdf/rbp/v29n4/a06v29n4.pdf>
18. Coelho FGM, Vital TM, Novais IP, Costa GA, Stella F, Santos-Galduroz RF. Desempenho cognitivo em diferentes níveis de escolaridade de adultos e idosos ativos. *Rev Bras Geriatr Gerontol* [Internet]. 2012 [cited 2016 Sep 07];15(1):7-15. Available from: <http://www.scielo.br/pdf/rbgg/v15n1/02.pdf>
19. Oliveira RS, Trezza BM, Busse AL, Jacob-Filho W. Efeito de aprendizagem de testes cognitivos computadorizados em idosos. *Einstein* [Internet]. 2014 [cited 2016 Sep 18];12(2):149-53. Available from: [http://www.scielo.br/pdf/eins/v12n2/pt\\_1679-4508-eins-12-2-0149.pdf](http://www.scielo.br/pdf/eins/v12n2/pt_1679-4508-eins-12-2-0149.pdf)
20. Ichihara-Takeda S, Taqueda K, Ikeda N, Matsuyama K, Funahashi S. Neuropsychological assessment of a new computerized cognitive task that was developed to train several cognitive functions simultaneously. *Front Psychol* [Internet]. 2016 [cited 2016 Aug 13];7:497. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4828453/pdf/fpsyg-07-00497.pdf>
21. Correia R, Nieto A, Ferreira D, Sabucedo M, Barroso J. Fund of information is more strongly associated with neuropsychological function than education in older spanish adults. *Arch Clin Neuropsychol* [Internet]. 2015 [cited 2016 Sep 10];30(4):310-21. Available from: <http://acn.oxfordjournals.org/content/30/4/310.full.pdf+html>
22. Alzheimer Disease International. World Alzheimer Report 2014: dementia and risk reduction: an analysis of prospective and modifiable factors [Internet]. London: Alzheimer Disease International; 2014 [cited 2016 Jul 05]. Available from: <https://www.alz.co.uk/research/WorldAlzheimerReport2014.pdf>
23. Ribeiro PCC, Oliveira BHD, Cupertino APFB, Neri AL, Yassuda MS. Desempenho de idosos na Bateria Cognitiva Cerad: relações com variáveis sociodemográficas e saúde percebida. *Psicol-Reflex Crit* [Internet]. 2010 [cited 2016 May 14];23(1):102-9. Available from: <http://www.scielo.br/pdf/prc/v23n1/a13v23n1.pdf>
24. Melo DM, Barbosa AJG. O uso do Mini-Exame do Estado Mental em pesquisas com idosos no Brasil: uma revisão sistemática. *Ciênc Saúde Colet* [Internet]. 2015 [cited 2016 Aug 24];20(12):3865-76. Available from: <http://www.scielo.br/pdf/csc/v20n12/1413-8123-csc-20-12-3865.pdf>
25. Freitas DHM, Campos FCA, Linhares LQ, Santos CR, Ferreira CB, Diniz BS, et al. Autopercepção da saúde e desempenho cognitivo em idosos residentes na comunidade. *Rev Psiq Clin* [Internet]. 2010 [cited 2016 Aug 19];37(1):49-52. Available from: <http://www.scielo.br/pdf/rpc/v37n1/a07v37n1.pdf>
26. Silva RJS, Smith-Menezes A, Tribess S, Rómo-Perez V, Virtuoso Júnior JS. Prevalência e fatores associados à percepção negativa da saúde em pessoas idosas no Brasil. *Rev Bras Epidemiol* [Internet]. 2012 [cited 2016 Aug 15];15(1):49-62. Available from: <http://www.scielo.br/pdf/rbepid/v15n1/05.pdf>
27. DeSalvo KB, Muntner P. Discordance between physician and patient self-rated health and all-Cause mortality. *Ochsner J* [Internet]. 2011 [cited 2016 Aug 22];11:232-40. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3179188/pdf/i1524-5012-11-3-232.pdf>
28. Borges AM, Santos G, Kummer JA, Fior L, Molim VD, Wibelinger LM. Autopercepção de saúde em idosos residentes em um município do interior do Rio Grande do Sul. *Rev Bras Geriatr Gerontol* [Internet]. 2014 [cited 2016 Sep 03];17(1):79-86. Available from: <http://www.scielo.br/pdf/rbgg/v17n1/1809-9823-rbgg-17-01-00079.pdf>
29. Agostinho MR, Oliveira MC, Pinto MEB, Balardin GU, Harzheim E. Autopercepção da saúde entre usuários da Atenção Primária em Porto Alegre, RS. *Rev Bras Med Fam Com* [Internet]. 2010 [cited 2016 Aug 22];5(17):9-15. Available from: <https://rbmfc.org.br/rbmfc/article/view/175/128>
30. Magalhães SS, Hamdan AC. The Rey Auditory Verbal Learning Test: normative data for the Brazilian population and analysis of the influence of demographic variables. *Psychol Neurosci* [Internet]. 2010 [cited 2016 Sep 10];3(1):85-91. Available from: <http://www.scielo.br/pdf/pn/v3n1/v3n1a11.pdf>