



Motricidade

ISSN: 1646-107X

director@revistamotricidade.com

Edições Desafio Singular

Portugal

Borba Neves, Eduardo  
Correlations between the simulated military tasks performance and physical fitness tests  
at high altitude  
Motricidade, vol. 13, núm. 2, 2017, pp. 12-17  
Edições Desafio Singular  
Vila Real, Portugal

Available in: <http://www.redalyc.org/articulo.oa?id=273053385003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Correlations between the simulated military tasks performance and physical fitness tests at high altitude

Eduardo Borba Neves<sup>1\*</sup>

ORIGINAL ARTICLE

## ABSTRACT

The aim of this study was to investigate the Correlations between the Simulated Military Tasks Performance and Physical Fitness Tests at high altitude. This research is part of a project to modernize the physical fitness test of the Colombian Army. Data collection was performed at the 13<sup>th</sup> Battalion of Instruction and Training, located 30km south of Bogota D.C., with a temperature range from 1°C to 23°C during the study period, and at 3100m above sea level. The sample was composed by 60 volunteers from three different platoons. The volunteers start the data collection protocol after 2 weeks of acclimation at this altitude. The main results were the identification of a high positive correlation between the 3 Assault wall in succession and the Simulated Military Tasks performance ( $r = 0.764$ ,  $p < 0.001$ ), and a moderate negative correlation between pull-ups and the Simulated Military Tasks performance ( $r = -0.535$ ,  $p < 0.001$ ). It can be recommended the use of the 20-consecutive overtaking of the 3 Assault wall in succession as a good way to estimate the performance in operational tasks which involve: assault walls, network of wires, military Climbing Nets, Tarzan jump among others, at high altitude.

**Keywords:** Simulated Military Tasks, Obstacle course, Physical Fitness Tests, Moderate Altitude.

## INTRODUCTION

The readiness and physical resilience of each soldier are essential for the good operational performance of a troop. Thus, an adequate and periodic assessment is a prerequisite for maintenance and improvement of a military readiness (Rohde, Sievert, Rüther, Witzki, & Leyk, 2015). Nowadays most military physical fitness tests (MPFT) still not adequate to reproduce and monitor physical requirements of military tasks (Jonas, O'Connor, Deuster, & Macedonia, 2010).

On the other hand, some nations start to developed additional mission-related test methods to assess operational readiness, such as: the Combat Fitness Test of the U.S. Marine Corps (United States of America, 2015) and FORCE Fitness Profile of the Canadian Armed Forces (Gagnon, Spivock, Reilly, Mattie, & Stockbrugger, 2015). This can be considered a change in MPFT philosophy, but these tests have to be feasible to assess a large number of military

in a few time (Wyss, Marti, Rossi, Kohler, & Mader, 2007).

However, some missions must be done in adverse conditions, as in high altitude (Derby & Weber, 2010; United States of America, 2002), and in this case, the hypoxia can alter the performance of troops in some activities, because it decrease the maximal consume of oxygen in 17% (Consolazio, Nelson, Matoush, & Hansen, 1966). Moreover, military tasks in cold weather at moderate altitude are accompanied by increases in several indicators of oxidative stress such as: breath pentane, oxygen radical absorption capacity, and lipid peroxides (LPO) (Chao, Askew, Roberts, Wood, & Perkins, 1999). Thus, it makes sense to know the response of troop to this kind of stimulus, nevertheless how to do it in places without large military areas or buildings which allow simulating military tasks? One solution can be the use of physical fitness tests which are strongly related with military tasks performance (Gagnon, 2016). In this sense, the aim of this

Manuscript received at March 15<sup>th</sup> 2017; Accepted at June 5<sup>th</sup> 2017

<sup>1</sup> Brazilian Army Research Institute of Physical Fitness, Rio de Janeiro, Brazil

\* Corresponding author: Av João Luis Alves s/nº Fortaleza de São João - Urca, CEP: 22291-090, Rio de Janeiro, RJ, Brazil. E-mail: neveseb@gmail.com

study was to investigate the Correlations between the Simulated Military Tasks Performance and Physical Fitness Tests at high altitude.

### METHOD

This research is part of a project to modernize the physical fitness test of the Colombian Army. Data collection was performed at the 13<sup>th</sup> Battalion of Instruction and Training, located 30km south of Bogota D.C., with a temperature range from 1°C to 23°C during the study period, and at 3100m above sea level.

The sample was composed by 60 volunteers from three different platoons. The volunteers start the data collection protocol after 2 weeks of

acclimation at this altitude. During these 2 weeks they have had different kind of operational and physical instructions. All volunteers signed written informed consent, and all procedures were carried out in accordance with the Declaration of Helsinki and national guidelines.

First, it was established the Simulated Military Tasks to serve as a gold standard, by the Colombian Army Direction of Training and Instruction. After, it was done a selection of the physical fitness test which could be done in small military areas, safe and easy applicable; and requires little material, time, and few personnel. The others phases of this study are presented in the Figure 1.




Figure 1. Study design

Three days before start data collection, volunteers were instructed to avoid strenuous physical activities for at least 24 hours prior to the

evaluation (Rosa et al., 2017). Data collection start on Monday (Day 1) and was carried out according to the sequence presented in Figure 2.




Figure 2: Data collection protocol

During anthropometric measurements, body mass was measured in kilograms (kg) using a Filizola scale with an accuracy of 100 g and a range of 150 kg. Height was measured in centimetres (cm), using the Frankfurt plane as a reference. From these measurements, it was obtained Body Mass Index (BMI) values

(weight/height<sup>2</sup>) (Neves, 2008). Abdomen circumference was measured on the umbilicus using a flexible steel tape and recorded in millimetres. The fat percentage (Fat%) was calculated by the following equation proposed by Salem (2008).

$$\text{Fat\%} = 0.061 \times \text{Age} + 16.002 \times (\text{AC})^{0.5} - 5.056 \times (\text{BM})^{0.5} - 91.222$$

Where: AC = Abdominal circumference, and BM = Body mass.




Figure 3: Simulated Military Tasks (Obstacle course from Colombian Army) used as gold standard in this study

Table 1.

*Physical fitness tests selected to be performed by the volunteers of this study*

| Test                         | Used by  | Observations  |
|------------------------------|--|---|
| Running                      | American College of Sports Medicine (ACSM, 2013)   | Distance of 3000m   |
| Abdominal                    | American College of Sports Medicine (ACSM, 2013)   | Abdominal 30°   |
| Push-ups                     | American College of Sports Medicine (ACSM, 2013)   | -   |
| Horizontal Long Jump         | Norway Army (Gagnon, 2016)   | The best of 3 attempt   |
| Sandbag Lift                 | Canadian Army (Gagnon et al., 2015)  | Time of 30 consecutive lifts of a 20 kg sandbag from the floor above a height of 1.0 m. |
| Pull-ups                     | Brazilian Army (Exército Brasileiro, 2015)   | -   |
| 3 Assault wall in succession | This is the last obstacle from the Obstacle run of Military Pentathlon (Exército Brasileiro, 2015) | Time of 20 consecutive overtaking   |




Figure 4: 3 Assault Walls in Succession (measures in meters)

The Simulated Military Tasks were selected from an obstacle course which already existed in the 13th Battalion of Instruction and Training. Nine from twelve obstacles were selected to be performed as presented in the Figure 3.

The physical fitness tests selected were presented in the Table 1. The last test, 3 Assault wall in succession was illustrated in Figure 4.

The data analysis started with the normality verification using the Shapiro-Wilk test. Once some variables show a nonparametric distribution, descriptive data are presented as minimum, maximum, mean, median, standard deviation, interquartile range. The correlation analysis was performed with Pearson correlation coefficient and Spearman Rank Correlation,

according to the type of distribution presented by each variable (Mukaka, 2012). The analyses were performed using the software Statistical Package for Social Sciences version 21.0. The significance level was set in 0.05.

## RESULTS

The Table 2 presents descriptive data from 60 Colombian militaries from three different platoons which did the simulated military tasks and physical fitness tests selected for this study. The Table 3 presents the correlation coefficients between the simulated military tasks performance (Obstacle course illustrated in the Figure 3) and the physical fitness tests selected.

Table 2.

*Descriptive data of 60 Colombian militaries from three different platoons which did the Simulated Military Tasks (Obstacle course) and Physical Fitness Tests, Colombia, 2017.*

|  | Shapiro-Wilk | Mínimo | Máximo | Average | Standard Deviation | Median | Interquartile range |
|--|--------------|--------|--------|---------|--------------------|--------|---------------------|
| Age (years)                              | 0.015        | 23.00  | 41.00  | 32.13   | 5.01               | 32.00  | 9,00                |
| Body mass (Kg)                           | 0.097        | 56.00  | 84.00  | 70.09   | 7.53               | 70.00  | 11.00               |
| Height (cm)                              | 0.706        | 154.00 | 185.00 | 169.33  | 6.78               | 170.00 | 9.50                |
| Body Mass Index                          | 0.205        | 20.00  | 29.00  | 24.42   | 2.14               | 24.00  | 2.75                |
| Abdominal Circunference (cm)             | 0.505        | 69.00  | 101.00 | 85.93   | 7.03               | 85.50  | 11.50               |
| Fat Percentage                           | 0.438        | 3.94   | 28.02  | 16.95   | 5.19               | 16.07  | 7.04                |
| Running (s)                              | 0.181        | 11.32  | 17.32  | 889.20  | 86.40              | 867.00 | 116.00              |
| Abdominal 30° (rep)                      | 0.382        | 13.00  | 70.00  | 38.02   | 13.07              | 37.50  | 18.00               |
| Push-ups (rep)                           | 0.235        | 9.00   | 60.00  | 32.00   | 9.79               | 31.00  | 12.50               |
| Horizontal Long Jump (cm)                | 0.810        | 162.00 | 272.00 | 205.98  | 22.31              | 207.00 | 28.25               |
| Sandbag Lift (s)                         | 0.126        | 51.00  | 86.00  | 64.32   | 8.24               | 64.00  | 9.75                |
| Simulated Military Tasks Performance (s) | 0.194        | 170.00 | 420.00 | 271.92  | 59.64              | 259.50 | 88.25               |
| Pull-ups (rep)                           | 0.004        | 1.00   | 20.00  | 6.49    | 4.33               | 5.00   | 7.00                |
| 3 Assault wall in succession (s)         | 0.320        | 159.00 | 298.00 | 217.94  | 32.45              | 216.50 | 41.75               |

Table 3.

*Correlation coefficients between the Simulated Military Tasks (Obstacle course) and the Physical Fitness Tests (N=60), Colombia, 2017.*

| Correlations                     | Simulated Military Tasks Performance (s) | p value |
|----------------------------------|--|---------|
| Running (s)                      | $r = 0.273^*$                            | 0.036   |
| Abdominal 30° (rep)              | $r = -0.189$                             | 0.151   |
| Push-ups (rep)                   | $r = -0.416^*$                           | 0.001   |
| Horizontal Long Jump (cm)        | $r = -0.391^*$                           | 0.002   |
| Sandbag Lift (s)                 | $r = 0.146$                              | 0.277   |
| Pull-ups (rep)                   | $\rho = -0.535^*$                        | <0.001  |
| 3 Assault wall in succession (s) | $r = 0.764^*$                            | <0.001  |

$\rho$  = Spearman Rank Correlation,  $r$  = Pearson correlation coefficient, \* Significant correlations.

Regarding to the anthropometric variables, the Pearson correlations coefficients with the simulated military tasks performance were: 0.279 with body mass ( $p=0.039$ ), 0.079 with height ( $p=0.565$ ), 0.271 Body Mass Index ( $p=0.046$ ), 0.527 with abdominal circumference ( $p<0.001$ ), and 0.505 with fat percentage ( $p<0.001$ ).

### DISCUSSION

The objective of this study was to investigate the correlations between the Simulated Military Tasks Performance and Commons Physical Fitness Tests at high altitude. The main results were the identification of a high positive correlation (Mukaka, 2012) between the 3 Assault wall in succession and the Simulated Military Tasks performance ( $r = 0.764$ ,  $p<0.001$ ), and a moderate negative correlation between pull-ups and the Simulated Military Tasks performance ( $r = -0.535$ ,  $p<0.001$ ).

These results agree with those found by Pihlainen, Santtila, Häkkinen, and Kyröläinen (2017) when studied soldiers from Finnish Defense Forces, and they reported a moderated correlation between the countermovement jump performed with the combat load and a military simulated test (MST) time ( $\rho = -0.659$ ,  $p<0.001$ ). They also find significant correlations among the MST time with Pull-ups ( $\rho = -0.435$ ,  $p<0.001$ ), and fat percentage ( $\rho = 0.534$ ,  $p<0.001$ ). In the present study, it was found a similar correlation coefficient among the simulated military tasks with Pull-ups ( $\rho = -0.535$ ,  $p<0.001$ ), and the fat percentage ( $r = 0.505$ ,  $p<0.001$ ).

The high positive correlation coefficient found between the 20 consecutive overtaking of the 3 Assault wall in succession and the simulated

military tasks reinforce the idea reported by Pihlainen et al. (2017) that the endurance capacity and the muscle power of the lower extremities are crucial performance components in anaerobic combat situations, even at high altitude (3100m above sea level).

However, the correlation coefficients depend of the operational tasks established as reference, i.e. those defined by Canadian Armed Forces (escape to cover, picking, digging, stretcher-carry and sandbag fortifications, among others...), in a project named FORCE program (Gagnon, 2016), in which they did similar analyses between operational tasks and physical fitness tests. The correlation coefficient between the stretcher-carry task and sandbag lift was 0.737, and between push-ups and the sandbag fortifications task was 0.599. In the present study, these physical fitness tests (sandbag lift and push-ups) did not show good correlations coefficients with the simulated military tasks composed by assault walls, network of wires, military Climbing Nets, Tarzan jump among others illustrated in Figure 3.

Most of the studies were done in low altitude. This study brings some results of correlations between simulated military tasks performance and physical fitness tests at 3100m above sea level. The sample size and the guarantee of the same level of subject's motivation in all days of this study protocol can be cited as a limitation of this study. Future procedures involve increase the sample size and the test of reliability of the physical fitness tests performed.

### PRACTICAL APPLICATIONS

For while, it can be recommended the use of the 20-consecutive overtaking of the 3 Assault

wall in succession as a good way to estimate the performance in operational tasks which involve: assault walls, network of wires, military Climbing Nets, Tarzan jump among others illustrated in Figure 3, at high altitude (3100m above sea level). This test can be done in small military areas, it is safe and easy applicable; and requires little material, time, and few personnel.

#### Acknowledgments:

We would to say thanks to the Chief of Colombian Army Department of Military Education, to the Director of Training and Instruction, to the Commander of 13th Battalion of Instruction and Training, and to all volunteers of this study

#### Conflict of interest:

Nothing to declare

#### Funding:

Nothing to declare

#### REFERENCES

- ACSM, American College of Sports Medicine. (2013). *ACSM's guidelines for exercise testing and prescription*: Lippincott Williams & Wilkins.
- Chao, Wei-Hsun, Askew, Eldon W., Roberts, Donald E., Wood, Steven M., & Perkins, James B. (1999). Oxidative Stress in Humans during Work at Moderate Altitude. *The Journal of Nutrition*, 129(11), 2009-2012.
- Consolazio, C. F., Nelson, R. A., Matoush, L. R., & Hansen, J. E. (1966). Energy metabolism at high altitude (3,475 m). *Journal of Applied Physiology*, 21(6), 1732-1740.
- Derby, Richard, & Weber, Kevin de. (2010). The athlete and high altitude. *Current sports medicine reports*, 9(2), 79-85.
- Exército Brasileiro, Estado Maior do Exército Brasileiro. (2015). *EB20-MC-10.350 Treinamento Físico Militar* (4ª ed.). Brasília: EGGCF.
- Gagnon, Patrick. (2016, 22 Set). *FORCEcombat – Adapting the Annual FORCE Evaluation for the Specific Demands of the Canadian Army's Combat Missions*. Paper presented at the Nordic Military Sports Leader Meeting and Symposium, Tuusula, Finland.
- Gagnon, Patrick, Spivock, Michael, Reilly, Tara, Mattie, Paige, & Stockbrugger, Barry. (2015). The FORCE Fitness Profile—Adding a Measure of Health-Related Fitness to the Canadian Armed Forces Operational Fitness Evaluation. *The Journal of Strength & Conditioning Research*, 29, S192-S198.
- Jonas, Wayne B, O'Connor, Francis, Deuster, Patricia, & Macedonia, Christian. (2010). Total Force Fitness for the 21st Century A New Paradigm. *Military Medicine*, 175( August Supplement ), 14-20.
- Mukaka, MM. (2012). A guide to appropriate use of correlation coefficient in medical research. *Malawi Medical Journal*, 24(3), 69-71.
- Neves, Eduardo Borba. (2008). Prevalência de sobrepeso e obesidade em militares do exército brasileiro: associação com a hipertensão arterial. *Ciê Saú de Coletiva*, 13(5), 1661-1668.
- Pihlainen, K, Santtila, M, Häkkinen, K, & Kyröläinen, H. (2017). Associations of physical fitness and body composition characteristics with simulated military task performance. *The Journal of Strength & Conditioning Research*, Epub ahead of print. doi: 10.1519/JSC.0000000000001921
- Rohde, Ulrich, Sievert, Alexander, Rütther, Thomas, Witzki, Alexander, & Leyk, Dieter. (2015). Concept for a predeployment assessment of basic military fitness in the German armed forces. *The Journal of Strength & Conditioning Research*, 29, S211-S215.
- Rosa, Claudio, Vilaça-Alves, José, Neves, Eduardo Borba, Saavedra, Francisco José Félix, Reckziegel, Miriam Beatris, Pohl, Hildegard Hedwig, . . . Reis, Victor Machado. (2017). The effect of weekly low frequency exercise on body composition and blood pressure of elderly women. *Archivos de Medicina del Deporte*, 34(1), 9-14.
- Salem, Marcelo. (2008). *Desenvolvimento e validação de equações e índices para a determinação da gordura corporal relativa, em militares brasileiros, a partir de medidas antropométricas*. (Tese de Doutorado), FIOCRUZ, Rio de Janeiro.
- United States of America. (2002). *Field Manual 3-97.61 Military Mountaineering*. Washington DC: Department of the Army (US).
- United States of America. (2015). *Marine Corps Order 6100.13 W/CH 2: Marine Corps Physical Fitness Program*. Washington DC: Headquarters United States Marine Corps.
- Wyss, Thomas, Marti, Bernard, Rossi, Sandro, Kohler, Urs, & Mader, Urs. (2007). Assembling and verification of a fitness test battery for the recruitment of the Swiss Army and nation-wide use. *Schweizerische Zeitschrift für Sportmedizin und Sporttraumatologie*, 55(4), 126-131.

