



Cardiocre

ISSN: 1889-898X

cardiocre@elsevier.com

Sociedad Andaluza de Cardiología
España

Pereira Silva, Ricardo; Pimentel Landim de Almeida, Camila; Bezerra da Silva Junior, Geraldo; Lucas Costa, Pedro; Cunha Costa, Pedro Sérgio; Gondim Rocha, Marilena
Electrocardiogram reports in a cardiology tests clinic
Cardiocre, vol. 50, núm. 1, enero-marzo, 2015, pp. 22-26
Sociedad Andaluza de Cardiología
Barcelona, España

Available in: <http://www.redalyc.org/articulo.oa?id=277041126006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Cardiocre

www.elsevier.es/cardiocre



Original article

Electrocardiogram reports in a cardiology tests clinic



Ricardo Pereira Silva^{a,b,*}, Camila Pimentel Landim de Almeida^a,
Geraldo Bezerra da Silva Junior^c, Pedro Lucas Costa^a, Pedro Sérgio Cunha Costa^a,
Marilena Gondim Rocha^a

^a Unicordis, Hospital São Mateus, Fortaleza, Ceará, Brazil

^b Department of Internal Medicine, School of Medicine, Federal University of Ceará, Fortaleza, Ceará, Brazil

^c School of Medicine, Post-Graduation Program in Collective Health, Health Sciences Center, University of Fortaleza, Fortaleza, Ceará, Brazil

ARTICLE INFO

Article history:

Received 11 May 2014

Accepted 4 September 2014

Available online 17 December 2014

Keywords:

Electrocardiogram

ECG

Cardiology tests

Ventricular repolarization

Coronary disease

ABSTRACT

Background: The electrocardiogram (ECG) is the oldest and the most frequently used test in cardiology. The aim of this study is to verify, in different patients' groups, which are the most frequent ECG reports in a cardiology tests clinic.

Methods: This is a retrospective evaluation of all ECGs performed in a cardiology clinic in Fortaleza, Ceará, Brazil. The main indications and results were evaluated.

Results: A total of 1150 ECGs were performed; 63% were normal, and the most frequent abnormalities were altered ventricular repolarization (AVR) and intraventricular conduction disturbance (16% of cases). In the preoperative groups, the ECG report was normal in 67% of cases, and the most frequent abnormality was intraventricular conduction disturbance (32.8%). In the clinical evaluation group, the report was normal in 60%, and the most frequent abnormality was also intraventricular conduction disturbance (18.4%). In the hypertension group, the report was normal in 56%, and the most commonly found abnormality was altered ventricular repolarization (22.5%). In the group with diagnosis or suspect coronary disease, the ECG was normal in 43%, and the most frequent abnormality was altered ventricular repolarization (34%). In the group with tachycardia, the ECG was normal in 65% of cases, and the most frequent abnormality was altered ventricular repolarization (11.7%). In the group of patients with history of syncope, the ECG was normal in all cases.

Conclusion: The ECG was normal in the majority of cases and the most frequent abnormality found was altered ventricular repolarization, independently of the indication for the test performance.

© 2014 SAC. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail addresses: ricardop@ufc.br, geraldobezerrajr@yahoo.com.br (R.P. Silva).
<http://dx.doi.org/10.1016/j.carcor.2014.09.002>

1889-898X/© 2014 SAC. Published by Elsevier España, S.L.U. All rights reserved.

Informes de electrocardiogramas en una clínica de exploraciones cardiológicas

R E S U M E N

Palabras clave:

Electrocardiograma
ECG
Exploraciones de cardiología
Repolarización ventricular
Enfermedad coronaria

Antecedentes: El electrocardiograma (ECG) es la exploración cardiológica más antigua y más comúnmente utilizada. El objetivo de este estudio es verificar, en diferentes grupos de pacientes, cuáles son los informes de ECG más frecuentes en una clínica de exploraciones cardiológicas.

Métodos: Se ha llevado a cabo una evaluación retrospectiva de todos los ECG realizados en una clínica de cardiología de Fortaleza, Ceará (Brasil). Se evaluaron las principales indicaciones y los resultados.

Resultados: Se realizó un total de 1150 ECG; un 63% fueron normales, y las anomalías más frecuentes fueron las siguientes: alteración de la repolarización ventricular (ARV) y trastorno de la conducción intraventricular (16% de los casos). En los grupos preoperatorios, el informe del ECG fue normal en el 67% de los casos, y la anomalía más frecuente fue el trastorno de la conducción intraventricular (32,8%). En el grupo de evaluación clínica, el informe fue normal en el 60%, y la anomalía más frecuente fue también el trastorno de la conducción intraventricular (18,4%). En el grupo de hipertensión, el informe fue normal en el 56%, y la anomalía más frecuente observada fue la alteración de la repolarización ventricular (22,5%). En el grupo con diagnóstico o sospecha de enfermedad coronaria, el ECG fue normal en el 43%, y la anomalía más frecuente fue la alteración de la repolarización ventricular (34%). En el grupo con taquicardia, el ECG fue normal en el 65% de los casos, y la anomalía más frecuente fue la alteración de la repolarización ventricular (11,7%). En el grupo de pacientes con antecedentes de síncope, el ECG fue normal en todos los casos.

Conclusión: El ECG fue normal en la mayoría de los casos y la anomalía observada con mayor frecuencia fue la alteración de la repolarización ventricular, con independencia de cuál fuera la indicación para realizar la exploración.

© 2014 SAC. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Among all tests used in cardiology, the electrocardiogram (ECG) is the oldest and the most frequently used. According to American Guidelines on ECG, this test should be done even for patients without cardiopathy suspect, included in the following conditions (class I): (1) for clinical evaluation of individuals older than 40 years; (2) to evaluate patients before the administration of cardiotoxic drugs; (3) to evaluate individuals before ergometric test; (4) to evaluate individuals at any age who have jobs that require high cardiovascular performance (e.g. astronauts, fireman, policeman) or whose performance is linked to public safety (e.g. pilots, bus drivers, air traffic controllers). There is no consensus (class II) regarding the need of ECG in the evaluation of competitive athletes. There is consensus that ECG should not be done (class III) in the routine evaluation of individuals younger than 40 years without risk factors.¹⁻⁴

As preoperative evaluation, the ECG should be performed (class I) in patients older than 40 years and in heart transplant donors or cardiopulmonary transplant receptors. There is no consensus (class II) regarding the ECG performance in preoperative evaluation in patients aged 30-40 years. There is consensus that ECG should not be done (class III) in patients younger than 30 years in the preoperative period, without risk factor for coronary disease.¹

The aim of this study is to verify, in different patients' groups, which are the most frequent ECG reports in a cardiology tests clinic.

Materials and methods

In the period between 1st August and 31st October 2012 a total of 1150 ECGs were performed in the Unicordis, a cardiology tests clinic in Fortaleza, Ceará, Brazil. The study was retrospective and all patients in the period were included in the study. The population was of ambulatory patients. The performance and interpretation of ECGs was done according to recent guidelines, published by the Brazilian Society of cardiology.⁵ Patients' age distribution is summarized in Fig. 1. The most prevalent age was 30-49 years. The majority of patients (61%) were female. The indications for ECG performance were preoperative evaluation (45%), clinical evaluation (30%), hypertension (13%), coronary insufficiency (4%), tachycardia/palpitation (4%), examination for job selection process (2%), mitral valve prolapse (1%) and syncope/pre-syncope (1%) (Table 1). All the ECGs were performed by a Cardiologist.

Results

A total of 1150 tests were analyzed; 444 were male (38.7%) and 706 were female (61.3%), with a mean age of 49.6 ± 19.2




Fig. 1 – Age distribution of patients undergoing ECG. *Mean age = 49.84 years.

years. Age distribution according to ECG indication was: preoperative evaluation – 48.4 ± 17.7 , clinical evaluation – 51.1 ± 20.2 , hypertension – 57.3 ± 19.0 , coronary insufficiency – 52.6 ± 20.5 , tachycardia/palpitation – 40.2 ± 21.1 , examination for job selection process – 42.0 ± 10.4 , mitral valve prolapse – 31.8 ± 11.7 and syncope/pre-syncope – 44.2 ± 26.1 years.

Considering the total of 1150 tests, 63% were normal, and the most frequent abnormalities were: altered ventricular repolarization (AVR) and intraventricular conductance disturbance, each one in 16% of cases (Table 2). In the preoperative groups, the ECG report was normal in 67% of cases, and the most frequent abnormality was intraventricular conductance disturbance, in 32.8% (Table 3). In the clinical evaluation group, the report was normal in 60%, and the most frequent abnormality was also intraventricular conductance disturbance, in 18.4% (Table 4). In the hypertension group, the report was normal in 56%, and the most commonly found abnormality was altered ventricular repolarization, in 22.5% (Table 5). In the group with diagnosis or suspect coronary disease, the ECG was normal in 43%, and the most frequent abnormality was altered ventricular repolarization, in 34% (Table 6). In the group with tachycardia, the ECG was normal in 65% of cases, and the most frequent abnormality was altered ventricular repolarization, in 11.7% (Table 7). In the group of individuals who performed ECG as a test for job selection, it was normal in 69% of cases, and the most frequent abnormality was altered ventricular repolarization, in 13.8% (Table 8). In the group of patients with mitral valve prolapse, the report was normal in 69.3%, and the

Table 1 – ECG indications.

	n	%
Preoperative evaluation	516	45
Clinical evaluation ^a	350	30
Hypertension	147	13
Coronary disease	44	4
Tachycardia/palpitation	43	4
Test for job selection	29	2
Mitral valve prolapse	13	1
Syncope	8	1

^a ECG performed when a person will begin a new job, and an ECG is required as a pre-requisite as part of health evaluation.

Table 2 – ECG reports.

	n	%
Normal	722	63
Altered ventricular repolarization	184	16
Conductance disturbance ^a	183	16
Extrasystoles	39	3.4
Heart chambers overload ^b	29	2.5
Sinus bradycardia	23	2.0
1st degree atrioventricular block	23	2.0
Inactive zone	22	1.9
Atrial fibrillation	7	0.6

^a Intraventricular conductance disturbances: 62 incomplete right bundle branch block (5.4%), 27 complete right bundle branch block (2.3%), 7 complete left bundle branch block (0.6%), 83 left anterior fascicular block (7.2%).

^b Heart chambers overload: 12 left atrium overloads, 8 left ventricle overload, 6 right ventricle overload, 2 right chambers overload and 1 left chamber overload.

Table 3 – ECG reports in the group of preoperative evaluation.

	n	%
Normal	346	67
Conductance disturbances ^a	66	32.8
Altered ventricular repolarization	71	13.7
Sinus bradycardia	14	2.8
Extrasystoles	8	1.6
Inactive zone	7	1.4
Heart chambers overload ^b	8	1.6
1st degree atrioventricular block	5	1.0
Atrial fibrillation	3	0.6

^a Intraventricular conductance disturbances: 29 incomplete right bundle branch block (5.6%), 6 complete right bundle branch block (21.2%), 1 complete left bundle branch block (0.2%), 30 left anterior fascicular block (5.8%).

^b Heart chambers overload: 3 left atrium overload, 1 left ventricular overload, 4 right ventricular overload.

Table 4 – ECG reports in the group “clinical evaluation”.

	n	%
Normal	210	60
Conductance disturbances ^a	64	18.4
Altered ventricular repolarization	55	15.7
Extrasystoles	22	6.3
Inactive zone	8	2.3
Heart chambers overload ^b	8	2.3
Sinus bradycardia	5	1.4
1st degree atrioventricular block	4	0.85
Atrial fibrillation	2	0.6

^a Intraventricular conductance disturbances: 16 incomplete right bundle branch block (4.6%), 17 complete right bundle branch block (4.9%), 27 left anterior fascicular block (7.7%), 4 complete left bundle branch block (1.2%).

^b Heart chambers overload: 2 left atrium overload, 3 left ventricular overload, 2 right atrium overload + right ventricular overload, 1 left atrium overload + left ventricular overload.

Table 5 – ECG reports in the group with hypertension.

	n	%
Normal	82	56
Altered ventricular repolarization	33	22.5
Conductance disturbances ^a	29	19.4
Heart chambers overload ^b	8	5.4
Inactive zone	3	2.0
Extrasystoles	2	1.3
Sinus bradycardia	1	0.6
1st degree atrioventricular block	0	0
Atrial fibrillation	0	0

^a Intraventricular conductance disturbances: 6 incomplete right bundle branch block (4%), 1 complete right bundle branch block (0.6%), 1 incomplete left bundle branch block (0.6%), 1 complete left bundle branch block (0.6%), 20 left anterior fascicular block (13.6%).

^b Heart chambers overload: 5 left atrium overload, 3 left ventricular overload.

Table 6 – ECG reports in the group with coronary disease.

	n	%
Normal	19	43
Altered ventricular repolarization	15	34
Conductance disturbances ^a	8	9.1
Inactive zones	3	6.8
Atrial fibrillation	2	4.5
Heart chambers overload ^b	2	4.5
Extrasystoles	1	2.3
Sinus bradycardia	1	0.6
1st degree atrioventricular block	0	0

^a Intraventricular conductance disturbances: 4 incomplete right bundle branch block (9%), 1 complete left bundle branch block (2.3%), 3 left anterior fascicular block (6.8%).

^b Heart chambers overload: 1 left atrium overload, 1 right ventricular overload.

most frequent abnormality was intraventricular conductance disturbance, in 30.8% (Table 9). In the group of patients with history of syncope, the ECG was normal in all cases.

Discussion

The present study evaluated the main indication and results of ECGs performed in a reference center of a metropolitan

Table 7 – ECG reports in the group of tachycardia.

	n	%
Normal	28	65
Altered ventricular repolarization	5	11.7
Extrasystoles	5	11.7
Conductance disturbances ^a	4	9.3
Heart chambers overload ^b	1	2.3
Sinus tachycardia	1	2.3
1st degree atrioventricular block	1	2.3
Inactive zone	1	2.3
Atrial fibrillation	0	0

^a Intraventricular conductance disturbances: 4 incomplete right bundle branch block (9.3%).

^b Heart chambers overload: 1 left ventricular overload.

Table 8 – ECG reports in the group performing tests for job selection.

	n	%
Normal	20	69
Altered ventricular repolarization	4	13.8
Conductance disturbances ^a	4	13.3
Extrasystoles	1	3.4
1st degree atrioventricular block	1	3.4
Heart chambers overload	0	0
Sinus bradycardia	0	0
Inactive zone	0	0
Atrial fibrillation	0	0

^a Intraventricular conductance disturbances: 1 incomplete right bundle branch block (3.4%), 1 complete right bundle branch block (3.4%).

area. The main indications were for preoperative evaluation and “clinical evaluation” (when ECG is done when a person will begin a new job, as part of health evaluation). The majority of ECGs were normal, and the most frequently found abnormalities were intraventricular conductance abnormality and altered ventricular repolarization. In a previous study evaluating the risks for an adult patient in non-cardiac elective surgery the ECG was performed due to the presence of cardiovascular comorbidities in 80% of cases, respiratory comorbidities in 42%, advanced age in 53% and as a routine in 10%.⁶ In our cohort, there was no specific cause for ECG performance in the preoperative period.

The altered ventricular repolarization (AVR) was the most frequent abnormality observed in our study in the general sample, in the groups of hypertension, coronary disease, tachycardia and in those who were undergoing the test for job selection.

“Strain” pattern in the ECG was evaluated in 8554 hypertensive patients in the LIFE study, and were “blind” treated with atenolol or losartan. Strain pattern, defined as the presence of ST segment depression with asymmetric and inverted T-wave in the derivations V5 and V6, was present in 971 patients (11%).⁷ In our group of patients with hypertension, AVR was observed with a higher incidence (22%), but we have to point that in these 22% are included ECGs with AVR with strain pattern and unspecific ventricular repolarization abnormalities.

Table 9 – ECG reports in the group with mitral valve prolapse.

	n	%
Normal	9	69.3
Conductance disturbances ^a	4	30.8
Altered ventricular repolarization	1	7.7
Extrasystoles	0	0
Heart chambers overload	0	0
Sinus bradycardia	0	0
1st degree atrioventricular block	0	0
Inactive zone	0	0
Atrial fibrillation	0	0

^a Intraventricular conductance disturbances: 2 incomplete right bundle branch block (15.4%), 1 complete right bundle branch block (7.7%), 1 left anterior fascicular block (7.7%).

Major and minor ECG abnormalities in asymptomatic women were compared with cardiovascular events and mortality in a *post-hoc* analysis in the study "Women's Health Initiative". Minor abnormalities were considered as: 1st and 2nd degree atrioventricular block, prolonged QT-interval, ST-segment minor abnormalities, left atrium overload, and frequent atrial or ventricular extrasystoles. Major ECG abnormalities were: atrial fibrillation, atrial flutter, 2nd degree atrioventricular block, left bundle branch block, right branch block, inactive electrical zone, and left ventricle hypertrophy with strain pattern. The adjusted risk for coronary events in patients who presented minor ECG abnormalities was 1.55, and for patients with major abnormalities it was 3.01.⁸

The "Reykjavic Study" evaluated the epidemiologic, clinical and prognostic characteristics of angina in the non-recognized acute myocardial infarction and detected that 1/3 of infarctions were not recognized.⁹ The finding of electrical inactive zone in our patients was not frequent. In the general sample, it occurred in 1.9% of cases. The sub-group with higher frequency of electrical inactive zone, as expected, was that of patients with suspected or diagnosed coronary disease, with a 6.8% frequency.

The Chicago Electric Company Study also evaluated the association between unspecific ST-segment abnormalities and cardiovascular mortality. A total of 1673 men were selected from the electric company workers, aged between 44 and 55 years, without evidence of coronary disease and without significant ECG abnormalities at the beginning of the study. Among the 1673 men, 173 developed ST-segment unspecific abnormalities during a 5-years follow-up. The authors linked the number of annual ECGs with ST-segment abnormalities, and found a relative risk for death of 2.28, 2.39, 2.30 and 1.6, respectively, for 4, 3, 2 or 1 ECG with this abnormality.¹⁰

The prevalence, prognosis and implications of unspecific ST-segment abnormalities were also studied in the elderly. A total of 233 out of 3224 participants of the "Cardiovascular Health Study" presented ST-segment unspecific abnormalities at the beginning of the study. After 39,518 patients-years of follow-up, the presence of unspecific ST-segment abnormalities was associated with mortality due to coronary disease (relative risk of 1.76), but there was no association with infarction incidence. This fact suggests that ST-segment unspecific abnormalities are particularly associated with risk for death due to primary arrhythmia.¹¹

In our cohort, the presence of atrial or ventricular extrasystoles was observed in 3.4% of cases of the general sample, and its higher frequency occurred in the groups of individuals who performed the test due to tachycardia (11.7%). It is known that the presence of extrasystoles has prognostic value for heart disease, and it has been shown that frequent ventricular extrasystoles after physical exercise works as a death predictor.¹²

A simple ECG can help to detect silent atrial fibrillation (AF) before the first cerebrovascular event, especially in patients with multiple cardiovascular conditions. AF occurred in 7 out of 132 patients in a case series (5.3%). Among these patients, there were four survivors from stroke, two with diabetes and one with hypertension. The prevalence of AF was higher in patients with multiple risk factors for stroke and AF. AF was found in 3% of patients with hypertension without other risk

factors, in 7% of patients with two risk factors and in 11% of patients with hypertension, diabetes and stroke.¹³ In our cohort, AF was present in 0.6% of cases in the general sample. The group with higher frequency was that of individuals with suspect or diagnosis of coronary disease (4.5%). In our group with hypertension, there was no case of AF.

In summary, the ECG was normal in the majority of cases and the most frequent abnormality found was altered ventricular repolarization, independently of the indication for the test performance.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Schlant R, Adolph RJ, DiMarco JP, et al. ACC/AHA Task Force Report. Guidelines for electrocardiography. *JACC*. 1992;19:473-81.
- Moyer VA, on behalf of the US Preventive Services Task Force. Screening for Coronary Heart Disease with electrocardiography: US preventive services task force recommendation statement. *Ann Intern Med*. 2012;157: 512-8.
- Chaitman BR. Should an electrocardiogram be included in routine preparticipation screening of young athletes? *Circulation*. 2007;116:2610-5.
- Pannell LM, Reys EM, Underwood SR. Cardiac risk assessment before non-cardiac surgery. *Eur Heart J Cardiovasc Imaging*. 2013;14:316-22.
- Sociedade Brasileira de Cardiologia. Diretrizes da Sociedade Brasileira de Cardiologia e Emissão de Relatórios de Exames Eletrocardiográficos 2009. *Arq Bras Cardiol*. 2009;93:1-19. Available at: http://publicacoes.cardiol.br/consenso/2009/diretriz_ecg_93supl02.pdf
- Bohmer AB, Defosse J, Geldner G, et al. Preoperative risk evaluation of adult patients for elective, noncardiac surgical interventions. Results of an on-line survey on the status in Germany. *Anaesthesist*. 2012;61:407-19.
- Okin PM, Devereux R, Nieminen MS, et al. Electrocardiographic strain pattern and prediction of cardiovascular morbidity and mortality in hypertensive patients. *Hypertension*. 2004;44:48-54.
- Denes P, Larson JC, Lloyd-Jones DM, et al. Major and minor ECG abnormalities in asymptomatic women and risk of cardiovascular events and mortality. *JAMA*. 2007;297:978-85.
- Sigurdsson E, Thorgeirsson G, Sigvaldason HM, et al. Unrecognized myocardial infarction: epidemiology, clinical characteristics. *Ann Intern Med*. 1995;122:96-102.
- Dalvig ML, Liao Y, Greenland P, et al. Association of nonspecific minor ST-T abnormalities with cardiovascular mortality: the Chicago Western Electric Study. *JAMA*. 1999;281:530-6.
- Kumar A, Prineas RJ, Arnold A, et al. Prevalence, prognosis and implications of isolated minor nonspecific ST segment and T-wave abnormalities in older adults. *Cardiovascular Health Study*. *Circulation*. 2008;118:2790-6.
- Frolkis JP, Pothier CE, Blackstone MD, et al. Frequent ventricular ectopy after exercise as a predictor of death. *N Engl J Med*. 2003;348:781-90.
- Samol A, Masin M, Gellner R, et al. Prevalence of unknown atrial fibrillation in patients with risk factors. *Europace*. 2013;15:657-62.