



Colombia Médica

ISSN: 0120-8322

colombiamedica@correounivalle.edu.co

Universidad del Valle

Colombia

Herrera, Julián Alberto

Primary prevention of preeclampsia: myth or reality?

Colombia Médica, vol. 46, núm. 4, octubre-diciembre, 2015, pp. 154-155

Universidad del Valle

Cali, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=28343295001>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Colombia Médica
colombiamedica.univalle.edu.co

Editorial

Primary prevention of preeclampsia: myth or reality?

Prevención primaria de preeclampsia: Mito o realidad?

Julián Alberto Herrera

Professor Emeritus of Medicine. Universidad del Valle, Cali, Colombia

Herrera JA: Primary prevention of preeclampsia: myth or reality?. *Colomb Med (Cali)*. 2015; 46(4): 154-155.

© 2015. Universidad del Valle. This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Article history: Received: 7 December 2015, Revised: 10 December 2015, Accepted: 18 December 2015

Preeclampsia is a disease of the human species, described more than a century ago, of unknown cause, which shares with atherosclerosis risk factors, biochemical abnormalities, endothelial dysfunction and a multifactorial origin. Reducing atherosclerosis lethality is achieved with secondary prevention; and similarly, in preeclampsia, early diagnosis and timely obstetric resolution (secondary prevention) prevents that the mother reaches the end of its natural history, where acute overlay of atherosclerotic plaques in the arteries spiral induces multiple organ failure and increases the maternal risk of death¹. The hospital care scheme implemented in Colombia achieved a reduction in maternal mortality by 9%, and when complemented with a preventive prenatal outpatient scheme (Biopsychosocial Model) designed to prevent progression of endothelial dysfunction, the reduction of maternal mortality increased up to 25%².

For decades, different approaches to the prevention of preeclampsia were investigated. One proposal was the prescription of acetylsalicylic acid in low doses (75-100 mg/day) to reduce inflammation and improve the balance between thromboxane A₂/prostacyclin. With this approach, a reduction of 10% was observed in the probability of preeclampsia appearing in the general obstetric population (OR= 0.90; 95% CI= 0.84 to 0.97), and a 25% reduction in the high-risk obstetric population (OR= 0.75; 95% CI= 0.66 to 0.85)³. Calcium supplementation was proposed to block the vasoconstrictor effect of parathyroid hormone (PTH). With this strategy, it was observed a decrease of 37% in the probability of having preeclampsia (OR= 0.63; 95% CI= 0.44 to 0.90); but two subsequent studies in which the supplementation began in the second quarter, using the same dose, did not show any protective effect (OR= 0.94; 95% CI= 0.76-1.16). But when the dose (<1 g/day elemental calcium) was reduced, it was estimated through nine studies that the probability of preeclampsia decreased up to 62% (OR= 0.38; 95% CI= 0.28 to 0.52)⁴.

Most studies evaluated the baseline calcium intake, but not the levels of maternal and placental hormones that regulate its metabolism, nor the circulating levels of ionized calcium. In a pregnant woman who has a high calcium intake and high supplementation, her supplemented calcium will be lost with the body fluids, since these hormones reduce its absorption and/or increase its excretion. In addition to lessening the effect of PTH, a proper balance in the vasomotor tone can be achieved by inducing calcium-dependent enzymes such as phospholipase A₂ and nitric oxide synthase, responsible for producing two potent vasodilators, nitric oxide and prostaglandin E₂.

In these studies the bioavailability of calcium was not controlled as a confounding factor. A proposal derived from it was investigated by adding conjugated linoleic acid (CLA) (450 mg/day) to the calcium supplementation (600 mg/day elemental calcium) in low doses in order to reduce lipid concentrations and inflammation⁵. This strategy reduced the risk by 80% (OR= 0.20; 95% CI= 0.44 to 0.82), taking into account studies in both food intake and the bioavailability of calcium⁶.

In the present issue of Colombia Medica, Alzate *et al.*⁷, compared calcium citrate supplementation in low doses and conjugated linoleic acid in pregnant women from the second quarter, compared with the use of calcium carbonate alone. They describe a preventive effect for preeclampsia in adolescents (OR= 0.000; 95% CI= 0.00 to 0.44) for the first treatment, compared with the usual scheme (calcium carbonate) (OR= 0.96; 95% CI= 0.73-1.27).

Why did the calcium supplementation with low doses plus linoleic acid show a greater risk reduction than calcium alone at high doses? Several studies in different populations have shown protective mechanisms for risk with the association of low doses of calcium plus linoleic acid by improving the balance between thromboxane A₂/prostaglandin E₂, by reducing intracellular free calcium, by antagonizing the activity of the renin-angiotensin-aldosterone, by reducing the endothelial dysfunction, and by reversing the metabolic syndrome^{5,6}, which has not been observed with calcium supplementation alone⁸.

Corresponding author:

Julian Herrera. Facultad de Salud. Universidad del Valle-Sede San Fernando, Calle 4B No 36-00 Cali, Colombia. Phone: +57 3155633740 Fax: +57 2 5565621 E-mail: Julian.herrera@correounivalle.edu.co.

Why was the risk reduction greater in adolescents than in older age groups? One could speculate that given their young age, they would be less exposed to chronic infections and gestational diabetes, factors associated with oxidative stress and endothelial dysfunction⁹; and because of their accelerated growth, the nutritional imbalance could be an important avenue for endothelial dysfunction¹⁰.

Primary prevention of disease is an important strategy because it reduces maternal mortality and prevents the leading cause of intrauterine growth restriction, i.e. preeclampsia, which is associated with fetal programming for cardiovascular diseases in the adult¹¹. There's no doubt that these preventive results are more relevant for African descendants and indigenous ethnic groups, in both of which teenage pregnancy is common, and that have lower attendance to prenatal care and institutional delivery, which would prevent perinatal hypoxia and cognitive deficits of children and adolescents¹². Preventing teen pregnancy and preventing maternal deaths are development goals which break this poverty circle⁹. The experience of our country with the Biopsychosocial model² has shown that it is possible to achieve the primary prevention of preeclampsia, by reducing the incidence of the disease in western Colombia in a pilot study, and then maternal mortality in the country in a sustainable way². The study by Alzate *et al.*⁷, makes an important light on the primary prevention of preeclampsia in adolescents. We need that the new generations of Colombians are not anymore exposed to fetal programming for life-threatening diseases in adverse perinatal environments, which very often increases disability and violence¹³. This is a challenge that must be undertaken for greater social equity, and to achieve the long-awaited consolidation of peace for Colombians.

References

1. Redman C. The six stages of preeclampsia. *Pregnancy Hypertens.* 2014;4(3):246–246.
2. Herrera JA, Herrera-Medina R, Herrera-Escobar JP, Nieto-Díaz A. Reduction of maternal mortality due to preeclampsia in Colombia-an interrupted time-series analysis. *Colomb Med (Cali)* 2014;45(1):25–31.
3. Hofmeyr GJ, Belizan JM, Von Dadelszen P, Calcium and Preeclampsia Study Group. Low-dose calcium supplementation for preventing pre-eclampsia: a systematic review and commentary. *BJOG.* 2014;121(8):951–957.
4. Campos A. The role of aspirin in preeclampsia prevention: State of the Art. *Acta Med Port.* 2015;28(4):517–524.
5. Campos Mondragón MG, Oliart Ros RM, Martínez Martínez A, Méndez Machado GF, Angulo Guerrero JO. Metabolic syndrome reversion by polyunsaturated fatty acids ingestion. *Med Clin (Barc)* 2013;141(12):513–518.
6. Herrera JA, Arevalo-Herrera M, Shahabuddin AKM, Ersheng G, Herrera S, *et al.* Calcium and conjugated linoleic acid reduces pregnancy-induced hypertension and decreases intracellular calcium in lymphocytes. *Am J Hypertens.* 2006;19:381–387.
7. Roberts JM, Bell MJ. If we know so much about preeclampsia, why haven't we cured the disease. *J Reprod Immunol.* 2013;99(1-2):1–9.
8. Herrera JA, Chaudhuri G, López-Jaramillo P. Is infection a major risk to preeclampsia. *Med Hypotheses.* 2001;57(3):393–397.
9. Schwartz A. Maternal Mortality. Risk factors, Anthropological Perspectives, Prevalence in Developing Countries and Preventive Strategies for Pregnancy-Related Deaths. New York: Nova Science Publishers Inc; 2015.
10. Herrera JA, Arévalo-Herrera M, Villegas A, Herrera S, Villalba M, Bromet A. Suplementación oral de calcio en adolescentes embarazadas de bajo nivel socioeconómico y su efecto en las concentraciones de calcio libre intracelular. *Colomb Med.* 2006;37(S1):9–15.
11. Henriksen T, Clausen T. The fetal origins hypothesis: placental insufficiency and inheritance versus maternal malnutrition in well-nourished populations. *Acta Obstet Gynecol Scand.* 2002;81(2):112–114.
12. Pallito EK, Kilbride HW. Perinatal outcome and later implications of intrauterine growth restriction. *Clin Obstet Gynecol.* 2006;49(2):257–269.
13. Ukleba KO, Pavlenishvili IV, Zurabashvili DZ. The influence of perinatal hypoxic-ischemic disorders of central nervous system on adolescent academic achievements and behavioural deviations. *Georgian Med News.* 2013;218:58–62.