



Acta Scientiae Veterinariae

ISSN: 1678-0345

ActaSciVet@ufrgs.br

Universidade Federal do Rio Grande do Sul
Brasil

Navarra Baticados, Waren; Inoue, Noboru; Sugimoto, Chihiro; Nagasawa, Hideyuki; Morales
Baticados, Abigail
Genomic Cloning and Sequence Analysis of Trypanosoma brucei rhodesiense Gene Encoding
Putative N-glycosylation Enzyme
Acta Scientiae Veterinariae, vol. 38, núm. 3, 2010, pp. 257-264
Universidade Federal do Rio Grande do Sul
Porto Alegre, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=289021902005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Genomic Cloning and Sequence Analysis of *Trypanosoma brucei rhodesiense* Gene Encoding Putative N-glycosylation Enzyme

Waren Navarra Baticados¹, Noboru Inoue², Chihiro Sugimoto³, Hideyuki Nagasawa² & Abigail Morales Baticados¹

ABSTRACT

Background: *Trypanosoma brucei rhodesiense* is a haemoflagellate parasite of zoonotic significance. Aside from its public health importance, this parasite subspecies gained notoriety because of their effective system to circumvent the immune response of vertebrate host. The parasite cell surface is covered with millions of VSG dimers, which serve as an almost infinite repertoire of biomolecules needed for evasion of host immune system. Around two decades ago, it was resolved that all trypanosome VSG is associated with one or more N-linked oligosaccharides, with a range of structures including high mannose and complex types. This complex process of protein modification known as N-linked glycosylation is catalyzed by oligosaccharyl transferase (OST). In general, the incorporation of glycan structures can alter protein's antigenic properties and recently it was established that glycan molecules associated with VSG were found to be important in several aspects of trypanosome-host interaction, especially during parasite evasion of the host defense mechanisms. Therefore, our major interest is to clone and characterize the trypanosome OST.

Materials, Methods & Results: The template genomic DNA for PCR amplification was extracted as described previously. In an attempt to clone *Trypanosoma brucei rhodesiense* putative oligosaccharyl transferase, an amplicon of ~2000 bp was obtained having an open reading frame of 2057 bp and deduced primary structure composed of 685 amino acid residues (TbrOST II). Comparison of TbrOST II ORF with annotated putative oligosaccharyl transferase in the genome of other organisms revealed sequence identity to other kinetoplastid. TbrOST II had high nucleotide (Ns) and amino acid (As) sequence similarity with the genomes of *T. brucei gambiense* (Ns:99%; As:78%) and *T. brucei* (Ns:95-98%; As:77%-98%). There was also significant nucleotide and amino acid sequence identity in the genomes of *T. cruzi* (Ns:74%; As:63%), *Leishmania infantum* (Ns:70-83%; As:46-57%), *L. braziliensis* (Ns:69-81%; As:46-55%) and *L. major* (Ns:69-80%; As:46-57%). Sequence similarity (71-77%) from other origins was also exhibited. The nucleotide sequence alignments and analysis were performed using the Oxford University Mac Vector 6.5 sequence analysis software and CLC Workbench 5.6 software.

Discussion: The nucleotide BLAST results indicate that sequence identity is higher between species of the same genus rather than of the same family. It is known that *T. brucei*, *T. gambiense* and *T. rhodesiense* are members of the *Brucei*-complex or *Brucei* group. Although *T. brucei brucei* has more similarities with *T. brucei rhodesiense* than *T. brucei gambiense*, these parasites are morphologically indistinguishable. This is the probable reason why high sequence identity was displayed by other subspecies of the *Brucei* group. In addition, the high percent identity possessed by TbrOST II with other trypanosomatids agrees with the evolutionarily conserved characteristics of the established OST. The DNA sequence data of TbrOST II showing similar sequences in the genome of other organisms further corroborate the previous reports regarding the ubiquitous nature of OST in other life forms. Based on the size of the amplicon and significant percentage of nucleotide and amino acid sequence identity to homologues within the genome of related species and various organisms, the results strongly indicate that TbrOST II is a trypanosome oligosaccharyl transferase gene candidate that should be fully characterized and subjected to functional genomic studies. The study reports the molecular cloning and sequencing of a potential oligosaccharyl transferase gene in *T. brucei rhodesiense* (TbrOST II). The sequence data has been deposited in the GenBank with accession number GU475126.

Keywords: *Trypanosoma brucei rhodesiense*, putative oligosaccharyl transferase, N-glycosylation.

INTRODUCTION

Trypanosoma brucei rhodesiense is a haemoflagellate parasite of zoonotic significance. A wide range of mammalian fauna, especially domestic livestock and wild bovids, serves as reservoir host [23,26]. Aside from its public health importance, this parasite subspecies gained notoriety because of their effective system to circumvent the immune response of vertebrate host. The parasite cell surface is covered with millions of VSG dimers, which serve as an almost infinite repertoire of biomolecules needed for evasion of host immune system. Unique VSG are alternately produced by sequential expression of about a thousand trypanosome VSG gene reservoir per parasite; a phenomenon described as antigenic variation [5,6]. Around two decades ago, it was resolved that all trypanosome VSG is associated with one or more N-linked oligosaccharides, with a range of structures including high mannose and complex types [7,19,21,24]. This complex process of protein modification is generally known as glycosylations, one of which is N-linked glycosylation.

The central event in the N-linked glycosylation process is catalyzed by oligosaccharyl transferase (OST) [11]. It catalyzes the co-translational addition of preassembled oligosaccharide complexes (Dol-PP-GlcNAc₂Man₃Glc₃) to an asparagine residue in an Asn-Xaa-Ser/Thr consensus sequon (Xaa can be any amino acid excluding proline) of the growing nascent polypeptide chain being translocated into the endoplasmic reticulum through a structure called translocon [4].

OST appears ubiquitous among eukaryotes and conserved throughout eukaryotic evolution. OST has been molecularly isolated and purified from mammalian sources, avian species and yeast *Saccharomyces cerevisiae* [3,11-14,22] as a multimeric enzyme.

In general, the incorporation of glycan structures to different protein moieties is precedent towards proper protein folding and stability, intracellular targeting, intercellular recognition, hormone synthesis, anti-apoptotic response, control of salt/osmotic stress, and cell surface expression of some glycoproteins [8,9,15,17]. Glycans can alter protein's antigenic properties and recently it was established that, glycan molecules associated with VSG were found to be important in several aspects of trypanosome-host interaction, especially during parasite evasion of the host defense mechanisms [16,20,21]. Therefore our major interest is to clone and characterize the trypanosome OST.

MATERIALS AND METHODS

Laboratory animals

Female 8-week-old BALB/c mice¹ were used in the study. The animal room was maintained at 22 ± 3°C with a 12:12 h of light-dark cycle. All experiments were conducted according to the guidelines for the care and use of laboratory animals, Obihiro University of Agriculture and Veterinary Medicine, Japan.

Trypanosomes and cultivation

Trypanosoma brucei rhodesiense IL2343 strain (Ivory Coast) stabilates were maintained in BALB/c mice. After reaching high parasitemia (~100 BSFs/field at magnification of x400), infected blood was collected by intracardiac puncture, cultured and maintained *in vitro* in HMI-9 medium supplemented with 10% fetal bovine serum² as previously described [10]. The HMI-9 *in vitro* cultivation medium was prepared using the following composition: Iscove's modified dulbecco's medium³ (IMDM), 10 mM bathocuproine disulfonic acid³, 100 mM pyruvic acid sodium salt³, 16 mM thymidine³, 40 mg/mL bovine serum albumin³, 1 mg/mL bovine holo-transferrin, 100 mM hypoxanthine⁴, 150 mM L-cysteine hydrochloride³, 14 mM 2-mercaptoethanol⁵, 25 mM HE-PES³ pH 7.2, 200 mM L-glutamine⁶, 10%v/v heat inactivated fetal bovine serum², 100 U/mL penicillin⁷ and 100 µg/mL streptomycin⁷.

DNA extraction

The template genomic DNA for PCR amplification was extracted as described previously [2]. Briefly, *T. brucei rhodesiense* IL2343 genomic DNA was extracted by adding 9 volumes of extraction buffer (0.2 M NaCl⁸, 10 mM Tris-HCl^{3,8} pH 8.0, 10 mM EDTA³ pH 8.0 and 1% SDS), proteinase K⁶ to a final concentration of 100 µg/mL and followed by 6 h incubation at 55°C with gentle agitation. Overnight incubation was performed after additional proteinase K was placed. Genomic DNAs were phenol-chloroform-isoamyl alcohol³ extracted, ethanol precipitated, and resuspended in Tris-EDTA buffer, pH 8.0 or deionized water. The concentration of the sample DNA was determined by spectrophotometry.

Polymerase chain reaction amplification of putative OST gene

The primers were designed from the nucleotide sequences of *T. brucei* genomic clones as guided by EMBL-EBI Parasite Genomes WU-Blast 2 database

search (www.ebi.ac.uk/blast/para-sites.html) for African trypanosomes with *L. major* putative OST STT3 subunit sequence as the query [EMBL Q9U5N8 (AJ251127.1)]. PCR amplification was performed using a forward primer⁹ (5'-TGG TAC GAC TAC ATG AGC TGG TAC CCG CT-3') and a reverse primer⁹ (5'-TGG ATC TCC TTC GCT GGC GGG TAC TG-3'). Distilled water was used as template for negative control reaction. The samples were programmed to a temperature-step cycle of 94°C at 10 min, 94°C at 30 s, 60°C at 30 s for a total of 30 cycles followed by 4 min extension at 72°C. The PCR products were analyzed by electrophoresis on 1% TAE (Tris-acetate-EDTA) agarose gel. The PCR product was then processed for cloning after agarose gel extraction using a commercial kit¹⁰ according to the manufacturer's instructions.

Cloning and sequencing of PCR products

The PCR product was ligated into *EcoRV* site of pT7 bluescript plasmid vector¹¹ using Takara solution I ligation kit¹². Ligation reaction was transformed into DH5 α competent *E.coli* cells and plated on Luria Britani's-ampicillin (LB-amp) agar dishes. The presence of insert was confirmed by restriction digest against *Hind* III & *Xba* I site from the cloning site of the plasmid vectors flanking the PCR product.

Prior to sequencing, twenty-five cycles of Bigdye PCR was carried out in a total volume of 5 μ L and were performed using the following standard condition: 96°C at 2 min, 96°C at 10 s, 50°C at 5 s and 60°C at 4 min. Sequencing was started by the single strand dideoxynucleotide-chain-termination method using a cycle sequencing kit¹³, DNA sequence analyzer¹³ and T7 promoter primer and pUC/M13 reverse primers⁹. The second set⁹ [forward (5'-GAC ATA CAG CGT CAG TTT GC-3'); reverse (5'-GAT GAA TGT GAG TGA AGA GAG C-3')] and the third set⁹ [forward (5'-CGT TCG GAT TCT TCA AAC CTA CAG-3') and reverse (5'-AAT ACG GGC ATC TTC AGG CG-3')] primers were used to obtain the partial nucleotide sequence. The nucleotide sequence alignments and analysis were performed using sequence analysis softwares^{14,15}.

RESULTS

A putative trypanosome oligosaccharyl transferase gene was successfully amplified using *T. brucei rhodesiense* crude DNA as the template. Genomic clone of approximately 2000 bp was obtained after PCR amplification (Figure 1). Nucleotide sequencing revealed

that the amplified band was composed of 2057 bp partial nucleotide sequence (Figure 2). The deduced partial primary structure of the *T. brucei rhodesiense* putative oligosaccharyl transferase clone II (TbrOST II) was composed of 685 amino acid sequence (Figure 2). Determination of the sequence homology in other organisms was carried out using NCBI Basic Local Alignment Search Tool (**BLAST**) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) [1].

Sequence analysis showed that TbrOST II had significant nucleotide (Ns) and amino acid sequence (As) percent identity to putative oligosaccharyl transferase subunit coming from other kinetoplastid genome in the NCBI public database.

When compared with the related trypanosome species, the DNA size of *T. brucei rhodesiense* putative oligosaccharyl transferase was 200 bp than *T. brucei gambiense* (1842 bp, FN554968.1). In addition, the sequence homologues from *T. brucei* (2466 bp, XM_839672.1) (2466 bp, XM_839671.1) (2406 bp, AC159432.1) (2406 bp, XM_839670.1), exceeded TbrOST II clone by >400 bp. The Latin American trypanosome species, *T. cruzi*, with percent identity to TbrOST II on the other hand had a molecular weight of 2397 bp. (XM_803446.1). Furthermore, the data showed that the annotated DNA size of the homologues under genus *Leishmania* was greater than that of genus *Trypanosoma* with a disparity ranging from




Figure 1. Polymerase Chain Reaction amplification of *Trypanosoma brucei rhodesiense* Putative Oligosaccharyl transferase clone II (TbrOST II). Lane (A), (B) & (C) represents 1 kbp DNA ladder, negative control reaction and ~ 2,000 bp PCR amplicon respectively.

~300-500 bp (*L. infantum*) and ~200- 500bp (*L. major* & *L. braziliensis*).

Leishmania infantum displayed the greatest difference in terms of DNA molecular weight to TbrOST II (2592 bp, XM_001468890.1) (2385 bp, XM_001468892.1) (2355 bp, XM_001468891.1) followed by *L. braziliensis* (2580 bp, XM_001567553.1) (2319 bp, XM_001568165.1) (2565 bp, XM_001568167.1) (2472 bp, XM_001568166.1) and the least was *L. major* (2574 bp, XM_838130.1) (2322 bp, XM_838128.1) (2559 bp, XM_838127.1) (2373 bp, XM_838129.1) (2502 bp, AJ251127.1).

The molecular size of the amplicon showed akin to several homologues under order kinetoplastida. Taking into consideration the reported DNA size of the homologues from closely related species, the data suggest that the full-length nucleotide sequence of *T. brucei rhodesiense* putative oligosaccharyl transferase is also not more than 3000 bp.

Furthermore, the results using BLAST nucleotide (Ns) and amino acid (As) homology search showed that

the *T. brucei rhodesiense* putative oligosaccharyl transferase had a very high percent sequence identity to the genome of *T. brucei* (Ns:98%, XM_839670.1; As:98%, XP_844763.1). Other homologues found in the genome of *T. brucei* (Ns:95%, XM_839672.1; As:78%, XP_844765.1) (Ns:96%, XM_839671.1; As:77%, XP_844764.1) and *T. brucei gambiense* (Ns:99%, FN554968.1; As:78%, CBH10989.1) also showed very high DNA sequence identity but with only fairly high amino acid sequence identity to TbrOST II. A moderate sequence similarity in the genome of *T. cruzi* (Ns:74%, XM_803446.1; As:63%, XP_808539.1) was also observed.

Subsequently, percent sequence identity hits in other members of family Trypanosomatidae, genus *Leishmania*, were also obtained (Figure 3). All the homologues from genus *Leishmania* displayed moderate and low nucleotide and amino acid sequence identity to TbrOST II respectively. The following are the different *Leishmania* clones from the GenBank with nucleotide and amino acid sequenced regions having local similarity

A	1	TGGTACGACT	ACATGAGCTG	GTACCCGCTT	GGCCGACCGG	TGGGACAAC	CATCTTCCCC
	61	GGAAATGCAGC	TTACCCGCTGT	AGCCATTTCAT	CGTGTGCTGG	AAATGCTCGG	CGCAGGTATG
	121	TCCATCAACA	ATATCTGTGT	GTACATTCTT	GCATGGTTCTG	GTAGTATTGC	CACGTGTTTG
	181	GCTGCTCTCA	TTGCGTACGA	ATCATCTAAT	TCGCTCAGTG	TCATGGCGTT	TACTGCGTAC
	241	TTTTTTTCCA	TCGTACCTGC	ACACCTGATG	CGATCAATGG	CTGGTGAATT	TGACAAATGAG
	301	TGTGTTGCAA	TGGCGGCGAT	GCTTCTGACG	TTCTACATGT	GGGTACGATC	GTACGCGACG
	361	TCAAAGTTCTG	GGCCCATTTGG	CGCTTTAGCT	GGTGTGGCAT	ACGGGTACAT	GGTGTCCACG
	421	TGGGGTGGTT	ATATTTTCTGT	GCTGAACATG	GTAGCCTTCC	ACGCTTCTGT	ATGTGTACTG
	481	CTTGATTGGG	CTCGTGGGAC	ATACAGCGTC	AGTTTGCTGA	GGGCGTATTCT	ACTGTTTTTT
	541	GTCAATTGGCA	CTGCCCTTGC	GATTGTGCTA	CCGCCAGTGG	AGTGGACGCC	CTTCCGGTGC
	601	CTGGAGCAAC	TGACAGCATT	GTTTGTCTTC	GTTTTCATGT	GGGCACTCCA	CTACTCGGAA
	661	TACCTGCGTG	AGCGTGCCCG	AGCGCCCAT	CACCTTCTTA	AAGCACTTCA	GATCCGTGCC
	721	CGCATTTTCA	TGGGCAACT	CTCCTTGTCT	TTGATTGTGG	CAAGTTTGTG	TGCCCGCTTC
	781	GGATTCCTCA	AACCTACAGC	GTACCCGCTC	CGTGCCTTGT	TCGTGAACAA	TACGCGTACC
	841	GGAAATCCCC	TCGTGGATTG	TGTGGCCGAG	CATCGGCCGA	GGACTGCGGG	GGCGTATCTG
	901	CGCTACTTTC	ATGTTTGTTA	CCCTCTTTGG	GGGTGCGGTG	GGCTCTCTAT	GTGGGTATTC
	961	ATGAAAAAGG	ACCGCTGGCG	CGCCATTGTT	TTTCTTGCTT	CACCTTCCAC	TGTTACGATG
	1021	TATTTACGCG	CCCGTATGTC	CGGATTACTT	CTGTTAGCGG	GTCCGCGAGC	AACGGCTTGC
	1081	GCCGGCATGT	TCATAGGGGG	GCTTTTGTAT	CTGGCGCTGT	CACAGTTTGG	TGATTTCGCT
	1141	AGCCCAAAAG	ATGCCCTCGG	CGATTCCGAT	CCCGCGGGAG	GGTCAAGACG	GGCAAGGGGC
	1201	AAAGTTGTTA	ATGAGCCTCC	CAAAAGAGCC	ATCCTTTAGTC	ACCGCTGGTT	TCAACGTTTA
	1261	GTGCAATCGT	TGCCCGTCCC	GCTACGACGT	GGTATCGCGG	TTGTGGTGTCT	CGTATGTCTC
	1321	TTGCGCAATG	CCATGAGACA	CTCATTCGAA	AAGTCCTGCG	AGAAAAATGGC	ACATGCACCT
	1381	TCATCTCCAA	GGATCATTGC	CGTGACTGAT	CTACCCAATG	GAGAGAGAGT	CCTCGCCGAT
	1441	GATTACTACG	TGTCGTACTT	GTGGCTGCGA	AACAATACGC	CTGAAGATGC	CGGTATTCTC
	1501	TCATGGTGGG	ACTACGGGTA	TCAAACTACT	GGAAATTGGCA	ATCGCACAAAC	CCTTGCAGAT
	1561	GGTAACACAT	GGAGTCACAA	GCACATAGCA	ACTATTGGAA	AGATGCTTAC	ATCCCTGTG
	1621	AAGGAGTCAC	ATGCTCTTAT	ACGCCATCTC	GCTGATTATG	TGCTGATATG	GTCTGGTCAA
	1681	GATGGCAGCG	ATTACTTAA	ATCGCCACAC	ATGGCTCGGA	TAGGCAACAG	TGTATATCGC
	1741	GATATGTGTT	CAGAAGACGA	TCCGCTGTGT	AGACAGTTCC	GGTTTTATAG	TGGTGACCTC
	1801	AATAAGCCTA	CGCCTATGAT	CGACGGGTCC	CTATTATACA	ATGACACAG	TTTGGTACG
	1861	GATGGCGGGA	AGACACAAC	GGATAAGAAC	ATGTTTCAAG	TCGCTCATGT	GTCAAAGTAT
	1921	GGTTTGGTGA	AGATCTACAA	GGTGATGAAT	GTGAGTGAAG	AGAGCAAGGC	GTGGGTTGCA
	1981	GACCCAAAGA	ACCGCGTATG	CGACCCGCC	GGATCTTGGA	TATGCGCCGG	CCAGTACCCG
	2041	CCAGCGAAGG	AGATCCA				
B	1	WYDYSWYPL	GRPVGTTIFP	GMQLTGVAIH	RVLEMLGRGM	SINNICYIYP	AWFGSIATVL
	61	AALAYESSN	SLSVMAFTAY	FFSIVPAHLM	RSMAGEFDNE	CVAMAAMLIL	FYMWVRSLSR
	121	SSSWPIGALA	GVAYGYMWST	WGGYIFVLM	VAFHASVCVL	LDWARGTYSV	SLLRAYSLEF
	181	VIGTALAICV	PPVEWTPERS	LEQLTALFVF	VFMWALHYSE	YLREARAPI	HSSKALQIRA
	241	RIEMGTLSLL	LIVASLLAPF	GFFKPTAYRW	RALEVKHTRT	GNPLVDVSAE	HRPTTAGAYL
	301	RYFHVCPFLW	GCGGLSMLVF	MKKDRWRATV	FLASLSVTVM	YFSAIRISRL	LLAGPAATAC
	361	AGMEIGGLFD	LALSQFGLR	SPKDGSDSD	PAGGSKRAKG	KVNEPPKRA	IFSHRWFLRL
	421	VQSLPVPLRR	GIADVVLVCL	FANPMRHSFE	KSCCKMAHAL	SSPRIIAVTD	LPNGERVLAD
	481	DYVSYLWLR	NNTPEDARIL	SWWDYGYQIT	GIGNRTTLAD	DMTWSHKHIA	TIGKMLTSPV
	541	KESHALIRHL	ADYVLWSDG	DGSDLLKSPH	MARIGNSVYR	GNCSEDDPLC	RQFGFYSDPL
	601	NKPTPMQRS	LLYHLHFGT	DGGKTQLDKN	MFQLAYVSKY	GLVKIYKVMN	VSEESKAWVA
	661	DPKNRVCDDP	GSWICAGQYP	PAKEI			

Figure 2. Partial Nucleotide (A) and Deduced Amino Acid (B) Sequence of *Trypanosoma brucei rhodesiense* Putative Oligosaccharyl transferase (TbrOST II) (GenBank: GU475126).

to TbrOST II: *L. infantum* (Ns:70%, XM_001468891.1; As:57%, XP_001468928.1) (Ns:70%, XM_001468890.1; As:56%, XP_001468927.1) (Ns:83%, XM_001468 892.1; As:46%, XP_001468929.1); *L. braziliensis* (Ns:69%, XM_001565753.1; As:54%, XP_00156 5803.1) (Ns:69%, XM_001568165.1; 55%,As:XP_001568215.1) (Ns:70%, XM_00156 8167.1; As:55%, XP_001568217.1) (Ns:81%, XM_0015681 66.1; As:46%, XP_001568216.1); *L. major* (Ns:69%, XM_838130.1; As:55%, XP_843223.1) (Ns:70%, XM_838128.1;As:57%, XP_843221.1) (Ns:70%, XM_838127.1; As:55%, XP_843220.1) (Ns:80%, XM_838129.1; As:46%, XP_843222.1) (Ns:80%, AJ251127.1; As:46%, CAB61 569.1).

```

TbrOSTII -----
Tbrucei  MTKGGKVAVTKGSAQSDGAGEGGMSSKAKSSTTFVATGGGSLPAWALKAVSTIVSAVILIY
Tbgam    MTKGGKVAVTKGSAQSDGAGEGGMSSKAKSSTTFVATGGGSLPAWALKAVSTIVSAVILIY

TbrOSTII -----WYDYSWYPLGRPVG
Tbrucei  SVHRAYDIRLTSVRLYGELIHEFDPWFNYRATQYLSDNWGRAFFQWYDYSWYPLGRPVG
Tbgam    SVHRAYDIRLTSVRLYGELIHEFDPWFNYRATQYLSDNWGRAFFQWYDYSWYPLGRPVG

TbrOSTII TTIFPGMQLTGVAIHRVLEMLGRGMSINNICYIIPAWFGSIATVLAALIAYESSNSLSVM
Tbrucei  TTIFPGMQLTGVAIHRVLEMLGRGMSINNICYIIPAWFGSIATVLAALIAYESSNSLSVM
Tbgam    TTIFPGMQLTGVAIHRVLEMLGRGMSINNICYIIPAWFGSIATVLAALIAYESSNSLSVM

TbrOSTII AFTAYFFSIVPAHLMRSMAGEFDNECVAMAAMLLTFYMWVRSLSRSSSWPIGALAGVAYG
Tbrucei  AFTAYFFSIVPAHLMRSMAGEFDNECVAMAAMLLTFYMWVRSLSRSSSWPIGALAGVAYG
Tbgam    AFTAYFFSIVPAHLMRSMAGEFDNECVAMAAMLLTFYMWVRSLSRSSSWPIGALAGVAYG

TbrOSTII YMVSTWGGYIFVLNMVAFHASVVCVLLDWARGTYSVSLLRAYSLEFFVIGTALAICVPPVEW
Tbrucei  YMVSTWGGYIFVLNMVAFHASVVCVLLDWARGTYSVSLLRAYSLEFFVIGTALAICVPPVEW
Tbgam    YMVSTWGGYIFVLNMVAFHASVVCVLLDWARGTYSVSLLRAYSLEFFVIGTALAICVPPVEW

TbrOSTII TPFRSLEQLTALFVFMWALHYSEYLRERARAPIHSSKALQIRARIFMGTLSELLIVAS
Tbrucei  TPFRSLEQLTALFVFMWALHYSEYLRERARAPIHSSKALQIRARIFMGTLSELLIVAS
Tbgam    TPFRSLEQLTALFVFMWALHYSEYLRERARAPIHSSKALQIRARIFMGTLSELLIVAI

TbrOSTII LLAPFGFFKPTAYRVRALFVKHTRTGNPLVDSVAEHRPTTAGAYLRYFHVVCYPLWCGGGL
Tbrucei  LLAPFGFFKPTAYRVRALFVKHTRTGNPLVDSVAEHRPTTAGAYLRYFHVVCYPLWCGGGL
Tbgam    YLFSTGYFRSFSSRVRALFVKHTRTGNPLVDSVAEHRPTTAGAFLRHLHVCYNGWIIGFF

TbrOSTII SMLVFMKKDRWRAIVFLASLSTVTMYFSARMSRLLLLLAGPAATACAGMFIGGLFDLALSQ
Tbrucei  SMLVFMKKDRWRAIVFLASLSTVTMYFSARMSRLLLLLAGPAATACAGMFIGGLFDLALSQ
Tbgam    FMSVSCFFHCTPGMSFLLYSILAYYFSLKMSRLLLLSAPVASILTGYVVGSIVDLAADC

TbrOSTII FGDLSRSPKDasGSDPAGGSKRAKGKVVNEPPKRAIF----SHRWFQRL-VQSLPVPLRR
Tbrucei  FGDLSRSPKDasGSDPAGGSKRAKGKVVNEPPKRAIF----SHRWFQRL-VQSLPVPLRR
Tbgam    FA-----ASGTEH--ADSKEHQGKARGKGQRQITVECGCHNPFYKLCNSFSSRLVV

TbrOSTII GIAVVVLVCLFANP--MRHSFEKSCEKMAHALSSPRIIAVTDLPNGERVLADDDYVSYLW
Tbrucei  GIAVVVLVCLFANP--MRHSFEKSCEKMAHALSSPRIIAVTDLPNGERVLADDDYVSYLW
Tbgam    GKFFVVVVLAIICGPTFLGSEFRAHCERFSLSVANPRIIS-SIRHSGKLVLADDDYVSYLW

TbrOSTII LRNNTPEDARILSWWDYGYQITGIGNRTTLADGNTWSHKHIATIGKMLTSPVKESHALIR
Tbrucei  LRNNTPEDARILSWWDYGYQITGIGNRTTLADGNTWSHKHIATIGKMLTSPVKESHALIR
Tbgam    LRNNTPEDARILSWWDYGYQITGIGNRTTLADGNTWNHEHIATIGKMLTSPVKESHALIR

TbrOSTII HLADYVLIWSGQDGDLLKSPHMARIGNSVYRDMCSEDDPLCRQFGFYSGDLNKPTMMQ
Tbrucei  HLADYVLIWAGEDRGDLLKSPHMARIGNSVYRDMCSEDDPRCRQFGFEGGDLNKPTMMQ
Tbgam    HLADYVLIWSGQDRGDLRKRSHMARIGNSVYRDMCSEDDPLCRQFGFYSGDLNKPTMMQ

TbrOSTII RSLLYNLHFRFGTDGDKTQLDKNMFQLAYVSKYGLVKIYKVMNVSEESKAWVADPKNRVCD
Tbrucei  RSLLYNLHFRFGTDGDKTQLDKNMFQLAYVSKYGLVKIYKVMNVSEESKAWVADPKNRVCD
Tbgam    RSLLYNLHFRFGTDGDKTQLDKNMFQLAYVSKYGLVKIYKVMNVSEESKAWVADPKNRKCD

TbrOSTII PPGSWICAGQYPPAKEI-----
Tbrucei  PPGSWICAGQYPPAKEIQDMLAKRFHYE-----
Tbgam    APGSWICAGQYPPAKEIQDMLAKRIDYELEDNRRNRSDAHYRAYMRQMG

```

Figure 3. Amino Acid sequence identity of *Trypanosoma brucei rhodesiense* Putative Oligosaccharyl transferase (TbrOST II) within genome of genus *Trypanosoma*, family Trypanosomatidae. Aligned sequences were from *T. brucei* -Tbrucei (98%, XP_844763.1) and *T. brucei gambiense* -Tbgam (78%, CBH10989.1) species.

DISCUSSION

The surface coat of trypanosome species were previously reported to have marked diversities in N-glycosylation [20]. This prompted us to hypothesize that trypanosomes also possess N-glycosylation enzyme. During the attempt to clone *T. brucei rhodesiense* putative oligosaccharyl transferase, a genomic clone of ~2000 bp was acquired.

Notably, *L. major* putative OST STT3 subunit sequence was used as the query during the primer design for PCR amplification of TbrOST II. The efficient amplification of the gene in genus *T. brucei rhodesiense* using primers designed from *L. major* stt3 gene as the query strongly indicate that TbrOST II seems to be conserved within the family Trypanosomatidae. The nucleotide BLAST results indicates that sequence identity is higher between species of the same genus rather than of the same family. It is known that *T. brucei*, *T. gambiense* and *T. rhodesiense* are members of the *Brucei*-complex or *Brucei* group. Although *T. brucei brucei* has more similarities with *T. brucei rhodesiense* than *T. brucei gambiense*, these parasites are morphologically indistinguishable [18,23]. This is the probable reason why high sequence identity was displayed by other subspecies of the *Brucei* group. This also indicates that the functional unit of this putative oligosaccharyl transferase is conserved within the *Brucei* complex trypanosomes. In addition, the high percent identity possessed by TbrOST II with other trypanosomatids agrees with the evolutionarily conserved characteristics of the established OST [25, 27]. Consequently, DNA identity searches within the public databases obtained homologues in the genome of mammalian, nematode, arthropod and algae species. Selected organisms include *Mus musculus* (Ns:71%, NM_024222.2; As:29%, NP_077184.2), *Schistosoma mansoni* (Ns:73%, XM_002577919.1; As:29%, XP_002577965.1), *Drosophila pseudoobscura pseudoobscura* (Ns:77%, XM_002134519.1; As:31%, XP_002134555.1), *Thalassiosira pseudonana* (Ns:77%, XM_002288187.1; As:46%, XP_002288223.1) and *Phaeodactylum tricornutum* (Ns:73%, XM_002185331.1; As:46%, XP_002185367.1). Moreover, obtained amino acid sequence identity in the genome of fly, nematode, mouse and

algae homologues also exhibited lower sequence similarity (29-46%) than the previously documented percent identity (~50%) of reported oligosaccharyl transferase in other eukaryotic species [25].

CONCLUSIONS

Finally, even currently regarded as putative, DNA sequence data of TbrOST II showing similar sequences in the genome of other organisms further corroborate the previous reports regarding the ubiquitous nature of OST in other life forms. Based on the size of the amplicon and significant percentage of nucleotide and amino acid sequence identity to homologues within the genome of related species and various organisms, the results strongly indicate that TbrOST II is a trypanosome oligosaccharyl transferase gene candidate that should be fully characterized and subjected to functional genomic studies. The study reports the molecular cloning and sequencing of a potential oligosaccharyl transferase gene in *T. brucei rhodesiense* (TbrOST II). The sequence data has been deposited in the GenBank with accession number of GU475126.

Acknowledgment. This work was supported by the Japanese Ministry of Education, Culture, Sports, Science and Technology (Monbukagakusho).

SOURCES AND MANUFACTURERS

¹CLEA Japan, Inc., Tokyo, Japan.

²Biosource International Inc., Camarillo, CA, USA.

³Sigma, St. Louis, USA.

⁴Calbiochem, La Jolla, CA, USA.

⁵BDH Chemicals, Poole, England.

⁶Invitrogen Co., Carlsbad, CA, USA.

⁷Meiji Seika Kaisha, Ltd., Tokyo, Japan.

⁸Wako Pure Chemical Industries Ltd., Osaka, Japan.

⁹Sigma-Genosys, Hokkaido, Japan.

¹⁰GeneClean kit II, Q Biogene, Vista, CA, USA.

¹¹Novagen, Madison, WI, USA.

¹²TAKARA Shuzo, Japan.

¹³Applied Biosystems, Foster City, CA, USA.

¹⁴Mac Vector 6.5 sequence analysis software, Oxford University, England.

¹⁵CLC Workbench 5.6 software. USA.

Baticados W.N., Inoue N., Sugimoto C., Nagasawa H. & Baticados A.M. 2010. Genomic Cloning and Sequence Analysis of *Trypanosoma brucei rhodesiense* Gene Encoding Putative N-glycosylation... *Acta Scientiae Veterinariae*. 38(3): 257-264.

REFERENCES

- 1 Altschul S.F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W. & Lipman D.J. 1997. "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs". *Nucleic Acids Research*. 25(17): 3389-3402.
- 2 Baticados W.N., Witola W.H., Inoue N., Kim J., Kuboki N., Xuan X., Yokoyama N. & Sugimoto C. 2005. Expression of a Gene Encoding *Trypanosoma congolense* Putative Abc1 Family Protein is Developmentally Regulated. *Journal of Veterinary Medical Science*. 67(2): 157-164.
- 3 Breuer W. & Bause E. 1995. Oligosaccharyl transferase is a constitutive component of an oligomeric protein complex from pig liver endoplasmic reticulum. *European Journal of Biochemistry*. 228(3): 689-696.
- 4 Brodsky J.L. 1997. Translocation of proteins across the endoplasmic reticulum membrane. *International Review of Cytology*. 178: 277-328.
- 5 Cross G.A.M. 2002. Antigenic variation in African trypanosomes and malaria. In: *Molecular Medical Parasitology*. Marr J.J., Nilsen T.W. & Komuniecki R.W. (Eds). Academic Press: New York, pp.89-110.
- 6 Donelson J.E., Hill K.L. & El-Sayed N.M.A. 1998. Multiple mechanism of immune evasion by African trypanosomes. *Molecular & Biochemical Parasitology*. 91(1): 51-66.
- 7 Ferguson M.A.J. 1997. The structure and biosynthesis of trypanosomatid glycosylphosphatidylinositols. In: Hide G. & Mottram J.C. (Eds). *Trypanosomiasis and Leishmaniasis*. U.K.: Commonwealth Agricultural Bureaux International, pp.65-77.
- 8 Garcia R.C., Cundell D.R., Tuomanen E.I., Kolakowski Jr. L.F., Gerard C. & Gerard N.P. 1995. The role of N-glycosylation for functional expression of the human platelet-activating factor receptor. Glycosylation is required for efficient membrane trafficking. *Journal of Biological Chemistry*. 270(11): 25178-25184.
- 9 Goder V., Bieri C. & Spiess M. 1999. Glycosylation can influence topogenesis of membrane proteins and reveals dynamic reorientation of nascent polypeptides within the translocon. *Journal of Cell Biology*. 147(2): 257-265.
- 10 Hirumi H. & Hirumi K. 1991. *In vitro* cultivation of *Trypanosoma congolense* bloodstream forms in the absence of feeder cell layers. *Parasitology*. 102(2): 225-236.
- 11 Kaplan H.A., Welply J.K. & Lennarz W.J. 1987. Oligosaccharyl transferase: the central enzyme in the pathway of glycoprotein assembly. *Biochimica et Biophysica Acta*. 906(2): 161-173.
- 12 Kelleher D.J. & Gilmore R. 1997. DAD1, the defender against apoptotic cell death, is a subunit of the mammalian oligosaccharyltransferase. *Proceedings of the National Academy of Sciences*. 94(10): 4994-4999.
- 13 Knauer R. & Lehle L. 1999. The oligosaccharyltransferase complex from *Saccharomyces cerevisiae*. *Journal of Biological Chemistry*. 274 (24): 17249-17256.
- 14 Kumar V., Heinemann F.S. & Ozols J. 1994. Purification and characterization of avian oligosaccharyltransferase. Complete amino acid sequence of the 50-kDa subunit. *Journal of Biological Chemistry*. 269(18): 13451-13457.
- 15 Levin D. & Bartlett-Heubusch E. 1992. Mutants in the *S. cerevisiae* *PKC1* gene display a cell cycle-specific osmotic stability defect. *Journal of Cell Biology*. 116(5): 1221-1229.
- 16 Lisowska E. 2002. The role of glycosylation in protein antigenic properties. *Cellular and Molecular Life Sciences*. 59(3): 445-455.
- 17 Live D.H., Kumar R.A., Beebe X. & Danishefsky S.J. 1996. Conformational influences of glycosylation of a peptide: A possible model for the effect of glycosylation on the rate of protein folding. *Proceedings of the National Academy of Sciences*. 93(23): 12759-12761.
- 18 Matthews K.R. 2005. The developmental cell biology of *Trypanosoma brucei*. *Journal of Cell Science*. 118(2): 283-290.
- 19 Mehler A., Bond C.S. & Ferguson M.A.J. 2002. The glycoforms of a *Trypanosoma brucei* variant surface glycoprotein and molecular modeling of glycosylated surface coat. *Glycobiology*. 12(10): 607-612.
- 20 Parodi A.J. 1993. N-glycosylation in trypanosomatid protozoa. *Glycobiology*. 3(3): 193-199.
- 21 Pays E. & Nolan D.P. 1998. Expression and function of surface proteins in *Trypanosoma brucei*. *Molecular & Biochemical Parasitology*. 91(1): 3-36.
- 22 Silberstein S., Kelleher D.J. & Gilmore R. 1992. The 48-kDa subunit of the mammalian oligosaccharyltransferase complex is homologous to the essential yeast protein WBP1. *Journal of Biological Chemistry*. 267(33): 23658-23663.
- 23 Soulsby E.J.L. 1982. Phylum: Sarcostigophora Honigberg and Balamuth, 1963, Subphylum: Mastigophora Diesing, 1866. In: Soulsby E.J.L. (Ed). *Helminths, Arthropods and Protozoa of Domesticated Animals*. 7th edn. London: Baillière Tindall, pp.514-582.
- 24 Strickler J.E. & Patton C.L. 1980. *Trypanosoma brucei brucei*: Inhibition of glycosylation of the major variable surface coat glycoprotein by tunicamycin. *Proceedings of the National Academy of Sciences*. 77(3): 1529-1533.

Baticados W.N., Inoue N., Sugimoto C., Nagasawa H. & Baticados A.M. 2010. Genomic Cloning and Sequence Analysis of *Trypanosoma brucei rhodesiense* Gene Encoding Putative N-glycosylation... *Acta Scientiae Veterinariae*. 38(3): 257-264.

25 Yan Q. & Lennarz W.J. 2002. Studies on the function of oligosaccharyl transferase subunits. *Journal of Biological Chemistry*. 277(49): 47692-47700.

26 Zoller T., Fèvre E.M., Welburn S.C., Odiit M. & Coleman P.G. 2008. Analysis of risk factors for *T. brucei rhodesiense* sleeping sickness within villages in south-east Uganda. *BMC Infectious Diseases*. 8(88): 1-9.

27 Zufferey R., Knauer R., Burda P., Stagljar I., Heesen S., Lehle L. & Aebi M. 1995. STT3, a highly conserved protein required for yeast oligosaccharyl transferase activity *in vivo*. *EMBO Journal*. 14(20): 4949-4960.