



Acta Scientiae Veterinariae

ISSN: 1678-0345

ActaSciVet@ufrgs.br

Universidade Federal do Rio Grande do
Sul
Brasil

Resende de Carvalho, Isadora; Scantamburlo Denadai, Daniela; Teramachi Trevisan, Juliane; Falcão Mendes, Heloisa Maria; Venturoli Perri, Sílvia Helena; Rodrigues Monteiro, Cristina Maria; Zafalon Ferreira, Joana; Rossi Feliciano, Marcus Antonio; Cestari Pierucci, Julia; Russiano Vicente, Wilter Ricardo; Burkhardt de Koivisto, Marion
Follicular Histomorphometry and Evaluation of Ovarian Apoptosis in Queens of Different Age Groups

Acta Scientiae Veterinariae, vol. 44, 2016, pp. 1-8

Universidade Federal do Rio Grande do Sul

Porto Alegre, Brasil

Disponible en: <http://www.redalyc.org/articulo.oa?id=289043697034>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica

Red de Revistas Científicas de América Latina, el Caribe, España y Portugal

Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso abierto

Follicular Histomorphometry and Evaluation of Ovarian Apoptosis in Queens of Different Age Groups

Isadora Resende de Carvalho¹, Daniela Scantamburlo Denadai¹, Juliane Teramachi Trevisan¹,
Heloisa Maria Falcão Mendes¹, Sílvia Helena Venturoli Perri¹, Cristina Maria Rodrigues Monteiro¹,
Joana Zafalon Ferreira¹, Marcus Antonio Rossi Feliciano², Julia Cestari Pierucci²,
Wilter Ricardo Russiano Vicente² & Marion Burkhardt de Koivisto¹

ABSTRACT

Background: In humans and bitches the age is another factor that may affect the size of ovarian structures, verifying alterations in the quality of the pool and size of follicular structures, which can compromise the use of these structures for *in vitro* maturation. There are no reports correlating the morphometric characteristics of the follicles and ovarian apoptosis at different ages in cats. The aim of this study was to evaluate the histomorphometric parameters of follicular growth and the relationship with the occurrence of apoptosis in ovarian tissue of young, adult and senile queens.

Materials, Methods & Results: Eighteen domestic queens, multiparous, of different breeds and age groups were used in this study and divided into three groups according to their ages: five months to one year - young (7.8 ± 1.0 months); one to six years - adults (2.8 ± 0.5 years); and more than six years - senile (8.0 ± 0.9 years). Vaginal cytology was performed in order to characterize the estrous phase associated with plasma concentrations of progesterone. The morphology and percentage of the vaginal epithelium cells were evaluated and queens were classified into estrous and non-estrous and plasma concentrations of progesterone were determined. Ovarian samples were collected after ovariohysterectomy to routine histological processing and all follicles were counted and categorized into two groups, non-atresic and atresic. The mean follicular and oocyte diameters were calculated between the measurement of the largest diameter and perpendicular diameter. The relationship between follicle and oocyte were determined using the measurements of diameter, area and perimeter. The apoptotic cells were detected and cells were considered positive when TUNEL reaction was detected. The morphometric index of 1039 follicles were evaluated. Primordial follicles in young animals showed larger diameter, follicular area and perimeter than the structures of adult queens, as well as the unilaminar primary follicles of the same group were larger compared with senile animals ($P < 0.05$). Comparing adult and younger queens, the first showed a significant decrease of oocyte diameter in primary and unilaminar primary follicles, as well as oocyte area and perimeter ($P < 0.05$). The values for follicular diameter, oocyte area and perimeter for multilaminar primary, secondary and pre-ovulatory structures did not present statistical differences between the groups ($P > 0.05$). For the pre-ovulatory follicles there was no positive correlation between the oocyte growths regarding the follicles ($P > 0.05$). Only in senile animals positive markers for apoptosis were identified in nuclei of primordial follicles. No significant differences concerning the number of follicles and TUNEL positive cells were observed between groups ($P > 0.05$).

Discussion: Considering the importance of this study for greater knowledge in the basic aspects for reproductive biotechnologies, we verified that secondary follicles showed the largest diameters and younger animals the largest values for diameter, area and perimeter, suggesting that this age group could be ideal for the use and manipulation of oocytes. The process of follicular atresia is characterized by the occurrence of apoptosis, or programmed cell death when the organism begins to efficiently eliminate dysfunctional cells. The study of follicular apoptosis in small animals, especially in cats, is very important for the development of reproduction biotechnologies. Phenomenon of apoptosis showed no relationship with age in queens, occurring in a physiological, continuous and proportionate manner considering the number of non-dominant follicles involved in each estrous cycle.

Keywords: feline, follicle, reproduction.

Received: 24 November 2015

Accepted: 18 May 2016

Published: 24 June 2016

¹Departamento de Clínica, Cirurgia e Reprodução Animal, Faculdade de Medicina Veterinária, UNESP, Campus de Araçatuba, SP, Brazil. ²Setor de Obstetrícia Veterinária, Departamento de Medicina Veterinária Preventiva e Reprodução Animal, Faculdade de Ciências Agrárias e Veterinárias - UNESP, Jaboticabal, SP. CORRESPONDENCE: M.A.R. Feliciano [marcusfeliciano@yahoo.com.br - Tel.: +55 (16) 3209-2626]. FCAV/UNESP, Via de Acesso Prof. Dr. Paulo Donato Castellani, Bairro Distrito Industrial, s/n. CEP 14.884-900 Jaboticabal, SP, Brazil.

INTRODUCTION

The domestic cat is an important and necessary experimental model for reproduction biotechnologies. Reliable methods for the *in vitro* maturation and *in vitro* fertilization have been developed through the use of oocytes obtained from domesticated cats [18]. The study of ovarian folliculogenesis provides better understanding of the reproductive physiology and may be useful for improving the preservation of endangered wild cats [3].

Recent study using prepubertal and sexually mature queens evaluated the ovarian condition and follicle growth *in vitro* maturation and concluded that size and quality of follicles and oocytes can affect maturational ability [16]. According to studies in humans [17] and bitches [5], age is another factor that may affect the size of ovarian structures, in other words, the morphometric characteristics of the follicles are related to the age of individuals, verifying alterations in the quality of the pool and size of follicular structures, which can compromise the use of these structures for *in vitro* maturation.

The process of follicular atresia in the ovarian tissue is characterized by the occurrence of apoptosis of granulosa or theca cells [4,14]. To our knowledge, there are no reports correlating the morphometric characteristics of the follicles and ovarian apoptosis at different ages in cats.

The aim of this study was to evaluate the histomorphometric parameters of follicular growth and the relationship with the occurrence of apoptosis in ovarian tissue of young, adult and senile queens.

MATERIALS AND METHODS

Eighteen domestic queens, multiparous, of different breeds and age groups, with mean body weight of 3 kg were used in this study. None of the animals had any report of reproductive diseases and were considered healthy after physical and clinical examinations. The animals were divided into three groups according to their ages: five months to one year - young; (7.8 ± 1.0 months); one to six years - adults (2.8 ± 0.5 years); and more than six years - senile (8.0 ± 0.9 years). The age was estimated by analyzing the dental arch [6].

Vaginal cytology was performed in order to characterize the estrous phase associated with plasma concentrations of progesterone.

These samplings were performed prior to the ovariohysterectomy of the queens. Pre-anesthetic medication consisted of intramuscular administration of acepromazine (0.05 mg/kg; Acepran®)¹, midazolam (0.2 mg/kg; Dormire®)², and morphine (0.3 mg/kg; Dimorf®)²; induction protocol involved intravenous Ketamine (5 mg/kg; Vetaset®)³ and xylazine (0.5 mg/kg; Dopaser®)⁴ and epidural anesthesia consisted of (0.3 mL/kg = lidocaine 2% of epinephrine; Xylestesin®)², bupivacaine 0.5% with epinephrine (Neocaína®)² and fentanyl (2 mcg/kg; Fentanest®)².

Vaginal smears were obtained using a cotton sterile swab previously moistened with saline solution. The smears were stained with a modified Wright-Giemsa stain (Diff-Quick®)⁵ and analyzed by light microscopy (Olympus BX61)⁶. The morphology and percentage of the vaginal epithelium cells were evaluated [15] and queens were classified into estrous and non-estrous.

Blood samples (3 mL) were collected by jugular vein puncture. The blood was centrifuged immediately and plasma was stored at -18°C until hormone analysis. Plasma concentrations of progesterone were determined, in duplicate, by solid-phase I125 radioimmunoassay⁷.

In this study only animals classified as in a non-estrus period were enrolled, based on the results of progesterone levels and vaginal cytology evaluations [15].

After ovariohysterectomy the ovaries were fixed in paraformaldehyde solution at 5% (pH of 7.2 - 7.4) for 24 h and samples submitted to routine histological processing. Five-micron serial sections were mounted onto plain glass slides and stained with hematoxylin and eosin for light microscope (Olympus BX61)⁶ evaluation. All follicles were counted and categorized into two groups, non-atresic and atresic, following the morphological criteria established [2].

The non-atresic follicles were those with intact granulosa membrane and the presence of a few pyknotic nuclei ($\leq 5\%$ pyknotic nuclei) and atresic follicles showed attenuated granulosa membrane, rupture of granulosa cells and increased number of pyknotic cells ($\geq 15\%$ pyknotic nuclei) according to literature [2].

Only non-atresic follicles were considered and classified [7]: primordial, primary oocytes surrounded by a single layer of follicular cells; unilaminar primary, oocytes surrounded by a single layer of cuboidal fol-

licular cells already with the formation of the zona pellucida; primary multilaminar or pre-antral, follicles characterized by the enlargement of the granulosa cells, after the formation of the zona pellucida and theca interna; secondary or antral follicles, enlargement of the structures' sizes, presence of follicular fluid and antrum, theca interna and externa, organization of granulosa cells begins; mature, pre-ovulatory or Graafian follicles, presence of the antrum, organized granulosa cells, theca interna and externa, corona radiata and oophorus cumulus.

For the morphometric study, histological sections were observed and photomicrographed by light microscopy (Olympus BX61)⁶ using the Image J 1.45 software. For each follicular stage, ten follicles were examined (or what was available in two histological sections). The mean follicular and oocyte diameters were calculated between the measurement of the largest diameter (maximum diameter) and perpendicular diameter (minimum diameter) [Figure 1].

The relationship between follicle and oocyte were determined using the measurements of diameter, area and perimeter. For each follicular stage three linear equations were established, representing the patterns of follicle and oocyte growth, where x = follicular measurement and y = oocyte measurement.

The samples were 10% buffered formalin-fixed and processed for further paraffin embedding. The apoptotic cells were detected by a commercial kit for in situ detection of apoptosis (FragEL™ DNA Fragmentation Detection Kit Colorimetric - TdTEnzyme)⁸, Sections of 5 μ m mounted on sig-

nalized glass microscope slides were deparaffinized and washed in TBS (Tris-buffered saline). The sections were treated with Proteinase K for 8 min, and washed again in TBS. After washing, endogenous peroxidase was inactivated by using 9% hydrogen peroxide and then washed in TBS and incubated in equilibration buffer for 20 min at room temperature. Subsequently, the sections were incubated with the TdT enzyme (terminal deoxynucleotidyltransferase) at 37°C for one-half h in a moist chamber. The reaction was interrupted and the sections incubated, with conjugated antibody, in the moist chamber for 1 h. After washing with TBS, the reaction was detected using diaminobenzidine (DAB) for 30 min. The sections were counterstained with methylgreen during 10 min. Images were scanned by Image-ProPlus 7.0 software, and obtained through camera (Q Color 5 Olympus)⁶, attached to an Olympus BX43 microscope⁶. Cells were considered positive when TUNEL reaction was detected.

Data was submitted to analysis of variance for group comparison, with the means being compared by Tukey test. Data was tested for normality and homogeneity of variances and in non-normal distribution the Kruskal-Wallis test was performed for group comparison, followed by Dunn multiple comparison test. The correlation coefficient (r) of determination (r^2) and the regression equation were calculated for the selected variables [20]. The values were considered significantly different when $P < 0.05$. Statistical analysis was performed using the program "GraphPad version 3.10".




Figure 1. Maximum (mx) and minimum (mn) diameter of multilaminar primary follicle in domestic cat: (A) follicular diameter and (B) oocyte diameter.

RESULTS

The morphometric index (maximum and minimum diameter, oocyte and follicular area and perimeter) of 1039 follicles were evaluated, including primordial follicle (n = 318), unilaminar primary (n = 271), multilaminar primary (n = 271), secondary (n = 124) and pre-ovulatory (n = 55).

Primordial follicles in young animals showed larger diameter, follicular area and perimeter than the structures of adult queens, as well as the unilaminar primary follicles of the same group were larger compared with senile animals ($P < 0.05$). Comparing adult and younger queens, the first showed a significant decrease of oocyte diameter in primary and unilaminar primary follicles, as well as oocyte area and perimeter ($P < 0.05$). The values for follicular diameter, oocyte area and perimeter for multilaminar primary, secondary and pre-ovulatory structures did not present statistical differences between the groups ($P > 0.05$) [Tables 1, 2 & 3].

For the pre-ovulatory follicles there was no positive correlation between the oocyte growths regard-

ing the follicles ($P > 0.05$). For primordial follicles, unilaminar and multilaminar primary and secondary, a correlation was observed between the measurements of oocytes as to the follicles (Table 4).

A total of 1975 follicles were histologically prepared and evaluated. The apoptosis signs of primordial follicles (1664), unilaminar primary (120), multilaminar primary (134), secondary (41) and pre-ovulatory follicles (16) were studied using Tunel analysis. Positive apoptosis was identified in oocytes and pre-granulosa cells of primordial follicles and in other follicle stages positive apoptosis was observed in both granulosa and theca cells. Only in senile animals positive markers for apoptosis were identified in nuclei of primordial follicles (Figure 2).

No significant differences concerning the number of follicles and Tunel positive cells were observed between groups ($P > 0.05$). Table 5 shows the number of follicles and Tunel positive cells at each follicular stage of 18 queens according to their age.

Table 1. Mean values (x), standard error of the mean (SEM) and median (Md) of the diameter (µm) of follicle and oocyte for female cats according to age.

| Stage | G1 (young) | | G2 (adults) | | G3 (senile) | | |
|----------|------------|----------------|-------------|----------------|-------------|-----------------|---------|
| | x ± SEM | Med | x ± SEM | Med | x ± SEM | Med | |
| Follicle | P | 45.13 ± 0.71a | 45.25 | 41.51 ± 0.65b | 41.4 | 43.24 ± 0.66ab | 43.55 |
| | UP | 70.35 ± 1.35 | 70.30A | 64.43 ± 1.53 | 63.35B | 68.11 ± 3.1 | 62.25B |
| | MP | 127.60 ± 4.3 | 114.95 | 132.53 ± 5.3 | 121.3 | 128.51 ± 6.6 | 119.43 |
| | SE | 452.66 ± 19.46 | 423.3 | 400.51 ± 18.28 | 371.6 | 388.11 ± 22.05 | 395.7 |
| | OP | 893.18 ± 43.49 | 885.45 | 887.99 ± 37.63 | 930.4 | 1014.0 ± 100.99 | 1036.1 |
| Oocyte | P | 40.55 ± 0.56a | 40.15 | 37.57 ± 0.59b | 36.6 | 39.41 ± 0.57ab | 40.25 |
| | UP | 47.97 ± 0.89 | 47.1A | 43.00 ± 1.02 | 42.45B | 45.24 ± 1.51 | 43.45AB |
| | MP | 62.35 ± 1.56 | 59.8 | 65.77 ± 2.55 | 61.22 | 62.05 ± 2.36 | 58.72 |
| | SE | 112.80 ± 7.1 | 103 | 98.620 ± 10.72 | 91.55 | 88.70 ± 6.11 | 89.3 |
| | OP | 96.63 ± 3.56 | 94.35 | 101.35 ± 3.77 | 97.6 | 107.89 ± 12.95 | 116.7 |

Primordial (P), unilaminar (UP), multilaminar primary (MP), secondary (SE) and pre-ovulatory (OP); a, b: means followed by different small letters in line differ significantly by Tukey test ($P < 0.05$); A, B: means followed by different uppercase letters in line differ significantly by Dunn's test ($P < 0.05$).

Table 2. Mean values (x), standard error of the mean (SEM) and median (Md) of the area (µm²) of follicle and oocyte for female cats according to age.

| Stage | G1 (young) | | G2 (adults) | | G3 (senile) | | |
|----------|------------|-----------------|-------------|-----------------|-------------|------------------|---------|
| | x ± SEM | Med | x ± SEM | Med | x ± SEM | Med | |
| Follicle | P | 1941 ± 51.85a | 1908.8 | 1652.4 ± 46.89b | 1590.3 | 1776.9 ± 48.44ab | 1723.3 |
| | UP | 4043.9 ± 159.05 | 3810.3A | 3375.6 ± 165.22 | 3137.9B | 4185.8 ± 541.1 | 3086.7B |
| | MP | 14239 ± 1061.6 | 9807 | 15522 ± 1508.3 | 11130 | 15046 ± 1606.4 | 10759 |
| | SE | 169435 ± 14726 | 134712 | 137060 ± 13732 | 107104 | 126038 ± 13570 | 125592 |
| | OP | 672390 ± 63002 | 634085 | 647261 ± 54270 | 675397 | 921159 ± 170350 | 956801 |
| Oocyte | P | 1320.4 ± 35.92a | 1305.1 | 1134.3 ± 36.25b | 1036.7 | 1241.3 ± 32.81ab | 1270.2 |
| | UP | 1851.9 ± 67.46 | 1702.2A | 1497.0 ± 71.02 | 1341.9B | 1721.0 ± 138.22 | 1461.8B |
| | MP | 3225.1 ± 162.81 | 2782.2 | 3823.3 ± 351.1 | 2892.9 | 4061.6 ± 848.16 | 2669.6 |
| | SE | 10157 ± 1119.1 | 8336.8 | 6688.2 ± 548.44 | 5903 | 6311.8 ± 876.32 | 4949.4 |
| | OP | 7537.9 ± 529.42 | 7046.5 | 8518.6 ± 656.64 | 7721 | 10087 ± 2399.5 | 10333 |

Primordial (P), unilaminar (UP), multilaminar primary (MP), secondary (SE) and pre-ovulatory (OP); a, b: means followed by different small letters in line differ significantly by Tukey test ($P < 0.05$); A, B: means followed by different uppercase letters in line differ significantly by Dunn's test ($P < 0.05$).

Table 3. Mean values (x), standard error of the mean (SEM) and median (Md) of the perimeter (µm) of follicle and oocyte for female cats according to age.

| Stage | | G1 (young) | | G2 (adult) | | G3 (senile) | |
|----------|----|----------------|--------|-----------------|---------|-----------------|---------|
| | | x ± SEM | Med | x ± SEM | Med | x ± SEM | Med |
| Follicle | P | 157.24 ± 2.13a | 158 | 145.56 ± 2.02b | 144.55 | 150.63 ± 2.02ab | 150.9 |
| | UP | 272.82 ± 34.55 | 222.4A | 204.29 ± 4.86 | 200.9B | 217.01 ± 9.71 | 201.05B |
| | MP | 407.25 ± 13.65 | 364 | 424.68 ± 17.21 | 390.3 | 395.65 ± 20.52 | 348.5 |
| | SE | 1487.5 ± 63.62 | 1405.7 | 1534.4 ± 2227.5 | 1210.1 | 1253.9 ± 72.45 | 1274.1 |
| | OP | 2958.2 ± 144 | 2995.9 | 2940.0 ± 128 | 3092.7 | 3430.8 ± 355.48 | 3749.4 |
| Oocyte | P | 129.90 ± 1.81a | 129.5 | 120.58 ± 1.88b | 116.45 | 126.37 ± 1.67ab | 128.5 |
| | UP | 152.99 ± 2.73 | 150.3A | 135.76 ± 3.3 | 132.75B | 143.18 ± 4.82 | 137.9B |
| | MP | 198.53 ± 5.07 | 194.1 | 208.80 ± 8.15 | 194.9 | 197.29 ± 7.34 | 187.35 |
| | SE | 342.28 ± 16.62 | 329.2 | 279.81 ± 12.65 | 277.5 | 282.70 ± 18.48 | 284.6 |
| | OP | 311.63 ± 11 | 302.65 | 327.35 ± 12.41 | 323.4 | 344.74 ± 43.82 | 363.9 |

Primordial (P), unilaminar (UP), multilaminar primary (MP), secondary (SE) and pre-ovulatory (OP); a, b: means followed by different small letters in line differ significantly by Tukey test ($P < 0.05$); A, B: means followed of different uppercase letters in line differ significantly by Dunn's test ($P < 0.05$).

Table 4. Linear equations (x = follicular measurement and y = oocyte measurement) and r² values for follicular and oocyte growth of female cats according to age.

| Stage | Measure | Linear equations | r ² |
|--------------------------------|-----------|----------------------|----------------|
| Primordial follicles | Diameter | y = 0.7405x + 7.0833 | 0.7655 |
| | Area | y = 0.6598x + 49.767 | 0.8608 |
| | Perimeter | y = 0.8043x + 3.9923 | 0.8582 |
| Unilaminar primary follicles | Diameter | y = 0.506x + 11.301 | 0.7695 |
| | Area | y = 0.2702x + 657.73 | 0.7677 |
| | Perimeter | y = 0.0422x + 134.89 | 0.0887 |
| Multilaminar primary follicles | Diameter | y = 0.3033x + 24.166 | 0.549 |
| | Area | y = 0.1504x + 1401 | 0.2163 |
| | Perimeter | y = 0.3024x + 77.633 | 0.5464 |
| Secondary follicles | Diameter | y = 0.0782x + 70.059 | 0.0297 |
| | Area | y = 0.0369x + 2603.7 | 0.3275 |
| | Perimeter | y = 0.0248x + 270.62 | 0.0616 |

Table 5. Mean values (x), standard error of the mean (SEM) and median (Md) of follicles and apoptosis positive cells of primordial (P), unilaminar (UP) and multilaminar primary (MP), secondary (SE) and pre-ovulatory (OP) follicles of female cats in different age groups.

| Structure | Type | G1 (young) | | G2 (adult) | | G3 (senile) | |
|----------------|------|----------------|-----|--------------|------|---------------|-----|
| | | x ± SEM | Med | x ± SEM | Med | x ± SEM | Med |
| Follicles | P | 100.17 ± 12.44 | 95 | 89.5 ± 16.48 | 95.5 | 87.66 ± 11.52 | 87 |
| | UP | 8.66 ± 1.38 | 8.0 | 5.33 ± 1.97 | 3.0 | 6.00 ± 2.04 | 4.5 |
| | MP | 1.32 ± 0.54 | 7.0 | 3.86 ± 1.57 | 8.5 | 5.92 ± 2.41 | 7.0 |
| | SE | 2.66 ± 0.49 | 2.5 | 2.16 ± 0.30 | 2.0 | 2.00 ± 0.57 | 2.0 |
| | OP | 0.83 ± 0.30 | 1.0 | 1.33 ± 0.33 | 1.0 | 0.50 ± 0.34 | 0.0 |
| Positive cells | P | 2.00 ± 1.26 | 0.0 | 0.00 ± 0.00 | 0.0 | 5.16 ± 3.68 | 1.0 |
| | UP | 2.16 ± 1.64 | 0.0 | 0.00 ± 0.00 | 0.0 | 0.33 ± 0.33 | 0.0 |
| | MP | 4.00 ± 2.11 | 2.0 | 0.66 ± 0.66 | 0.0 | 0.66 ± 0.42 | 0.0 |
| | SE | 2.83 ± 1.81 | 0.5 | 2.16 ± 0.47 | 2.0 | 1.16 ± 1.16 | 0.0 |
| | OP | 4.16 ± 2.56 | 1.5 | 13.16 ± 5.88 | 8.5 | 2.50 ± 1.58 | 0.0 |

Significance level of 5%.




Figure 2. TUNEL positive cells in primordial follicles of senile domestic cat ovary (longitudinal section 200x; A - oocyte; B - pre-granulosa cells).

DISCUSSION

Considering the importance of this study for greater knowledge in the basic aspects for reproductive biotechnologies, we verified that secondary follicles showed the largest diameters and younger animals the largest values for diameter, area and perimeter, suggesting that this age group could be ideal for the use and manipulation of oocytes. These characteristics were also observed in female dogs by literature [5], who evaluated the effects of age and oocyte size upon subsequent oocyte maturation (IVM). They verified that oocytes smaller than 100 μm in diameter had less nuclear material and meiotic competence and younger bitches showed optimized oocyte maturation.

Age had an important effect on oocyte morphometry in queens, given that remaining follicles and their respective oocytes decrease in size with the process of ageing. This fact is justified through the morphometric parameters of unilaminar primary follicles, which are smaller in older cats compared with the younger group (diameter - G1: 70.30 μm ; G3: 62.25 μm) (area - G1: 3810.3 μm^2 ; G3: 3086.7 μm^2) (Perimeter- G1: 222.4 μm ; G3: 201.05 μm). This pattern was also observed in women [17], who found a reduction in mean follicular diameter of primordial and primary follicles in women older than 36 years (39.4 μm) in comparison to younger groups (20 to 27 years: 41.9 μm ; 27 to 36 years: 41.5 μm).

As described by literature [11], the oocyte and follicle growth in cats showed a biphasic pattern. During the first stage of follicular development, a significant correlation between follicle and oocyte growth was verified, resulting in a linear equation $y = 0.3048x$

$+ 25.018$, $r^2 = 0.7239$ (x = follicular diameter y = oocyte diameter). After antral formation and development of secondary and pre-ovulatory follicles, the oocyte growth pattern significantly decreased compared to follicular growth, resulting in a new linear correlation: $y = 0.0072x + 98.001$, $r^2 = 0.0015$. These results are superior to those described in cats [11].

In adult animals, the mean follicular diameter was lower in primordial follicles, unilaminar and multilaminar primary follicles and higher in secondary or antral follicles (41.51 μm , 64.43 μm , 132.53 μm and 400.51 μm , respectively), when compared to the results of literature [11] (44.3 μm , 86.2 μm , 155.6 μm and 223.8 μm , respectively). In contrast, the results in this study were superior when comparing to other works, where queens showed a decrease in the mean diameter of primordial follicles, unilaminar primary and secondary follicles of 28.3 μm , 41 μm and 74.6 μm , respectively [3], and 38.8 μm , 63.9 μm and 98.7 μm , respectively [1]. Different values were also observed in the mean follicular diameter of other species such as adult sheep, in which primordial follicles showed a mean of 39.4 μm [9] and bitches with values for primordial, unilaminar and multilaminar primary follicles of 44.3 μm , 50.7 μm and 148.9 μm , respectively [1].

When we compared our findings with the results of other species we observed that the mean follicular diameter was higher in primordial follicles, unilaminar and multilaminar primary follicles (45.13 μm , 70.35 μm and 127.60 μm) than diameters obtained in other species as in pre-pubertal gilts (33.8 μm , 40.4 μm and 84.5 μm) [13] and young women (< 13 years) (39 μm and 44.1 μm) [17]; the mean follicular

diameter of older queens was higher in primordial and unilaminar primary follicles (43.24 μm and 68.11 μm , respectively) compared to post-pubertal female buffaloes (7 - 10 years) (35 μm and 41.8 μm , respectively) [10]; the mean oocyte diameter in young animals was higher in primordial, unilaminar and multilaminar primary follicles (40.55 μm , 47.97 μm and 62.35 μm , respectively) when comparing with pre-pubertal gilts (26 μm , 27.3 μm and 39.1 μm) [13] and women (< 13 years) with values for primordial and unilaminar primary follicles (34.3 μm and 41 μm , respectively) [17].

The process of follicular atresia is characterized by the occurrence of apoptosis, or programmed cell death when the organism begins to efficiently eliminate dysfunctional cells [12]. The study of follicular apoptosis in small animals, especially in cats, is very important for the development of reproduction biotechnologies.

The TUNEL technique is widely used to detect DNA fragmentation as an indicative factor of apoptosis. Cells that had suffered apoptosis were detected in human embryos using the chromatin condensation state and DNA fragmentation, which was subsequently confirmed by TUNEL [8]. To our knowledge, this is the first study evaluating the frequency of apoptosis with the TUNEL assay in ovarian tissue in queens according to their age. The TUNEL method was effective labeling oocytes of primordial follicles and pre-granulosa cells in older cats and multilaminar primary and secondary follicles or antral follicles in granulosa cells of young animals.

In our study, there was no significant difference of apoptosis in the different age groups, indicating that none of the older, adult or younger queens showed an increase or decrease of apoptosis. These results in felines are contrary to those verified by literature [19], who demonstrated that apoptosis rate of human oocytes was significantly higher in a group of older women (41 - 50 years) compared to younger women (21 to 40 years).

CONCLUSION

Young queens presented larger values for unilaminar primary follicles than older animals. We suggest that the pool of remaining small follicles and their respective oocytes decrease with age and in senile phase these follicles are of lower quality, when compared with the beginning of reproductive life. On the other hand, the phenomenon of apoptosis showed no relationship with age in animals, occurring in a physiological, continuous and proportionate manner considering the number of non-dominant follicles involved in each estrous cycle.

The results of this study regarding histomorphometry, morphology and apoptosis of ovarian tissue may contribute as a model for the study of physiological parameters or set up of assisted reproductive technologies of endangered wild cats.

MANUFACTURERS

¹Vetnil Indústria e Comércio de Produtos Veterinários. Louveira, SP, Brazil.

²Cristália Produtos Químicos Farmacêuticos. São Paulo, SP, Brazil.

³Zoetis Indústria de Produtos Veterinários Ltda. Campinas, SP, Brazil.

⁴Hertape Calier. Juatuba, MG, Brazil.

⁵Laboratório Aids Pty. Narrabeen, Australia.

⁶Olympus Optical do Brasil. São Paulo, SP, Brazil.

⁷Coat-A-Count Diagnostic Products Cooperation. Los Angeles, CA, USA.

⁸Merck Millipore. Darmstadt, Germany.

Acknowledgements. The authors would like to thank FAPESP for the financial support (grants 2011/18506-2 & 2011/18550-1) and Professor Anilton César Vasconcelos for providing his lab and Lorinaldo Lopes de Moraes and Marta Paccanaro Peres for performing histological stain and slides.

Ethical Approval. The trial was conducted following approval by the Institutional Animal Care and Use Committee of the UNESP Araçatuba, SP, Brazil (protocol number 00717-2012).

Declaration of interest. The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

REFERENCES

- 1 Barber M.R., Lee S.M., Steffens W.L., Ard M. & Fayrer-Hosken R.A. 2001. Immunolocalization of zona pellucid antigens in the ovarian follicle of dogs, cats, horses and elephants. *Theriogenology*. 55: 1705-1717.
- 2 Blondin P., Dufour M. & Sidard M.A. 1996. Analysis of atresia in bovine follicles using different methods: flow cytometry, enzyme-linked immunosorbent assay, and classic histology. *Biology of Reproduction*. 54: 631-637.
- 3 Carrijo Júnior O.A., Marinho A.P., Campos A.A., Amorim C.A. & Bão S.N. 2010. Morphometry, estimation and ultrastructure of ovarian preantral follicle population in queens. *Cells Tissues Organs*. 191: 152-160.

- 4 Feranil J.B., Isobe N. & Nakao T. 2005.** Apoptosis in the antral follicles of swamp buffalo and cattle ovary: tunel and caspase-3 histochemistry. *Reproduction in Domestic Animals*. 40: 111-116.
- 5 Hewitt D.A. & England G.C.W. 1998.** The effect of oocyte size and bitch age upon oocyte nuclear maturation *in vitro*. *Theriogenology*. 49: 957-966.
- 6 Ishizaki M.N., Nascimeno A.A., Kaneto C.N., Montano T.R.P., Perri S.H.V., Vasconcelos R.O. & Bresciani K.D.S. 2006.** Frequência e Intensidade parasitária de Helminthos gastrintestinais em felinos da zona urbana do município de Araçatuba-Sp. *ARS Veterinaria*. 22: 212-216.
- 7 Junqueira L.C. & Carneiro J. 2011.** Aparelho Reprodutor Feminino. In: Junqueira L.C. & Carneiro J. (Eds). *Histologia básica*. Rio de Janeiro: Guanabara Koogan, pp.432-439.
- 8 Jurisicov A., Varmuza S. & Casper R.F. 1996.** Programmed cell death and human embryo fragmentation. *Molecular Human Reproduction*. 2: 93-98.
- 9 Lundy T., Smith P., O'Connell A., Hudson N.L. & Mc Natty K.P. 1999.** Populations of granulose cells in small follicles of the sheep ovary. *Journal of Reproduction and Fertility*. 115: 251-262.
- 10 Mondadori R.G., Luque M.C.A., Santin T.R. & Bão SN. 2007.** Ultrastructural and morphometric characterization of buffalo (*Bubalus bubalis*) ovarian preantral follicles. *Animal Reproduction Science*. 97: 323-333.
- 11 Reynaud K., Gicquel C., Thoumire S., Ficheux C., Bestandji M. & Chastant- Maillard S. 2009.** Folliculogenesis and morphometry of oocyte and follicle growth in the feline ovary. *Reproduction in Domestic Animals*. 44: 174-179.
- 12 Robbins S.L., Collins T., Cotran R.S. & Kumar V.M.D. 2004.** *Patologia Estrutural e Funcional*. 6th edn. Rio de Janeiro: Elsevier, 1251p.
- 13 Silva R.C., Bão S.N., Jivago J.L.P.R. & Lucci C. 2011.** Ultrastructural characterization of porcine oocytes and adjacent follicular cells during follicle development: Lipid component evolution. *Theriogenology*. 76: 1647-1657.
- 14 Tilly J.L., Billig H., Kowalski K.I. & Hsueh A.J.W. 1992.** Epidermal growth factor and basic fibroblast growth factor suppress the spontaneous onset of apoptosis in cultured rat ovarian granulose cells and follicles by a tyrosine kinase-dependent mechanism. *Molecular Endocrinology*. 6: 1942-1950.
- 15 Toniollo G.H., Cury S.R., Vicente W.R.R., Camacho A.A., Garcia J.M. & Vantini R. 1995.** Colpocitologia do ciclo estral em gatas. *Brazilian Journal of Veterinary Research and Animal Science*. 32: 125-129.
- 16 Uchikura K., Nagano M. & Hishinuma M. 2011.** The effect of ovarian status and follicular diameter on maturational ability of domestic cat oocytes. *The Journal of Veterinary Medical Science*. 73: 561-566.
- 17 Westergaard C.G., Byskov A. & Andersen C.Y. 2007.** Morphometric characteristics of the primordial to primary follicle transition in the human ovary in relation to age. *Human Reproduction*. 22: 2225-2231.
- 18 Wlodarczyk R., Bukowska M., Jackwska M., Mucha S. & Jaskowski J.M. 2009.** *In vitro* maturation and degeneration of domestic cat oocytes collected from ovaries temperatures. *Veterinary Medicine - Czech*. 54: 491-497.
- 19 Wu J., Zhang L. & Wang X. 2000.** Maturation and apoptosis of human oocytes *in vitro* are age-related. *Fertility and Sterility*. 74: 1137-1141.
- 20 Zar J.H. 1999.** *Biostatistical analysis*. 4th edn. New Jersey: Prentice-Hall, 930p.