



Journal of Human Sport and Exercise

E-ISSN: 1988-5202

jhse@ua.es

Universidad de Alicante

España

RYBÁROVÁ, SILVIE; NOVOTNÝ, JAN

Skin temperature changes of muscle regions in training swimmers

Journal of Human Sport and Exercise, vol. 10, núm. 1, 2015, pp. S192-S197

Universidad de Alicante

Alicante, España

Available in: <http://www.redalyc.org/articulo.oa?id=301043404005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Skin temperature changes of muscle regions in training swimmers

SILVIE RYBÁŘOVÁ , JAN NOVOTNÝ

Faculty of Sport Studies, Masaryk University, Brno, Czech Republic

## ABSTRACT

Rybářová, S., & Novotný, J. (2015). Skin temperature changes of muscle regions in training swimmers. *J. Hum. Sport Exerc.*, 9(Proc1), pp.S192-S197. Purpose: Our aim was to detect changes in infrared radiation, overloaded structures of the musculoskeletal system at the main part of the shoulder girdle and upper body with the help of elite swimmers of Kometa Brno. Methods: First measurement was done before training and second 15 minutes after training in water in the swimming pool. The group consisted of seven Czech national swimmers. Athletes participated in testing during six months. Every measurement contains four positions. Front and back side, right and left side. Every athlete went through 13 measurements. We directed infrared thermograph camera FlukeTiR at 10 muscular groups that are most used in swimming. Besides we have form about training with kilometres, other exercise out of water, competitions, illness and be absent at training and other pain muscles, ligaments etc. Results: We have analysed all temperatures only of one swimmer. Here was significant increased temperature after swimming only in deltoideus anterior at right side (from  $33.4 \pm 1.02^{\circ}\text{C}$  to  $34.0 \pm 0.69^{\circ}\text{C}$ ). The other temperature was significant decreased: muscles groups of pectoralis major and minor (right side from  $33.6 \pm 0.92$  to  $33.1 \pm 0.61^{\circ}\text{C}$ ; left side from  $33.8 \pm 0.82$  to  $33.1 \pm 0.69^{\circ}\text{C}$ ), latissimus dorsi and erector spinae - pars lumbalis at both sides together. Conclusion: Five of twenty muscle regions have notable tendencies of increased temperatures, including a front part of deltoideus at right side which is very active by spreading arm forward and beginning of swimming stroke. In nine areas, including the main agonist for swimmers movement forward - triceps brachii, we found out only no significant lowering of temperatures. That was caused by cooling of the swimmer in the water. We have next six swimmers to analysis. **Key words:** SWIMMING, INFRARED THERMOGRAPHY, MUSCULAR WORK.



**Corresponding author.** Žerotínovo nám. 617/9, 601 77 Brno, Republika Čechy

E-mail: 160531@mail.muni.cz

9th INSHS International Christmas Sport Scientific Conference, 4-6 December 2014. International Network of Sport and Health Science. Szombathely, Hungary.

JOURNAL OF HUMAN SPORT & EXERCISE ISSN 1988-5202

© Faculty of Education. University of Alicante

doi:10.14198/jhse.2015.10.Proc1.05

## INTRODUCTION

A very good method to measure individual muscle activity is electromyography. This method provides information on electric muscular activity which is prior to metabolic activity itself. In recent years, results of electromyographical studies have been published on muscular activity during swimming (Conceição et al., 2013; Lauer et al., 2013).

Scientific articles on thermal changes as a result of swimming in water are rare. (Wade & Veghte, 1977) observed four professional swimmers after a 500-meter free style swim in water of 23.5 °C. (Zaidi, Taiar, Fohanno & Polidori, 2007) described the temperature changes in one swimmer. The swimmer used in turns all four swimming styles. Each for 1 minute and there were 10-minute breaks between them. They defined the temperature of large body segments and they did not evaluate the areas corresponding to particular muscles. For example, dorsal arm included dorsal deltoid muscles and triceps.

We want to help with solution measuring and show metabolic activities in progress and muscles groups on swimming. For measuring our dates we have used infrared thermography. Our aim was to detect changes in infrared radiation, overloaded structures of the musculoskeletal system at the main part of the shoulder girdle and upper body with the help of elite swimmers of Kometa Brno.

Than find out if there is some new thermally interesting areas before and after training in local area or hole muscle.

Purpose is to find if thermograph can find or prevent injury of movement musculoskeletal system. Healthy is key to the top athletes.

## MATERIAL AND METHODS

### *Athletes group*

Over group consisted of seven Czech national swimmers of Kometa Brno. The group are 5 men and 2 women, from 20 to 34 years old. They had 11 weeks of training in water and 3 units of dry land. Training in water is around 50-70km per week. It is around 2000 km per year. They are specializing in freestyle for long (5-25 km) and short distance (50 m, 100 m).

We chose one swimmer for a complete analysis of measurement. He is 24 years old, 183 cm high and 76 kg weight. His main style is free-style in open water. Mains discipline is 5 km and 10k m. He is Czech national swimmer and he was at European and World championships. He had this year 2200 km per year. He started with swimming 16 years ago.

### *Measurement*

1. Measurement was done before training.
2. 15 minutes after training

Training was in water in the swimming pool. There was temperature of water 27°C. One unit was 2 hour long and measuring was done regularly in training since 6 to 8 in the morning. Before first measuring was the body 15 minutes nude a after training the swimmer has use the towel and sit for 15 minutes in the pool. Athletes participated in monitoring during six months. Measurements were made from January to June. Every athlete went through 13 measurements. We have 104 pictures of one swimmer. Every measurement contains four positions. Front and back side, right and left side.




Figure 1. Thermograms of front side and back side




Figure 2. Thermograms of right side and left side

We directed infrared thermograph camera FlukeTiR at 10 muscular groups that are most used in swimming.

Camera is hand-held, infrared spectral band 7.5  $\mu\text{m}$  to 14  $\mu\text{m}$  portable, 23 x 17  $^{\circ}\text{C}$  lens, manual focus, sensitivity of 0.1  $^{\circ}\text{C}$ , LCD display with a resolution of 480 x 640 pixels. At the display there was the temperature distribution of body surface.



Figure 3. Camera FlukeTiR

We measured the skin infrared radiation of selected 10 muscle groups of the upper body for swimmers that are most involved in the movement in the water.

Table 1. Selected regions of interest (ROI)

ROI	Muscles, their parts or groups
Da	m. deltoideus - pars anterior
Dp	m. deltoideus – pars posteriori
DI	m. deltoideus – pars lateralis
Bb	m. biceps brachii
Tb	m. triceps brachii
Ts	m. trapezius – pars superior
P	m. pectoralis major et minor
R-Ti	m. rhomboideus major et minor, and m. trapezius – pars inferior
Ld	m. latissimus dorsi
Esl	m. erector spinae - pars lumbalis

## RESULTS OF SELECTED SWIMMER

Table 2. Temperatures of selected regions

ROI	Left side				Right side			
	Before		After		Before		after	
	X	S	X	S	X	S	X	S
Da	33.6	0.93	34.0	0.56	33.4	1.02	34.0	0.69
Dp	33.8	1.05	34.2	0.62	33.9	0.96	34.1	0.70
DI	33.8	0.78	33.5	0.79	33.9	0.77	33.6	0.77
Bb	33.7	0.92	33.5	0.61	33.6	1.00	33.2	0.68
Tb	32.7	0.90	32.6	0.65	32.7	1.03	32.8	0.65
Ts	34.2	0.73	34.1	0.62	34.4	0.74	34.3	0.62
P	33.8	0.82	33.1	0.69	33.6	0.92	33.1	0.61
R-Ti	34.3	0.77	34.1	0.84	34.4	0.73	33.9	0.77
Ld	33.9	0.61	32.9	0.70	34.2	0.52	33.1	0.43
Esl	34.5	0.49	32.7	0.86	34.5	0.47	32.8	0.98

Key: x-average; s-the standard deviation

We have analysed all temperatures only of one swimmer. Here was significant increased temperature after swimming only in deltoideus anterior at right side (from  $33.4 \pm 1.02^{\circ}\text{C}$  to  $34.0 \pm 0.69^{\circ}\text{C}$ ) (Fig. 4.). The other temperature was significant decreased: muscles groups of pectoralis major and minor (right side from  $33.6 \pm 0.92$  to  $33.1 \pm 0.61^{\circ}\text{C}$ ; left side from  $33.8 \pm 0.82$  to  $33.1 \pm 0.69^{\circ}\text{C}$ ) (Fig. 5.), latissimus dorsi and erector spinae - pars lumbalis at both sides together.




Figure 4. Temperature before and after swimming in region of deltoideus anterior at right side




Figure 5. Temperature before and after swimming in region of pectoralis major and minor at left side

## DISCUSSION

Our swimmer has only warming on his muscles deltoideus. He is right handed, and it is visible on his hand work in the water. Stroke is stronger and the hand is not so much lose. You can see it on his technique. Deltoideus muscle is really active in use under water as up the water stroke.

We haven't found any change of body temperature during hole measuring, the swimmer have no problems or pain.

Other muscular parts and groups have degrees of temperature. That is because cold water and air is colder than the body temperature. We had the other measurement, but it was so early after training and at skin was too much water. The measurement has been canceled. It is not applicable.




Figure 6. Skin with water at front and back side after training

## CONCLUSIONS

We got results by measuring only one person. Five of twenty muscle regions have notable tendencies of increased temperatures, including a front part of deltoideus at right side which is very active by spreading arm forward and beginning of swimming stroke. In nine areas of main agonist swimmers movement forward included – triceps brachii, we have find out only no significant lowering temperatures. That because cooling of the swimmer in the water. We haven't found any source or areas showing injury or damage muscle system. We need to analyze more swimmers for better results. We have next six swimmers to analysis.

## REFERENCES

1. Conceição, A., Silva, A., Barbosa, T.M., & Louro, H. (2013). Observation and technical characterization in swimming: 200 m breaststroke. *Revista Brasileira de Medicina do Esporte*, 19(1), pp.56-61.
2. Lauer, J., Figueiredo, P., Vilas-Boas, J.P., Fernandes, R.J., & Rouard, A.H. (2013). Phase-dependence of elbow muscle coactivation in front crawl swimming. *Journal of Electromyography & Kinesiology*, 23(4), pp.820-826.
3. Wade, C.E., & Veghte, J.H. (1977). Thermographic evaluation of the relative heat loss by area in man after swimming. *Aviation, Space and Environmental Medicine*, 48, pp.16-18.
4. Zaidi, H., Taiar, R., Fohanno, S., & Polidori, G. (2007). The influence of swimming type on the skin-temperature maps of a competitive swimmer from infrared thermography. *Acta Bioengineering and Biomechanics*, 9(1), pp.47-51.