

Acta Scientiarum. Technology

ISSN: 1806-2563

eduem@uem.br

Universidade Estadual de Maringá  
Brasil

Saoud, Mahmoud

Domination numbers of the complete grid graphs  $P_k \times P_n$

Acta Scientiarum. Technology, vol. 34, núm. 3, julio-septiembre, 2012, pp. 321-324

Universidade Estadual de Maringá

Maringá, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=303226542010>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



# Domination numbers of the complete grid graphs $P_k \times P_n$

Mahmoud Saoud

Ecole Normale Supérieure, Département de mathématiques, BP 92 Kouba, 16050, Alger, Algérie. E-mail: saoud\_m@yahoo.fr

**ABSTRACT.** This paper concerns the domination numbers  $\gamma(P_k \times P_n)$  for the complete  $P_k \times P_n$  grid graphs for  $k = 7, 8, 9$  and for all  $n \geq 1$ . These numbers were previously established (BONDY; MURTY, 2008; CHANG; CLARK, 1993). Here we present dominating sets by other method.

**Keywords:** dominating Set, domination number, transformation of a dominating set, cartesian product of two paths.

## Números de dominação de gráficos *grid* completos $P_k \times P_n$

**RESUMO.** Este artigo diz respeito aos números de dominação  $\gamma(P_k \times P_n)$  para gráficos completos  $P_k \times P_n$ , para  $k = 7, 8, 9$  e para todo  $n \geq 1$ . Estes números foram previamente estabelecidos por (BONDY; MURTY, 2008; CHANG; CLARK, 1993). Aqui apresentamos conjuntos dominados por outro método.

**Palavras-chave:** conjunto de dominação, número de dominação, transformação de um conjunto dominante, produto cartesiano de dois caminhos.

### Introduction

A dominating set in a graph is a set of vertices having the property that every vertex not in the set is adjacent to a vertex in the set. The domination number  $\gamma(G)$  of a graph  $G$  is the cardinality of a smallest dominating set in  $G$ .

The problem of finding the domination number of a arbitrary grid graph (= subgraph of  $P_k \times P_n$ ) is NP-complete (CLARK et al., 1990).

In this paper, we introduce the concept of transforming the domination from a vertex in a dominating set  $D$  of a graph  $G = (V, E)$  to a vertex in  $V - D$ , where  $G$  is a simple connected graph. We give an algorithm using this transformation to obtain a dominating set of a graph  $G$ .

A graph  $G = (V, E)$  is a mathematical structure which consists of two sets  $V$  and  $E$ , where  $V$  is finite and nonempty, and every element of  $E$  is an unordered pair  $\{u, v\}$  of distinct elements of  $V$ ; we simply write  $uv$  instead of  $\{u, v\}$ .

The elements of  $V$  are called vertices, while the elements of  $E$  are called edges (BONDY; MURTY, 2008).

Two vertices  $u$  and  $v$  of a graph  $G$  are said to be adjacent if  $uv \in E$ .

The neighborhood of  $v$  is the set of all vertices of  $G$  which are adjacent to  $v$ ; the neighborhood of  $v$  is denoted by  $N(v)$ . The closed neighborhood of  $v$  is  $\bar{N}(v)$ ,  $\bar{N}(v) = N(v) \cup \{v\}$ .

The degree  $d(v)$  of a vertex  $v$  is the cardinality  $|N(v)|$ ,  $d(v) = |N(v)|$ .

### Material and methods

#### Definitions

Let  $D$  be a dominating set of a graph  $G = (V, E)$ .

1. We define the function  $C_D$ , which we call the weight function, as follows:  $C_D: V \rightarrow N$ , where  $N$  is the set of natural numbers,  $C_D(v) = |\tilde{N}(v)|$ , where  $\tilde{N}(v) = \{w \in D: vw \in E \text{ or } w = v\}$ .

i.e. the weight of  $v$  is the number of vertices in  $D$  which dominate  $v$ .

2. We say that  $v \in D$  has a moving domination if there exists a vertex  $w \in N(v) - D$  such that  $wu \in E$  for every vertex  $u$ ,  $u \in \{x \in N(v): C_D(x) = 1\}$ .

3. We say that a vertex  $v \in D$  is a redundant vertex of  $D$  if  $C_D(w) \geq 2$  for every vertex  $w \in \bar{N}(v)$ .

4. If  $v \in D$  has a moving domination, we say that  $v$  is inefficient if transforming the domination from  $v$  to any vertex in  $N(v)$  would not produce any redundant vertex.

#### Complete grid graph $P_k \times P_n$ :

For two vertices  $v_o$  and  $v_n$  of a graph  $G$ , a  $v_o - v_n$  walk is an alternating sequence of vertices and edges  $v_o, e_1, v_1, \dots, e_n, v_n$  such that consecutive vertices and edges are incident.

A path is a walk in which no vertex is repeated. A path with  $n$  vertices is denoted by  $P_n$ , it has  $n - 1$  edges; the length of  $P_n$  is  $n - 1$ ; the cartesian product  $P_k \times P_n$  of two paths is the complete grid graph with vertex set  $V = \{(i, j): 1 \leq i \leq k, 1 \leq j \leq n\}$

where  $(u_1, u_2) (v_1, v_2)$  is a edge of  $P_k \times P_n$  if  $|u_1 - v_1| + |u_2 - v_2| = 1$  (CHANG; CLARK, 1993).

If  $D$  is a dominating set of  $P_k \times P_n$  which has no redundant vertex, then a vertex  $v \in D$  has a moving domination if and only if one of the following two cases occurs:

Case (1): for every vertex  $w \in N(v)$ , we have  $C_D(w) \geq 2$ .

In this case, the domination of  $v$  can be transformed to any vertex in  $N(v) - D$ .

Case(2): there exists exactly one vertex  $u \in N(v)$  such that  $C_D(u) = 1$  in this case, the domination of  $v$  can be transformed only to  $u$ .

#### An algorithm for finding a dominating set of a graph $P_k \times P_n$ using a transformation of domination of vertices

1. Let  $P_k \times P_n = (V, E)$  be a graph of order greater than 1;  $|V| = m$ .

2. Let  $D = V$  be a dominating set of  $P_k \times P_n$ , then, for any vertex  $v \in D$  we have  $C_D(v) = d(v) + 1 \geq 2$ .

3. Pick a vertex  $v_1$  of  $D$ , and delete from  $D$  all vertices  $w, w \in N(v_1)$  then, for  $1 < n < \frac{m}{2}$ , pick a vertex  $v_n \in D - \bigcup_{i=1}^{n-1} \overline{N}(v_i)$  and delete from  $D$  all vertices  $w, w \in N(v_n) - \bigcup_{i=1}^{n-1} \overline{N}(v_i)$ .

4. If  $D$  contains a redundant vertex, then delete it.

Repeat this process until  $D$  has no redundant vertex.

5. Transform domination from vertices of  $D$  which have moving domination to vertices in  $V - D$  to obtain redundant vertices and go to step 4.

If no redundant vertex can be obtained by a transformation of domination of vertices of  $D$ , then stop, and the obtained dominating set  $D$  satisfies:

for every  $v \in D, \exists w \in \overline{N}(v)$  such that  $C_D(w) = 1$ .

#### Example

1. Let  $(k, n)$  be the vertex in the  $k$ -th row and in the  $n$ -th column of the graph  $G = P_7 \times P_{16}; |V| = 112$ .

2. Let  $D = V$ , dominating set of  $G$ .

3. Pick a vertex  $v_1 = (2, 1) \in D$ , and delete from  $D$  all vertices  $w, w \in N(v_1)$ , then, for  $1 < n < \frac{112}{2}$ , pick a vertex  $v_n, v_n \in D - \bigcup_{i=1}^{n-1} \overline{N}(v_i)$ , and delete from  $D$  all vertices  $w$ ,

$w \in N(v_n) - \bigcup_{i=1}^{n-1} \overline{N}(v_i)$ . We obtain the dominating set  $D$  (black circles) in figure 1.





Figure 1. Dominating set (black circles).

4. Since for every vertex  $v \in D, \exists w \in \overline{N}(v)$  such that  $C_D(w) = 1$ ,  $D$  has no redundant vertices.

5. Transform the domination from the vertex (6, 16) to the vertex (5, 16) and delete, from  $D$ , the resulting redundant vertex (5, 15).

Therefore, the set  $D$  indicated in figure 2 (black circles) is a dominating set of  $G = P_7 \times P_{16}$ .


Note that  $D$  minimum dominating set (see CHANG et al., 1994)  $\gamma(P_7 \times P_{16}) = 27$ .




$$\begin{aligned} \gamma(P_7 \times P_{16}) &= 27 \\ &= 2(n - 5) + 5 \\ &= 2n - 5 \end{aligned}$$

Figure 2. Domination number (black circles).

And so, we gradually get domination numbers of  $P_7 \times P_n$ .





**Table 1.** Domination numbers  $\gamma(P_7 \times P_n), n \geq 1$ .


Hence:

$$\gamma(P_7 \times P_n) = \begin{cases} 3 & \text{for } n = 1 \\ 4 & \text{for } n = 2 \\ 6 & \text{for } n = 3 \\ 2n - t & \text{for } n = 3t + 1, 3t + 2, 3t + 3; t \geq 1 \end{cases}$$


where  $t$  is a positive integer.

#### Dominating sets for $P_8 \times P_n$ Grid graph


By self-method, we give in Table 2,  $\gamma(P_8 \times P_n), n \geq 1$ .




And by gradually, we find that:  $\gamma(P_8 \times P_n) = 2n$  for  $6 \leq n \leq 14$ .




And by gradually, we find that  $\gamma(P_8 \times P_n) = 2n - 1$  for  $15 \leq n \leq 22$ .



And by gradually, we find that  $\gamma(P_8 \times P_n) = 2n - 2$  for  $23 \leq n \leq 30$ .



And by gradually, we find that  $\gamma(P_8 \times P_n) = 2n - 3$  for  $31 \leq n \leq 38$ .



And by gradually, we find that  $\gamma(P_8 \times P_n) = 2n - 4$  for  $39 \leq n \leq 46$ .

Hence:


$$\gamma(P_8 \times P_n) = \begin{cases} 2n + 1 & \text{for } n = 1, 2, 3, 5 \\ 2n & \text{for } n = 4 \text{ or } 6 \leq n \leq 14 \\ 2n - (t + 1) & \text{for } n = 15 + 8t, 16 + 8t, \dots, 22 + 8t; t \geq 0 \end{cases}$$

where  $t$  is a integer.

**Table 2.** Domination numbers  $\gamma(P_8 \times P_n), n \geq 1$ .

### Dominating sets for $P_9 \times P_n$ Grid graph


By self-method, we give in Table 3,  $\gamma(P_9 \times P_n), n \geq 1$



$n = 12$

$$\begin{aligned}\gamma(P_9 \times P_{12}) &= 26 \\ &= 2(n - 8) + 18 \\ &= 2n + 2\end{aligned}$$


And by gradually, we find that  $\gamma(P_9 \times P_n) = 2n + 2$  for  $4 \leq n \leq 12$ .



$n = 13$

$$\begin{aligned}\gamma(P_9 \times P_{13}) &= 29 \\ &= 2(n - 6) + 15 \\ &= 2n + 3\end{aligned}$$

And by gradually, we find that  $\gamma(P_9 \times P_n) = 2n + 3$  for  $13 \leq n \leq 23$ .



$n = 24$

$$\begin{aligned}\gamma(P_9 \times P_{24}) &= 52 \\ &= 2(n - 13) + 30 \\ &= 2n + 4\end{aligned}$$

And by gradually, we find that  $\gamma(P_9 \times P_n) = 2n + 4$  for  $24 \leq n \leq 34$ .

Hence:

$$\gamma(P_9 \times P_n) = \begin{cases} 2n + 1 & \text{for } 1 \leq n \leq 3 \\ 2n + 2 & \text{for } 4 \leq n \leq 12 \\ 2n + (t + 3) & \text{for } n = 13 + 11t, 14 + 11t, \dots, 23 + 11t; t \geq 0 \end{cases}$$

where  $t$  is a integer.

**Table 3.** Domination numbers  $\gamma(P_9 \times P_n), n \geq 1$ .

We plan to deal with  $P_k \times P_n$  grid graphs, for  $n \geq 1$  and  $k \geq 10$ .

### Results and discussion

$$\gamma(P_7 \times P_n) = \begin{cases} 3 & \text{for } n = 1 \\ 4 & \text{for } n = 2 \\ 6 & \text{for } n = 3 \\ 2n - t & \text{for } n = 3t + 1, 3t + 2, 3t + 3; t \geq 1 \end{cases}$$

**Table1.** Domination numbers  $\gamma(P_7 \times P_n), n \geq 1$ .

$$\gamma(P_8 \times P_n) = \begin{cases} 2n + 1 & \text{for } n = 1, 2, 3, 5 \\ 2n & \text{for } n = 4 \text{ or } 6 \leq n \leq 14 \\ 2n - (t + 1) & \text{for } n = 15 + 8t, 16 + 8t, \dots, 22 + 8t; t \geq 0 \end{cases}$$

**Table2.** Domination numbers  $\gamma(P_8 \times P_n), n \geq 1$ .

$$\gamma(P_9 \times P_n) = \begin{cases} 2n + 1 & \text{for } 1 \leq n \leq 3 \\ 2n + 2 & \text{for } 4 \leq n \leq 12 \\ 2n + (t + 3) & \text{for } n = 13 + 11t, 14 + 11t, \dots, 23 + 11t; t \geq 0 \end{cases}$$

**Table3.** Domination numbers  $\gamma(P_9 \times P_n), n \geq 1$ .

### Conclusion

From these results, we note that it is possible to calculate the domination numbers quickly and without the need of a calculator in most cases, however the results published previously, you must necessarily use a calculator.

We tried this method on large grid graphs, and we obtained similar results quickly

### References

- BONDY, J. A.; MURTY, U. S. R. **Graph theory**, Springer, 2008.
- CHANG, T. Y.; CLARK, W. E. The domination numbers of  $5 \times n$  and  $6 \times n$  grid graphs. **Journal of Graph Theory**, v. 17, n. 1, p. 81-107, 1993.
- CHANG, T. Y.; CLARK, W. E.; HARE, E. O. Domination numbers of complete grid graph, I. **Ars Combinatoria**, v. 38, p. 97-111, 1994.
- CLARK, B. N.; COLBOURN, C. J.; JOHNSON, D. S. Unit disk graphs. **Discrete Math**, v. 86, n. 1, p. 165-177, 1990.

Received on November 6, 2010.

Accepted on October 14, 2011.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.